


TABLE OF CONTENTS



1/643

Calculus is build on two operations: **differentiation** and **integration**.

Differentiation


- ▶ Slope of a line
- ▶ Rate of change
- ▶ Optimization
- ▶ Numerical Approximations

Integration

- ▶ Area under a curve
- ▶ "Reverse" of differentiation
- ▶ Solving differential equations
- ▶ Calculate net change from rate of change
- ▶ Volume of solids
- ▶ Work (in the physics sense)

2/643

Approximate the area of the shaded region using rectangles.



We're going to be doing a lot of adding.

3/643

SUMMATION (SIGMA) NOTATION

$$\sum_{i=a}^b f(i)$$

- ▶ a, b (integers with $a \leq b$) "bounds"
- ▶ i "index:" integer which runs from a to b
- ▶ $f(i)$ "summands:" compute for every i , add

$$\sum_{i=a}^b f(i) = f(a) + f(a+1) + f(a+2) + \cdots + f(b)$$

4/643

SIGMA NOTATION

Expand $\sum_{i=2}^4 (2i + 5)$.



5/643

SIGMA NOTATION

Expand $\sum_{i=1}^4 (i + (i - 1)^2)$.



6/643

Write the following expressions in sigma notation:

▶ $3 + 4 + 5 + 6 + 7$

▶ $8 + 8 + 8 + 8 + 8$

▶ $1 + (-2) + 4 + (-8) + 16$



7/643

ARITHMETIC OF SUMMATION NOTATION

Let c be a constant.

▶ Adding constants: $\sum_{i=1}^{10} c =$

▶ Factoring constants: $\sum_{i=1}^{10} 5(i^2) =$

▶ Addition is Commutative: $\sum_{i=1}^{10} (i + i^2) =$

8/643

Theorem 1.1.5: Arithmetic of Summation Notation

ARITHMETIC OF SUMMATION NOTATION

Let c be a constant.

► Adding constants: $\sum_{i=1}^{10} c = 10c$

► Factoring constants: $\sum_{i=1}^{10} 5(i^2) = 5 \sum_{i=1}^{10} (i^2)$

► Addition is Commutative: $\sum_{i=1}^{10} (i + i^2) = \left(\sum_{i=1}^{10} i \right) + \left(\sum_{i=1}^{10} i^2 \right)$

COMMON SUMS

Let $n \geq 1$ be an integer, a be a real number, and $r \neq 1$.

$$\sum_{i=0}^n ar^i = a + ar + ar^2 + \cdots + ar^n = a \frac{1 - r^{n+1}}{1 - r}$$

$$\sum_{i=1}^n i = 1 + 2 + \cdots + n = \frac{n(n+1)}{2}$$

$$\sum_{i=1}^n i^2 = 1^2 + 2^2 + \cdots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\sum_{i=1}^n i^3 = 1^3 + 2^3 + \cdots + n^3 = \frac{n^2(n+1)^2}{4}$$

Let $n \geq 1$ be an integer, a be a real number, and $r \neq 1$.

$$\sum_{i=0}^n ar^i = a + ar + ar^2 + \cdots + ar^n = a \frac{1 - r^{n+1}}{1 - r}$$

$$\sum_{i=1}^n i = 1 + 2 + \cdots + n = \frac{n(n+1)}{2}$$

$$\sum_{i=1}^n i^2 = 1^2 + 2^2 + \cdots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\sum_{i=1}^n i^3 = 1^3 + 2^3 + \cdots + n^3 = \frac{n^2(n+1)^2}{4}$$

Simplify: $\sum_{i=1}^{13} (i^2 + i^3)$



Let $n \geq 1$ be an integer, a be a real number, and $r \neq 1$.

$$\sum_{i=0}^n ar^i = a + ar + ar^2 + \cdots + ar^n = a \frac{1 - r^{n+1}}{1 - r}$$

$$\sum_{i=1}^n i = 1 + 2 + \cdots + n = \frac{n(n+1)}{2}$$

$$\sum_{i=1}^n i^2 = 1^2 + 2^2 + \cdots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\sum_{i=1}^n i^3 = 1^3 + 2^3 + \cdots + n^3 = \frac{n^2(n+1)^2}{4}$$

Simplify: $\sum_{i=1}^{50} (1 - i^2)$




(OPTIONAL) PROOF OF A COMMON SUM

Here is a derivation of $\sum_{i=0}^n r^i = \frac{1-r^{n+1}}{1-r}$:

13/643

(OPTIONAL) PROOF OF ANOTHER COMMON SUM


$$\sum_{i=1}^{10} i = 1 + 2 + 3 + 4 + 5 + 6 + 7 + 8 + 9 + 10 =$$



14/643

(OPTIONAL) PROOF OF A COMMON SUM

$$\sum_{i=1}^n i = 1 + 2 + 3 + \dots + n =$$



15/643

The purpose of these sums is to describe areas.

16/643

Notation

The symbol

$$\int_a^b f(x) dx$$

is read “the definite integral of the function $f(x)$ from a to b .”


- ▶ $f(x)$: integrand
- ▶ a and b : limits of integration
- ▶ dx : differential

17/643

If $f(x) \geq 0$ and $a \leq b$, one interpretation of

$$\int_a^b f(x) dx$$

is “the area of the region bounded above by $y = f(x)$, below by $y = 0$, to the left by $x = a$, and to the right by $x = b$.”




18/643

If $f(x) \geq 0$ and $a \leq b$, one interpretation of

$$\int_a^b f(x) dx$$


is the **signed** area of the region between $y = f(x)$ and $y = 0$, from $x = a$ to $x = b$. Area **above** the axis is **positive**, and area **below** it is **negative**.



19/643

RIEMANN SUMS


A **Riemann sum** approximates the area under a curve by cutting it into equal-width segments, and approximating each segment as a rectangle.




There are different ways to choose the height of each rectangle.

20/643


TYPES OF RIEMANN SUMS (RS)



Left RS




Right RS



Midpoint RS


21/643

Approximate $\int_2^4 \log(x) dx$ using a **right Riemann sum** with $n = 4$ rectangles. For now, do not use sigma notation.




22/643

Approximate $\int_{-1}^0 e^x dx$ using a **left Riemann sum** with $n = 3$ rectangles. For now, do not use sigma notation.




23/643

Approximate $\int_0^{\sqrt{\pi}} \sin(x^2) dx$ using a **midpoint Riemann sum** with $n = 5$ rectangles. For now, do not use sigma notation.




24/643

Approximate $\int_1^{17} \sqrt{x} \, dx$ using a **midpoint Riemann sum** with 8 rectangles. Write the result in sigma notation.



$$\sum_{i=1}^8 2\sqrt{2i} = \underbrace{2\sqrt{2}}_{i=1} + \underbrace{2\sqrt{4}}_{i=2} + \underbrace{2\sqrt{6}}_{i=3} + \underbrace{2\sqrt{8}}_{i=4} + \underbrace{2\sqrt{10}}_{i=5} + \underbrace{2\sqrt{12}}_{i=6} + \underbrace{2\sqrt{14}}_{i=7} + \underbrace{2\sqrt{16}}_{i=8}$$




Riemann sum with n rectangles

$$\int_a^b f(x) \, dx \approx \sum_{i=1}^n \Delta x \cdot f(x_{i,n}^*)$$

where $\Delta x = \frac{b-a}{n}$ and $x_{i,n}^*$ is an x -value in the i th rectangle.


$$\sum_{i=1}^n \Delta x \cdot f(x_{i,n}^*) = \Delta x \cdot f(x_{1,n}^*) + \Delta x \cdot f(x_{2,n}^*) + \Delta x \cdot f(x_{3,n}^*) + \cdots + \Delta x \cdot f(x_{n,n}^*)$$



Right Riemann sum with n rectangles


$$\int_a^b f(x) \, dx \approx \sum_{i=1}^n \Delta x \cdot f(\quad)$$

where $\Delta x = \frac{b-a}{n}$ and $x_{i,n}^*$ =



TYPES OF RIEMANN SUMS (RS)

What height would you choose for the i th rectangle?



Right RS

Left RS

Midpoint RS

29/643

Riemann sums with n rectangles. Let $\Delta x = \frac{b-a}{n}$

The **right** Riemann sum approximation of $\int_a^b f(x) dx$ is:

$$\sum_{i=1}^n \Delta x \cdot f(a + i\Delta x)$$

The **left** Riemann sum approximation of $\int_a^b f(x) dx$ is:

$$\sum_{i=1}^n \Delta x \cdot f(a + (i-1)\Delta x)$$

The **midpoint** Riemann sum approximation of $\int_a^b f(x) dx$ is:

$$\sum_{i=1}^n \Delta x \cdot f\left(a + \left(i - \frac{1}{2}\right) \Delta x\right)$$

30/643 Definition 1.1.11

Riemann sums with n rectangles: Let $\Delta x = \frac{b-a}{n}$

The **right** Riemann sum approximation of $\int_a^b f(x) dx$ is:

$$\sum_{i=1}^n \Delta x \cdot f(a + i\Delta x)$$

Give a right Riemann Sum for the area under the curve $y = x^2 - x$ from $a = 1$ to $b = 6$ using $n = 1000$ intervals.



31/643

Riemann sums with n rectangles: Let $\Delta x = \frac{b-a}{n}$

The **midpoint** Riemann sum approximation of $\int_a^b f(x) dx$ is:

$$\sum_{i=1}^n \Delta x \cdot f\left(a + \left(i - \frac{1}{2}\right) \Delta x\right)$$

Give a midpoint Riemann Sum for the area under the curve $y = 5x - x^2$ from $a = 6$ to $b = 9$ using $n = 1000$ intervals.



32/643

EVALUATING RIEMANN SUMS

▶ SKIP RIEMANN EVALUATIONS

$$\sum_{i=1}^n i = \frac{n(n+1)}{2}$$

$$\sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\sum_{i=1}^n i^3 = \frac{n^2(n+1)^2}{4}$$

Give the right Riemann sum of $f(x) = x^2$ from $a = 0$ to $b = 10$, $n = 100$:

$$\sum_{i=1}^n \Delta x \cdot f(a + i\Delta x) =$$



33/643

EVALUATING RIEMANN SUMS IN SIGMA NOTATION

$$\sum_{i=1}^n i = \frac{n(n+1)}{2}$$


$$\sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\sum_{i=1}^n i^3 = \frac{n^2(n+1)^2}{4}$$


Give the right Riemann sum of $f(x) = x^3$ from $a = 0$ to $b = 5$, $n = 100$:



35/643



34/643



36/643

Definition

Let a and b be two real numbers and let $f(x)$ be a function that is defined for all x between a and b . Then we define $\Delta x = \frac{b-a}{N}$ and


$$\int_a^b f(x) \, dx = \lim_{N \rightarrow \infty} \sum_{i=1}^N f(x_{i,N}^*) \cdot \Delta x$$

when the limit exists and when the choice of $x_{i,N}^*$ in the i^{th} interval doesn't matter.

\sum, \int both stand for "sum"

$\Delta x, dx$ are tiny pieces of the x -axis, the bases of our very skinny rectangles

It's understood we're taking a limit as N goes to infinity, so we don't bother specifying N (or each location where we find our height) in the second notation.



$$\sum_{i=1}^n i = \frac{n(n+1)}{2}$$

$$\sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$$


$$\sum_{i=1}^n i^3 = \frac{n^2(n+1)^2}{4}$$

Give the right Riemann sum of $y = x^2$ from $a = 0$ to $b = 5$ with n slices, and simplify:

We found the right Riemann sum of $y = x^2$ from $a = 0$ to $b = 5$ using n slices:

$$\frac{125}{6} \cdot \frac{2n^2 + 3n + 1}{n^2}$$

Use it to find the exact area under the curve.



REFRESHER: LIMITS OF RATIONAL FUNCTIONS

$$\lim_{n \rightarrow \infty} \frac{n^2 + 2n + 15}{3n^2 - 9n + 5} =$$

When the degree of the top and bottom are the same, the limit as n goes to infinity is the ratio of the leading coefficients.

$$\lim_{n \rightarrow \infty} \frac{n^2 + 2n + 15}{3n^3 - 9n + 5} =$$

When the degree of the top is smaller than the degree of the bottom, the limit as n goes to infinity is 0.

$$\lim_{n \rightarrow \infty} \frac{n^3 + 2n + 15}{3n^2 - 9n + 5} =$$

When the degree of the top is larger than the degree of the bottom, the limit as n goes to infinity is positive or negative infinity.

41/643

$$\sum_{i=1}^n i = \frac{n(n+1)}{2}$$

$$\sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\sum_{i=1}^n i^3 = \frac{n^2(n+1)^2}{4}$$


Evaluate $\int_0^1 x^2 dx$ exactly using midpoint Riemann sums.

42/643

Let's see some special cases where we can use geometry to evaluate integrals without Riemann sums.

43/643


$$\int_0^5 2x dx$$



44/643




$$\int_3^5 2x \, dx$$




45/643

$$\int_{-2}^2 \sin x \, dx$$




46/643

$$\int_{-1}^1 |x| \, dx$$




47/643

$$\int_0^1 \sqrt{1-x^2} \, dx$$



48/643


$$\int_{10}^{10} \log x \, dx$$



A car travelling down a straight highway records the following measurements:

Time	12:00	12:10	12:20	12:30	12:40	12:50	1:00
Speed (kph)	80	100	100	90	90	75	100

Approximately how far did the car travel from 12:00 to 1:00?



The computation

$$\text{distance} = \text{rate} \times \text{time}$$

looks a lot like the computation

$$\text{area} = \text{base} \times \text{height}$$

for a rectangle. This gives us another interpretation for an integral.


ANOTHER INTERPRETATION OF THE INTEGRAL

Let $x(t)$ be the position of an object moving along the x -axis at time t , and let $v(t) = x'(t)$ be its velocity. Then for all $b > a$,

$$x(b) - x(a) = \int_a^b v(t) \, dt$$

That is, $\int_a^b v(t) \, dt$ gives the *net distance* moved by the object from time a to time b .

TABLE OF CONTENTS



We defined the definite integral as

$$\int_a^b f(x) dx = \lim_{N \rightarrow \infty} \sum_{i=1}^N \Delta x \cdot f(x_{i,N}^*)$$

where $\Delta x = \frac{b-a}{N}$ and $x_{i,N}^*$ is a point in the interval $[a + (i-1)\Delta x, a + i\Delta x]$.

We have seen in previous classes that limits don't always exist. We will verify that this limit does indeed exist, and is equal to the desired area (at least in the most common cases).

We'll start with some general ideas that appear in the proof.

Suppose x and y are both in the interval $[a, b]$. What is the maximum possible value of $|x - y|$?




Proposition 1: distance between two numbers in an interval


If $a \leq x \leq b$ and $a \leq y \leq b$, then

Proposition 2: area inequality


Let $f(x)$ be a function, defined over the interval $[a, b]$. If $m \leq f(x) \leq M$ over the entire interval $[a, b]$, then the (signed) area between the curve $y = f(x)$ and the x -axis, from a to b , is between $m(b - a)$ and $M(b - a)$.



Intuition: If $f'(x)$ is bounded on (a, b) and $b - a$ is small, then $f(b) - f(a)$ is also small.



The Mean Value Theorem provides a more explicit connection between these quantities.



Mean Value Theorem

Let a and b be real numbers with $a < b$. Let f be a function such that

- ▶ $f(x)$ is continuous on the closed interval $a \leq x \leq b$, and
- ▶ $f(x)$ is differentiable on the open interval $a < x < b$.

Then there is a c in (a, b) such that

$$f(b) - f(a) = f'(c)(b - a).$$

Equivalently: $f'(c) = \frac{f(b) - f(a)}{b - a}$.

Triangle Inequality

For any real numbers x_1, x_2, \dots, x_n :

$$\left| \sum_{i=1}^n x_i \right| \leq \sum_{i=1}^n |x_i|$$

Intuition: If some terms are positive and some are negative, they “cancel each other out” and make the overall sum smaller.

$ 1 + 2 $	$ 1 + 2 $
$ 1 + (-2) $	$ 1 + -2 $
$ (-1) + (-2) $	$ -1 + -2 $

Triangle Inequality

For any real numbers x_1, x_2, \dots, x_n :

$$\left| \sum_{i=1}^n x_i \right| \leq \sum_{i=1}^n |x_i|$$

Proof outline:

REQUIREMENTS

We will consider

$$\int_a^b f(x) \, dx$$


where:

- ▶ $a < b$
- ▶ $f(x)$ is continuous over the interval $[a, b]$
- ▶ $f(x)$ is differentiable over the interval (a, b)
- ▶ $f'(x)$ is bounded over the interval (a, b) . That is, there exists a positive constant number F such that $|f'(x)| \leq F$ for all x in the interval (a, b) .

61/643

ERROR IN A SINGLE SLICE

Consider approximating the area of single slice, from x_0 to x_1 .




- ▶ A is the actual area of the slice
- ▶ $f(\bar{x})$ and $f(\underline{x})$ are the largest and smallest function values over the slice
- ▶ Our slice has width $x_1 - x_0$

Then we can bound our area:

62/643 Proposition 2: area inequality

ERROR IN A SINGLE SLICE

Consider approximating the area of single slice, from x_0 to x_1 .




- ▶ $f(x^*) \cdot (x_1 - x_0)$ is our approximation of the area of the slice, for some x^* in the interval $[x_0, x_1]$.
- ▶ $f(\bar{x})$ and $f(\underline{x})$ are the largest and smallest function values over the slice, so

$$f(\underline{x}) \leq f(x^*) \leq f(\bar{x})$$

Then we can bound our approximation:

63/643 using Proposition 2: area inequality

ERROR IN A SINGLE SLICE




$$\begin{aligned} f(\underline{x}) \cdot (x_1 - x_0) &\leq A &\leq f(\bar{x}) \cdot (x_1 - x_0) \\ f(\underline{x}) \cdot (x_1 - x_0) &\leq f(x^*) \cdot (x_1 - x_0) &\leq f(\bar{x}) \cdot (x_1 - x_0) \end{aligned}$$

$$\underbrace{|A - f(x^*) \cdot (x_1 - x_0)|}_{\text{error in slice}} \leq$$

64/643 using Proposition 1: difference between two numbers in a given interval

ERROR IN A SINGLE SLICE

- ▶ The error in our single slice is at most $[f(\bar{x}) - f(\underline{x})] \cdot (x_1 - x_0)$
- ▶ We want to show that our total error is not too large.
- ▶ Intuitively, if $|f'(x)|$ is never very large, and $x_1 - x_0$ is not very large, then $f(\bar{x}) - f(\underline{x})$ is not very large.



65/643

ERROR IN A SINGLE SLICE

Mean Value Theorem

Let a and b be real numbers with $a < b$. Let f be a function such that

- ▶ $f(x)$ is continuous on the closed interval $a \leq x \leq b$, and
- ▶ $f(x)$ is differentiable on the open interval $a < x < b$.

Then there is a c in (a, b) such that

$$f(b) - f(a) = f'(c)(b - a)$$

There exists some c in (x_0, x_1) such that

$$f(\bar{x}) - f(\underline{x}) = f'(c) \cdot (\bar{x} - \underline{x})$$

Since $|f'(x)|$ is never larger than the positive constant F in (a, b) ,

$$|f(\bar{x}) - f(\underline{x})| \leq F \cdot |\bar{x} - \underline{x}| \leq F \cdot |x_1 - x_0|$$

66/643 CLP1 Theorem 2.13.4, the mean value theorem, and Proposition 1

ERROR IN A SINGLE SLICE

All together,

$$\underbrace{|A - f(x^*) \cdot (x_1 - x_0)|}_{\text{error in slice}} \leq$$


67/643

We have shown that the error on a **single** slice can't be worse than some amount.

Now let's consider adding up slices.

68/643

What we did for a single slice, we now do for all slices.
Updated notation for slice j :



Slice error bound:


$$|A_j - f(x_j^*) \cdot (x_j - x_{j-1})| \leq F \cdot (x_j - x_{j-1})^2$$

69/643

(POSSIBLY IRREGULAR) PARTITIONS

Consider partitioning the interval $[a, b]$ into n subintervals, not necessarily the same size. Let the points at the edges of the slices be $a = x_0, x_1, x_2, \dots, x_{n-1}, x_n = b$.

In each part, choose a vertex x_i^* to sample the height of the function.



The approximation of $\int_a^b f(x) dx$ depends on how you choose your subintervals, and where you choose your sample points. Let


$$\mathbb{P} = (n, x_1, x_2, \dots, x_{n-1}, x_1^*, x_2^*, \dots, x_n^*)$$

denote these choices.

70/643

Let $\mathcal{I}(\mathbb{P})$ be the approximation that arises from \mathbb{P} :

$$\mathcal{I}(\mathbb{P}) = \sum_{i=1}^n f(x_i^*) (x_i - x_{i-1})$$



Let $M(\mathbb{P})$ be the maximum width of any subinterval.
If $M(\mathbb{P})$ is small, then *every* subinterval is small (narrow).

71/643


Define the integral as the limit

$$\int_a^b f(x) dx = \lim_{M(\mathbb{P}) \rightarrow 0} \mathcal{I}(\mathbb{P})$$


(Compare to our previous Riemann sum definition.)

We will show that the limit exists and is equal to the signed area under the curve.

72/643



Actual area: $\int_a^b f(x) dx = \sum_{i=1}^n A_i$



Approximation: $\mathcal{I}(\mathbb{P}) = \sum_{i=1}^n f(x_i^*) \cdot (x_i - x_{i-1})$

$$\underbrace{\left| \int_a^b f(x) dx - \mathcal{I}(\mathbb{P}) \right|}_{\text{overall error}} = \left| \sum_{i=1}^n A_i - \sum_{i=1}^n f(x_i^*) \cdot (x_i - x_{i-1}) \right|$$

$$= \left| \sum_{i=1}^n [A_i - f(x_i^*) \cdot (x_i - x_{i-1})] \right|$$

$$0 \leq \underbrace{\left| \int_a^b f(x) dx - \mathcal{I}(\mathbb{P}) \right|}_{\text{overall error}} \leq F \cdot M(\mathbb{P}) \cdot (b - a)$$

$$\lim_{M(\mathbb{P}) \rightarrow 0} 0 = 0 \qquad \lim_{M(\mathbb{P}) \rightarrow 0} [F \cdot M(\mathbb{P}) \cdot (b - a)] = 0$$

So, by the squeeze theorem,

$$\lim_{M(\mathbb{P}) \rightarrow 0} \underbrace{\left| \int_a^b f(x) dx - \mathcal{I}(\mathbb{P}) \right|}_{\text{overall error}} = 0$$

That is,

$$\lim_{M(\mathbb{P}) \rightarrow 0} \mathcal{I}(\mathbb{P}) = \int_a^b f(x) dx$$

COMPARING DEFINITIONS

Here, we defined

$$\int_a^b f(x) dx = \lim_{M(\mathbb{P}) \rightarrow 0} \mathcal{I}(\mathbb{P})$$

for “nice” functions $f(x)$.

Originally, we used a slightly different definition:

Definition 1.1.9 (abridged)

For “nice” functions $f(x)$:

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_{i,n}^*) \cdot \frac{b-a}{n}$$

when the limit exists and takes the same value for all choices of the $x_{i,n}^*$'s.


COMPARING DEFINITIONS

We showed that **all** families of partitions “work,” as long as their largest subintervals shrink to length 0.

If all families of partitions “work,” then we might as well choose a simple one. The (arguably) simplest choices are regular partitions, cutting the interval $[a, b]$ into n subintervals of length $\frac{b-a}{n}$.

77/643

TABLE OF CONTENTS



78/643

We defined the definite integral using a limit and a sum.

Definition

Let a and b be two real numbers and let $f(x)$ be a function that is defined for all x between a and b . Then we define $\Delta x = \frac{b-a}{N}$ and


$$\int_a^b f(x) \, dx = \lim_{N \rightarrow \infty} \sum_{i=1}^N f(x_{i,N}^*) \cdot \Delta x$$

when the limit exists and when the choice of $x_{i,N}^*$ in the i^{th} interval doesn't matter.

Many of the operations that work nicely with sums and limits will also work nicely with integrals.


79/643

ADDING (AND SUBTRACTING) FUNCTIONS




80/643

ADDING (AND SUBTRACTING) FUNCTIONS



81/643


ADDING (AND SUBTRACTING) FUNCTIONS



$$\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$$

82/643


ADDING (AND SUBTRACTING) FUNCTIONS



$$\int_a^b (f(x) - g(x)) dx = \int_a^b f(x) dx - \int_a^b g(x) dx$$


83/643

MULTIPLYING A FUNCTION BY A CONSTANT



84/643

MULTIPLYING A FUNCTION BY A CONSTANT



85/643

ARITHMETIC OF INTEGRATION

When a , b , and c are real numbers, and the functions $f(x)$ and $g(x)$ are integrable on an interval containing a and b :

$$(a) \int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx$$


$$(b) \int_a^b [f(x) - g(x)] dx = \int_a^b f(x) dx - \int_a^b g(x) dx$$

$$(c) \int_a^b c \cdot f(x) dx = c \int_a^b f(x) dx \quad \text{when } c \text{ is constant}$$

86/643 Theorem 1.2.1: Arithmetic of Integration

ARITHMETIC OF INTEGRATION

Suppose $\int_{-1}^1 f(x) dx = -6$ and $\int_{-1}^1 g(x) dx = 10$.




$$\int_{-1}^1 (2f(x) + g(x)) dx =$$

|| ∞


87/643

INTERVAL OF INTEGRATION



88/643


INTERVAL OF INTEGRATION



What rule do you think is being illustrated?

89/643

WHAT HAPPENS IN $\int_a^b f(x) dx$ WHEN $b < a$?



$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_{i,n}^*) \cdot \frac{b-a}{n}$$

This is the definition of a definite integral *whether or not* $a < b$.

90/643

PROPERTY OF DEFINITE INTEGRALS

$$\int_a^b f(x) dx = - \int_b^a f(x) dx$$


As strictly a measure of area, not usually a super useful fact – but helps later when we do arithmetic with integrals.

It's also useful that the definition works without having to worry about which limit of integration (a or b) is larger.

91/643


ARITHMETIC FOR DOMAIN OF INTEGRATION

When a , b , and c are constants, and $f(x)$ is integrable over a domain containing all three:

(a) $\int_a^a f(x) dx = 0$ 


(b) $\int_a^b f(x) dx = - \int_b^a f(x) dx$ $\Delta x = \frac{b-a}{n} = - \frac{a-b}{n}$

(c) $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$ for constant c



92/643 Theorem 1.2.3: Arithmetic for the Domain of Integration

Suppose $\int_{-1}^0 f(x) dx = 1$, $\int_0^1 f(x) dx = -3$, and $\int_{-1}^1 g(x) dx = 10$.




$$\int_{-1}^1 (2f(x) + g(x)) dx =$$

93/643



Suppose $\int_{-1}^0 f(x) dx = 1$ and $\int_0^1 f(x) dx = -3$.



$$\int_{-1}^3 f(x) dx + \int_3^0 f(x) dx =$$


94/643



Even and Odd Functions


Let $f(x)$ be a function.

- ▶ We say $f(x)$ is **even** when $f(x) = f(-x)$ for all x , and
- ▶ we say $f(x)$ is **odd** when $f(x) = -f(-x)$ for all x .



95/643 Definition 1.2.9 in CLP-2; Definition 3.6.6 and 3.6.7 in CLP-1

INTEGRALS OF EVEN FUNCTIONS




Suppose $f(x)$ is **even**. Then

$$\int_a^b f(x) dx =$$

96/643

INTEGRALS OF EVEN FUNCTIONS




Suppose $f(x)$ is **even**. Then

$$\int_{-b}^b f(x) \, dx =$$

97/643

INTEGRALS OF ODD FUNCTIONS



Suppose $f(x)$ is **odd**. Then

$$\int_{-b}^b f(x) \, dx =$$

98/643

Theorem 1.2.12 (Even and Odd)

Let $a > 0$.

(a) If $f(x)$ is an **even** function, then

$$\int_{-a}^a f(x) \, dx = 2 \int_0^a f(x) \, dx$$

(b) If $f(x)$ is an **odd** function, then

$$\int_{-a}^a f(x) \, dx = 0$$


99/643

Integral Inequality

Let $a \leq b$ be real numbers and let the functions $f(x)$ and $g(x)$ be integrable on the interval $a \leq x \leq b$.

If $f(x) \leq g(x)$ for all $a \leq x \leq b$, then

$$\int_a^b f(x) \, dx \leq \int_a^b g(x) \, dx$$




100/643 Theorem 1.2.13

Integral Inequality

Let $a \leq b$ and $m \leq M$ be real numbers and let the function $f(x)$ be integrable on the interval $a \leq x \leq b$.


If $m \leq f(x) \leq M$ for all $a \leq x \leq b$, then

$$m(b-a) \leq \int_a^b f(x) dx \leq M(b-a)$$




Find a lower bound c and an upper bound d such that

$$c \leq \int_1^5 f(x) dx \leq d$$



Find a lower bound c and an upper bound d such that $d - c \leq 3$ and


$$c \leq \int_0^6 f(x) dx \leq d$$



ABSOLUTE VALUES

$$f(x) \leq |f(x)| \text{ for any } f(x)$$


$$-f(x) \leq |f(x)| \text{ for any } f(x)$$



$$\left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx$$

because $\left| \int_a^b f(x) dx \right|$ is either $\int_a^b f(x) dx$ or $-\int_a^b f(x) dx$.

TABLE OF CONTENTS



105/643

REVIEW: AREA UNDER A CURVE

Methods for finding the area under a curve.

► Limit of a Riemann Sum

► **Conceptually** easy – cut into rectangles


► **Computationally** rough $\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x; \sum_{i=1}^n i = \frac{n(n+1)}{2}$



► Use Geometry

► **Computationally** nice when it's available! (Circles, triangles, symmetry, etc.)

► Often not available – most functions don't make such nice shapes.



► Up next: Fundamental Theorem of Calculus

► **Conceptually** less obvious – we'll spend about a day explaining **why** it works

► **Computationally** generally nicer than Riemann sums

► Doesn't work for every function

106/643

Fundamental Theorem of Calculus, Part 1

Let $a < b$ and let $f(x)$ be a function which is defined and continuous on $[a, b]$. Let

$$A(x) = \int_a^x f(t) dt$$

for any x in $[a, b]$. Then the function $A(x)$ is differentiable and


$$A'(x) = f(x).$$

FTC(I) gives us the derivative of a very specific function (subject to some fine print).

It shows a close relationship between integrals and derivatives.


107/643 Theorem 1.3.1

AREA FUNCTION: $A(x) = \int_a^x f(t) dt$ FOR $a \leq x \leq b$



108/643

AREA FUNCTION: $A(x) = \int_a^x f(t) dt$ FOR $a \leq x \leq b$



Notation: the function A depends on the variable x .

We need to know how the function f behaves on the whole interval $(0, x)$ to find $A(x)$. That's why we use $f(t)$, not $f(x)$.



109/643

AREA FUNCTION NOTATION

It might look strange at first to see two different variables. Let's consider the alternatives:

$$A(x) = \int_0^x f(t) dt \quad B(x) = \int_0^x f(x) dt \quad C(x) = \int_0^x f(x) dx$$

$$A(1) = \int_0^1 f(t) dt \quad B(1) = \int_0^1 f(1) dt \quad C(1) = \int_0^1 f(1) \underbrace{d1}_{??}$$



110/643

Fundamental Theorem of Calculus, Part 1

Let $a < b$ and let $f(x)$ be a function which is defined and continuous on $[a, b]$. Let

$$A(x) = \int_a^x f(t) dt$$


for any x in $[a, b]$. Then the function $A(x)$ is differentiable and

$$A'(x) = f(x).$$

Question: Why is it true?

111/643 Theorem 1.3.1


DERIVATIVE OF AREA FUNCTION, $A(x) = \int_a^x f(t) dt$



$$A'(x) = \lim_{\Delta x \rightarrow 0} \frac{\Delta A}{\Delta x} = \lim_{h \rightarrow 0} \frac{A(x+h) - A(x)}{h} = \lim_{h \rightarrow 0} \frac{hf(x)}{h} = f(x)$$

112/643

DERIVATIVE OF AREA FUNCTION, $A(x) = \int_a^x f(t) dt$



$$A'(x) = \lim_{\Delta x \rightarrow 0} \frac{\Delta A}{\Delta x} = \lim_{h \rightarrow 0} \frac{A(x+h) - A(x)}{h} = \lim_{h \rightarrow 0} \frac{hf(x)}{h} = f(x)$$

When h is very small, the purple area looks like a rectangle with base h and height $f(x)$, so $A(x+h) - A(x) \approx hf(x)$ and $\frac{A(x+h) - A(x)}{h} \approx f(x)$. As h tends to zero, the error in this approximation approaches 0.

113/643

Fundamental Theorem of Calculus, Part 1

Let $a < b$ and let $f(x)$ be a function which is defined and continuous on $[a, b]$. Let

$$A(x) = \int_a^x f(t) dt$$

for any x in $[a, b]$. Then the function $A(x)$ is differentiable and

$$A'(x) = f(x).$$

Suppose $A(x) = \int_2^x \sin t dt$. What is $A'(x)$?

Suppose $B(x) = \int_x^2 \sin t dt$. What is $B'(x)$?

114/643 Theorem 1.3.1



Fundamental Theorem of Calculus, Part 1

Let $a < b$ and let $f(x)$ be a function which is defined and continuous on $[a, b]$. Let

$$A(x) = \int_a^x f(t) dt$$

for any x in $[a, b]$. Then the function $A(x)$ is differentiable and

$$A'(x) = f(x).$$

Suppose $C(x) = \int_2^{e^x} \sin t dt$. What is $C'(x)$?




115/643 Theorem 1.3.1


It's possible to have two different functions with the same derivative.

$$A(x) = \int_0^x 2t dt = x^2$$

$$B(x) = \int_1^x 2t dt = x^2 - 1$$



$$A'(x) = 2x$$




$$B'(x) = 2x$$

When two functions have the same derivative, they differ only by a constant.

In this example: $B(x) = A(x) - 1$

116/643 Lemma 1.3.8



If two continuous functions have the same derivative, then one is a constant plus the other.

Two clues for finding $A(x) = \int_a^x f(t) dt$:

- ▶ If $A(x) = \int_a^x f(t) dt$, then¹ $A'(x) = f(x)$
- ▶ If $F'(x) = A'(x)$, then $A(x) = F(x) + C$ for some constant C .

$A(x) = \int_a^x e^t dt$. What functions could $A(x)$ be?

¹(as long as $f(t)$ is continuous on $[a, x]$)



Two clues for finding $A(x) = \int_a^x f(t) dt$:

- ▶ If $A(x) = \int_a^x f(t) dt$, then¹ $A'(x) = f(x)$
- ▶ If $F'(x) = A'(x)$, then $A(x) = F(x) + C$ for some constant C .

$A(x) = \int_a^x \cos t dt$. What functions could $A(x)$ be?

¹(as long as $f(t)$ is continuous on $[a, x]$)



Two clues for finding $A(x) = \int_a^x f(t) dt$:


- ▶ If $A(x) = \int_a^x f(t) dt$, then¹ $A'(x) = f(x)$
- ▶ If $F'(x) = A'(x)$, then $A(x) = F(x) + C$ for some constant C .

$A(x) = \int_{-2}^x 5t^4 dt$. What functions could $A(x)$ be?

¹(as long as $f(t)$ is continuous on $[a, x]$)




$$A(x) = \int_{-2}^x 5t^4 dt = x^5 + 32$$



121/643


$$A(x) = \int_{-2}^x 5t^4 dt = x^5 + 32$$



$$A(-1) = \int_{-2}^{-1} 5t^4 dt = (-1)^5 + 32 = 31$$

122/643


$$A(x) = \int_{-2}^x 5t^4 dt = x^5 + 32$$



$$A(0) = \int_{-2}^0 5t^4 dt = (0)^5 + 32 = 32$$

123/643


$$A(x) = \int_{-2}^x 5t^4 dt = x^5 + 32$$



$$A(1) = \int_{-2}^1 5t^4 dt = (1)^5 + 32 = 33$$

124/643


$$A(x) = \int_{-2}^x 5t^4 dt = x^5 + 32$$



$$A(2) = \int_{-2}^2 5t^4 dt = (2)^5 + 32 = 64$$

125/643

$$A(x) = \int_{-2}^x 5t^4 dt = x^5 + 32$$



$$A(3) = \int_{-2}^3 5t^4 dt = (3)^5 + 32 = 275$$

126/643

Two clues for finding $A(x) = \int_a^x f(t) dt$:

- ▶ If $A(x) = \int_a^x f(t) dt$, then¹ $A'(x) = f(x)$
- ▶ If $F'(x) = A'(x)$, then $A(x) = F(x) + C$ for some constant C .

$A(x) = \int_a^x f(t) dt$. What functions could $A(x)$ be?

¹(as long as $f(t)$ is continuous on $[a, x]$)

127/643

Two clues for finding $A(x) = \int_a^x f(t) dt$:

- ▶ If $A(x) = \int_a^x f(t) dt$, then¹ $A'(x) = f(x)$
- ▶ If $F'(x) = A'(x)$, then $A(x) = F(x) + C$ for some constant C .


$A(b) = \int_a^b f(t) dt$. What functions could $A(b)$ be?

- ▶ $A'(x) = f(x)$.
- ▶ Guess a function with derivative $f(x)$: $F(x)$.
- ▶ Then $A(x) = F(x) + C$ for some constant C .
- ▶ Also $A(a) = 0$, so $0 = F(a) + C$, so $C = -F(a)$
- ▶ So, $A(b) = F(b) - F(a)$

¹(as long as $f(t)$ is continuous on $[a, x]$)


128/643

$$\int_a^b f(t) dt = F(b) - F(a) \quad \text{where} \quad F'(x) = f(x)$$



129/643


$$\int_a^b f(t) dt = F(b) - F(a) \quad \text{where} \quad F'(x) = f(x)$$



$$\int_{-2}^1 f(t) dt = F(1) - F(-2)$$

130/643


$$\int_a^b f(t) dt = F(b) - F(a) \quad \text{where} \quad F'(x) = f(x)$$



$$\int_{-2}^2 f(t) dt = F(2) - F(-2)$$

131/643

$$\int_a^b f(t) dt = F(b) - F(a) \quad \text{where} \quad F'(x) = f(x)$$



$$\int_{-3}^3 f(t) dt = F(3) - F(-3)$$

132/643

Fundamental Theorem of Calculus, Part 2

Let $F(x)$ be differentiable, defined, and continuous on the interval $[a, b]$ with $F'(x) = f(x)$ for all $a < x < b$. Then

$$\int_a^b f(x) dx = F(b) - F(a)$$

Examples:


$$\frac{d}{dx} \{5x^7\} = 35x^6, \text{ so}$$

$$\int_0^3 35x^6 dx =$$

$$\frac{d}{dx} \{\tan x\} = \sec^2 x, \text{ so}$$


$$\int_0^{\pi/4} \sec^2 x dx =$$

$$\int_0^3 35x^6 dx = F(b) - F(a) \quad \text{where} \quad F(x) = 5x^7$$



$$\int_0^3 35x^6 dx = 5(3)^7 - 5(0)^7$$

$$\int_0^{\pi/4} \sec^2 x dx = F(b) - F(a) \quad \text{where} \quad F(x) = \tan x$$



$$\int_0^{\pi/4} \sec^2 x dx = \tan\left(\frac{\pi}{4}\right) - \tan 0 = 1$$

RELEVANT VOCABULARY

Definition

If $F(x)$ is a function whose derivative is $f(x)$, we call $F(x)$ an **antiderivative** of $f(x)$.

Examples:

The derivative of x^2 is $2x$, so:
 x^2 is an **antiderivative** of $2x$.

When $x > 0$, the derivative of $\log x$ is $\frac{1}{x}$, so:

For all x , the derivative of $\log |x|$ is $\frac{1}{x}$, so:

An antiderivative of $\sin x$ is



CONVENIENT NOTATION

Definition

$$f(x) \Big|_a^b = f(b) - f(a)$$

The function $f(x)$ evaluated from a to b

Examples:

$$(5x + x^2) \Big|_1^2 =$$

$$\frac{x^2}{x+2} \Big|_5^{-1} =$$

FTC Part 2, Abridged

$$\int_a^b f(x) dx = F(x) \Big|_a^b$$

where $F(x)$ is an antiderivative of $f(x)$

137/643

Definition

The **indefinite integral** of a function $f(x)$:

$$\int f(x) dx$$

means the *most general* antiderivative of $f(x)$.

Examples:

$$\int 2x dx =$$

$$\int \frac{1}{x} dx =$$

Remember: two functions with the same derivative differ by a constant, so we include the “+C” for indefinite integrals.

138/643

DEFINITE VS INDEFINITE INTEGRALS

For each pair of properties below, decide which applies to **definite** integrals, and which to **indefinite** integrals.

No limits (or bounds) of integration, $\int f(x) dx$	
Limits (or bounds) of integration, $\int_a^b f(x) dx$	
Area under a curve	
Antiderivative	
Number	
Function	

139/643

ANTIDIFFERENTIATION BY INSPECTION

1. $\int e^x dx$
2. $\int \cos x dx$
3. $\int -\sin x dx$
4. $\int \frac{1}{x} dx$
5. $\int 1 dx$
6. $\int 2x dx$
7. $\int nx^{n-1} dx$ ($n \neq 0$, constant)
8. $\int x^n dx$ ($n \neq -1$, constant)

140/643

Power Rule for Antidifferentiation

$$\int x^n dx = \frac{1}{n+1}x^{n+1} + C$$

if $n \neq -1$ is a constant

Example:


$$\int (5x^2 - 15x + 3) dx =$$



ANTIDERIVATIVES TO RECOGNIZE

- ▶ $\int x^n dx = \frac{1}{n+1}x^{n+1} + C$
- ▶ $\int a dx = ax + C$
- ▶ $\int e^x dx = e^x + C$
- ▶ $\int \frac{1}{x} dx = \log|x| + C$
- ▶ $\int \sin x dx = -\cos x + C$
- ▶ $\int \cos x dx = \sin x + C$
- ▶ $\int \sec^2 x dx = \tan x + C$
- ▶ $\int \sec x \tan x dx = \sec x + C$
- ▶ $\int \csc x \cot x dx = -\csc x + C$
- ▶ $\int \csc^2 x dx = -\cot x + C$
- ▶ $\int \frac{1}{1+x^2} dx = \arctan x + C$
- ▶ $\int \frac{1}{\sqrt{1-x^2}} dx = \arcsin x + C$

TABLE OF CONTENTS



ANTIDERIVATIVES

Fact:

$$\frac{d}{dx} \{\sin(x^2 + x)\} =$$

Related Fact:

$$\int (2x + 1) \cos(x^2 + x) dx =$$

ANTIDERIVATIVES

Chain Rule:

$$\frac{d}{dx} \left\{ \underbrace{\sin(x^2 + x)}_{\text{inside function}} \right\} = \underbrace{(2x + 1)}_{\substack{\text{derivative of} \\ \text{inside function}}} \cos(\underbrace{x^2 + x}_{\text{inside function}})$$

Hallmark of the chain rule: an “inside” function, with that function’s derivative multiplied.

145/643

SOLVE BY INSPECTION

$$\int 2xe^{x^2+1} dx$$

$$\int \frac{1}{x} \cos(\log x) dx$$

$$\int 3(\sin x + 1)^2 \cos x dx$$

(Look for an “inside” function, with its derivative multiplied.)

146/643

UNDOING THE CHAIN RULE

Chain Rule:

$$\frac{d}{dx} \{f(u(x))\} = f'(u(x)) \cdot u'(x)$$

(Here, $u(x)$ is our “inside function”)

Antiderivative Fact:

$$\int f'(u(x)) \cdot u'(x) dx = f(u(x)) + C$$

147/643

UNDOING THE CHAIN RULE

Antiderivative Fact:

$$\int f'(u(x)) \cdot u'(x) dx = f(u(x)) + C$$

Shorthand: call $u(x)$ simply u ;
since $\frac{du}{dx} = u'(x)$, call $u'(x) dx$ simply du .

$$\int f'(u(x)) \cdot u'(x) dx = \int f'(u) du \Big|_{u=u(x)} = f(u(x)) + C$$

This is the **substitution rule**.

148/643

We saw these integrals before, and solved them by inspection. Now try using the language of substitution.

$$\int 2xe^{x^2+1} dx$$

$$\int \frac{1}{x} \cos(\log x) dx$$

$$\int 3(\sin x + 1)^2 \cos x dx$$

149/643

$$\int (3x^2) \sin(x^3 + 1) dx =$$

150/643

$$\int (3x^2) \sin(x^3 + 1) dx = \int \sin(u) du \Big|_{u=x^3+1}$$

“Inside” function: $x^3 + 1$. Its derivative: $3x^2$
Shorthand: $x^3 + 1 \rightarrow u$, $3x^2 dx \rightarrow du$

151/643

$$\begin{aligned} \int (3x^2) \sin(x^3 + 1) dx &= \int \sin(u) du \Big|_{u=x^3+1} \\ &= -\cos(u) + C \Big|_{u=x^3+1} \\ &= \cos(x^3 + 1) + C \end{aligned}$$

“Inside” function: $x^3 + 1$. Its derivative: $3x^2$
Shorthand: $x^3 + 1 \rightarrow u$, $3x^2 dx \rightarrow du$

Warning 1: We don't just change dx to du . We need to couple dx with the derivative of our inside function.
After all, we're undoing the chain rule! We need to have an “inside derivative.”

Warning 2: The final answer is a function of x .

152/643

We used the substitution rule to conclude

$$\int (3x^2) \sin(x^3 + 1) dx = -\cos(x^3 + 1) + C$$

We can check that our antiderivative is correct by differentiating.

153/643

We saw:

$$\int 3x^2 \sin(x^3 + 1) dx = -\cos(x^3 + 1) + C$$

So, we can evaluate:

$$\int_0^1 3x^2 \sin(x^3 + 1) dx$$

154/643

NOTATION: LIMITS OF INTEGRATION

$$\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin^3 x} dx$$

Let $u = \sin x$, $du = \cos x dx$. Note the limits (or bounds) of integration $\pi/4$ and $\pi/2$ are values of x , not u : they follow the differential, unless otherwise specified.

$$\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin^3 x} dx$$

$x = \frac{\pi}{2}$
 $x = \frac{\pi}{4}$

155/643

TRUE OR FALSE?

1. Using $u = x^2$,

$$\int e^{x^2} dx = \int e^u du$$

2. Using $u = x^2 + 1$,

$$\int_0^1 x \sin(x^2 + 1) dx = \int_0^1 \frac{1}{2} \sin u du$$

156/643

Evaluate $\int_0^1 x^7 (x^4 + 1)^5 dx$.

Time permitting, more examples using the substitution rule

Evaluate $\int \sin x \cos x dx$.



CHECK OUR WORK

We can check that $\int \sin x \cos x dx =$ _____ by differentiating.

We can check that $\int \sin x \cos x dx =$ _____ by differentiating.

Evaluate $\int \frac{\log x}{3x} dx$.



CHECK OUR WORK

We can check that $\int \frac{\log x}{3x} dx =$ by differentiating.

Evaluate $\int \frac{e^x}{e^x + 15} dx$.

Evaluate $\int x^4(x^5 + 1)^8 dx$.



CHECK OUR WORK

We can check that $\int \frac{e^x}{e^x + 15} dx =$ by differentiating.

We can check that $\int x^4(x^5 + 1)^8 dx =$ by differentiating.

Evaluate $\int_4^8 \frac{s}{s-3} ds$. Be careful to use correct notation.



165/643

Evaluate $\int x^9(x^5 + 1)^8 dx$.



166/643

CHECK OUR WORK

We can check that $\int x^9(x^5 + 1)^8 dx =$
by differentiating.

167/643

PARTICULARLY TRICKY SUBSTITUTION

Evaluate $\int \frac{1}{e^x + e^{-x}} dx$.




168/643

CHECK OUR WORK

We can check that $\int \frac{1}{e^x + e^{-x}} dx =$ by differentiating.


169/643

TABLE OF CONTENTS




170/643

Find the area between $f(x)$ and $g(x)$ from $x = -1$ to $x = 2$.




171/643

Find the (unsigned) area between $f(x)$ and $g(x)$ from $x = -\frac{\pi}{2}$ to $x = \frac{\pi}{2}$.



172/643


Find the (unsigned) area of the finite region bounded by $f(x)$ and $g(x)$.



173/643




Find the (unsigned) area in the figure below between the curves $f(x)$ and $g(x)$ from $x = 0$ to $x = 1$.



174/643




Set up, but do not evaluate, integral(s) to find the (unsigned) area of the finite region bounded by $x = 1 + y^2$ and $y = x - 3$.



175/643




Set up, but do not evaluate, integral(s) to find the (unsigned) area of the finite region bounded by $x = 1 + y^2$ and $y = x - 3$.




176/643

TABLE OF CONTENTS




177/643

QUICK REFRESHER: VOLUMES OF CYLINDERS



The volume of a cylinder with radius r and height h is:




The volume of a washer, with outer radius R , inner radius r , and height h is:

178/643


QUICK REFRESHER: VOLUMES OF CYLINDERS

More generally, if we have a shape of area A , and we extrude it into a solid of height h , the resulting solid has volume:



179/643

Consider the volume, V , enclosed by rotating the curve $y = \sqrt{x}$, from $x = 0$ to $x = 4$, around the x -axis.




We cut the solid into slices, and approximate the volume of each slice. Each thin slice is *approximately* a cylinder.

If we use n slices, the width of each is:

The radius of the slice at $x = x_i^*$ is:


180/643

Consider the volume, V , enclosed by rotating the curve $y = \sqrt{x}$, from $x = 0$ to $x = 4$, around the x -axis.




$$V \approx \sum_{i=1}^n (\text{volume of each slice})$$

Consider the volume, V , enclosed by rotating the curve $y = \sqrt{x}$, from $x = 0$ to $x = 4$, around the x -axis.



Informally, we think of one slice, at position x , as having thickness dx . So, we can write the volume of this slice as:

Summing up the volumes of slices from $x = 0$ to $x = 4$, our total volume is:




Let h and r be positive constants.

1. What familiar solid results from rotating the line segment from $(0,0)$ to (r,h) around the y -axis?
2. In the informal manner of the last example, describe the volume of a horizontal slice of the solid taken at height y .
3. What is the volume of the entire solid?

Observation

When we rotated around the **horizontal** axis, the width of our cylindrical slices was dx , and our integrand was written in terms of x .


When we rotated around the **vertical** axis, the width of our cylindrical slices was dy , and we integrated in terms of y .



Vertical slices are approximately cylinders

Horizontal slices are approximately cylinders


In this question, we will find the volume enclosed by rotating the curve $y = 1 - x^2$, from $x = -1$ to $x = 2$, about the line $y = 4$.



1. Sketch the surface traced out by the rotating curve.
2. Sketch a cylindrical slice. (Consider: will it be horizontal or vertical?)
3. Give the volume of your slice. Use dx or dy for the width, as appropriate.
4. Integrate (with the appropriate limits of integration) to find the volume of the solid.

185/643

In this question, we will find the volume enclosed by rotating the curve $y = 1 - x^2$, from $x = -1$ to $x = 2$, about the line $y = 4$.




186/643

To find the volume of the entire object, we “add up” the slices from $x = -1$ to $x = 2$ by integrating.

$$\int_{-1}^2 \pi(3 + x^2)^2 dx =$$

187/643


Let A be the area between the curve $y = \log x$ and the x -axis, from $(1, 0)$ to $(e, 1)$. In this question, we will consider the volume of the solid formed by rotating A about the y -axis.



1. Sketch A .
2. Sketch a washer-shaped slice of the solid. (Should it be horizontal or vertical?)
3. Give the volume of your slice. Use dx or dy for the width, as appropriate.
4. Integrate to find the volume of the entire solid.


188/643

Let A be the area between the curve $y = \log x$ and the x -axis, from $(1, 0)$ to $(e, 1)$. In this question, we will consider the volume of the solid formed by rotating A about the y -axis.




189/643

To find the volume of the entire object, we “add up” the slices from $y = 0$ to $y = 1$ by integrating.



190/643


So far, we’ve found the volume of solids formed by rotating a curve. When a point rotates about a fixed centre, the result is a circle, so we could slice those solids up into pieces that are approximately cylinders.



We can find the volumes of other shapes, as long as we can find the areas of their cross-sections.

191/643


The corner of a room is sealed off as follows: On both walls, a parabola of the form $z = (x - 1)^2$ is drawn, where z is the vertical axis and x is the horizontal. They start one metre above the corner, and end one metre to the side of the corner. Taught ropes are strung *horizontally* from one parabola to the other, so the horizontal cross-sections are right triangles. **How much volume is enclosed?**



192/643




A pyramid with height h metres has a square base with side-length b metres. At an elevation of y metres above the base, $0 \leq y \leq h$, the cross-section of the pyramid is a square with side-length $\frac{b}{h}(h - y)$. What is the volume of the pyramid?




OPTIONAL: CHALLENGE QUESTION

A paddle fixed to the x -axis has two flat blades. One blade is in the shape of $f(x) = \frac{8}{3}(x - 1)(x - 5)$, from $x = 1$ to $x = 5$. The other blade is in the shape of $g(x) = x(6 - x)$, $0 \leq x \leq 6$. The paddle turns through a gelatinous fluid, scraping out a hollow cavity as it turns. What is the volume of this cavity? You may leave your answer as an integral, or sum of integrals.



The size of the cavity at a point x along the paddle is determined by the **larger** of $|f(x)|$ and $|g(x)|$.



The size of the cavity at a point x along the paddle is determined by the **larger** of $|f(x)|$ and $|g(x)|$.





TABLE OF CONTENTS



197/643

REVERSE THE PRODUCT RULE

Product Rule:

$$\frac{d}{dx}\{u(x) \cdot v(x)\} = u'(x) \cdot v(x) + u(x) \cdot v'(x)$$

Related fact:

$$\int [u'(x) \cdot v(x) + u(x) \cdot v'(x)] dx = u(x) \cdot v(x) + C$$

Rearrange:

$$\Rightarrow \int [u'(x)v(x)] dx + \int [u(x)v'(x)] dx = u(x)v(x) + C$$

$$\Rightarrow \int [u(x)v'(x)] dx = u(x)v(x) - \int [v(x)u'(x)] dx + C$$

198/643

INTEGRATION BY PARTS

$$\int [u(x)v'(x)] dx = u(x)v(x) - \int [v(x)u'(x)] dx$$

Example: $\int xe^x dx$



199/643

CHECK OUR WORK

In the previous slide, we evaluated

$$\int xe^x dx = xe^x - e^x + C$$

for some constant C . We can check that this is correct by differentiating.

200/643

INTEGRATION BY PARTS (IBP): A CLOSER LOOK

$$\int [u(x)v'(x)] dx = u(x)v(x) - \int [v(x)u'(x)] dx + C$$
$$\underbrace{\int xe^x dx}_{\text{How to integrate??}} = x(e^x) - \underbrace{1 \int e^x dx}_{\text{Easy to integrate!}} + C$$

We start and end with an integral. IBP is only useful if the new integral is somehow an improvement.

We **differentiate** the function we choose as $u(x)$, and **antidifferentiate** the function we choose as $v'(x)$

201/643

CHOOSING $u(x)$ AND $v(x)$

$$\int [u(x)v'(x)] dx = u(x)v(x) - \int [v(x)u'(x)] dx + C$$
$$\int [x \sin x] dx =$$

Option A:	Option B:
$u(x) = x$	$u(x) = \sin x$
$v'(x) = \sin x$	$v'(x) = x$



202/643

CHECK OUR WORK

To check our work, we can calculate $\frac{d}{dx} \{ -x \cos x + \sin x + C \}$. It should work out to be $x \sin x$.

203/643

CHOOSING $u(x)$ AND $v(x)$

$$\int [u(x)v'(x)] dx = u(x)v(x) - \int [v(x)u'(x)] dx + C$$
$$\int [x^2 \log x] dx =$$

Option A:	Option B:
$u(x) = x^2$	$u(x) = \log x$
$v'(x) = \log x$	$v'(x) = x^2$



204/643

CHECK OUR WORK

To check our work, we can calculate $\frac{d}{dx} \left\{ \frac{1}{3}x^3 \log x - \frac{1}{9}x^3 + C \right\}$. It should work out to be $x^2 \log x$.

$$\int [u(x)v'(x)] dx = u(x)v(x) - \int [v(x)u'(x)] dx + C$$
$$\int \left[\frac{1}{2}xe^{6x} \right] dx =$$

Option A:	Option B:
$u(x) = \frac{1}{2}x$	$u(x) = e^{6x}$
$v'(x) = e^{6x}$	$v'(x) = \frac{1}{2}x$



CHECK OUR WORK

We check that $\int \left[\frac{1}{2}xe^{6x} \right] dx = \frac{1}{12}xe^{6x} - \frac{1}{72}e^{6x} + C$ by differentiating.

MNEMONIC

$$\int [u(x)v'(x)] dx = u(x)v(x) - \int [v(x)u'(x)] dx + C$$
$$\int u dv = uv - \int v du + C$$

We abbreviate:

- ▶ $u(x) \rightarrow u$
- ▶ $u'(x) dx \rightarrow du$
- ▶ $v(x) \rightarrow v$
- ▶ $v'(x) dx \rightarrow dv$

CHOOSING u , dv IN YOUR HEAD

Choose u and dv to evaluate the integral below:

$$\int (3t + 5) \cos(t/4) dt$$

Thoughts: $\int u dv = uv - \int v du$
 u gets differentiated, and dv gets antidifferentiated.

209/643

Evaluate, using IBP or Substitution

$$\int u dv = uv - \int v du + C$$

▶ $\int x e^{x^2} dx$

▶ $\int x^2 e^x dx$

▶ $\int e^{x+e^x} dx$



210/643

DEFINITE INTEGRATION BY PARTS

Method 1: Antidifferentiate first, then plug in limits of integration.

Method 2: Plug as you go.

211/643

Evaluate $\int_1^e \log^2 x dx$

212/643

SPECIAL TECHNIQUE: $v'(x) = 1$

$$\int u \, dv = uv - \int v \, du + C$$

Evaluate $\int \log x \, dx$ using integration by parts.



CHECK OUR WORK

Let's check that $\int \log x \, dx = x \log x - x + C$.

$$\int u \, dv = uv - \int v \, du + C$$

Evaluate $\int \arctan x \, dx$ using integration by parts.



CHECK OUR WORK

Let's check that $\int \arctan x \, dx = x \arctan x - \frac{1}{2} \log |1 + x^2| + C$.

Setting $dv = 1 dx$ is a very specific technique. You'll probably only see it in situations integrating logarithms and inverse trigonometric functions.

$$\int \log x dx, \quad \int \arcsin x dx, \quad \int \arccos x dx, \quad \int \arctan x dx, \quad \text{etc.}$$

Evaluate $\int e^x \cos x dx$ using integration by parts.

Let $u = e^x$ and $dv = \cos x dx$. Then $du = e^x dx$ and $v = \sin x$:

$$\int e^x \cos x dx = e^x \sin x - \int e^x \sin x dx$$

Let $u = e^x$ and $dv = \sin x dx$. Then $du = e^x dx$ and $v = -\cos x$:

$$= e^x \sin x - \left[-e^x \cos x - \int -e^x \cos x dx \right]$$

$$= e^x \sin x + e^x \cos x - \int e^x \cos x dx$$

$$\int e^x \cos x dx = e^x \sin x + e^x \cos x - \int e^x \cos x dx$$

$$2 \int e^x \cos x dx = e^x \sin x + e^x \cos x + C$$

$$\int e^x \cos x dx = \frac{1}{2} (e^x \sin x + e^x \cos x) + C$$



INTEGRATING AROUND IN A CIRCLE

We can use this technique to antidifferentiate products of two functions that almost, but don't quite, stay the same under (anti)differentiation.

Use integration by parts a number of times, ending up with an expression involving (a scalar multiple of) the original integral.

To do this, be consistent with your choice of u and dv .

Evaluate $\int \cos(\log x) dx$.



CHECK OUR WORK

We check that $\int \cos(\log x) dx = \frac{x}{2} [\cos(\log x) + \sin(\log x)] + C$ by differentiating.


Evaluate $\int e^{2x} \sin x dx$ using integration by parts.



CHECK OUR WORK

We can check our work by differentiating $\frac{1}{5}e^{2x}[2 \sin x - \cos x] + C$. We should end up with $e^{2x} \sin x$.

TABLE OF CONTENTS



1.8 TRIGONOMETRIC INTEGRALS

Recall:

- ▶ $\sin^2 x + \cos^2 x = 1$
- ▶ $\tan^2 x + 1 = \sec^2 x$
- ▶ $\sin^2 x = \frac{1}{2}(1 - \cos 2x)$
- ▶ $\cos^2 x = \frac{1}{2}(1 + \cos 2x)$
- ▶ $\sin(2x) = 2 \sin x \cos x$

INTEGRATING PRODUCTS OF SINE AND COSINE

$$\int \sin x \cos x \, dx =$$



INTEGRATING PRODUCTS OF SINE AND COSINE

$$\int \sin x \cos x \, dx =$$

$$\int \sin^{10} x \cos x \, dx =$$



CHECK OUR WORK

If we are correct that $\int \sin x \cos x \, dx =$ _____, then it should be true that $\frac{d}{dx} \left\{ \right\} = \sin x \cos x$.

CHECK OUR WORK

If we are correct that $\int \sin^{10} x \cos x \, dx =$ _____, then it should be true that $\frac{d}{dx} \left\{ \right\} = \sin^{10} x \cos x$.

229/643

INTEGRATING PRODUCTS OF SINE AND COSINE

$$\int_0^{\frac{\pi}{2}} \sin^{\pi+1} x \cos x \, dx =$$



230/643

CHECK OUR WORK

If we are correct that $\int \sin^{\pi+1} x \cos x \, dx =$ _____, then it should be true that $\frac{d}{dx} \left\{ \right\} = \sin^{\pi+1} x \cos x$.

231/643

INTEGRATING PRODUCTS OF SINE AND COSINE

Let $u = \sin x$, $du = \cos x \, dx$.

$$\int \sin^{10} x \cos^3 x \, dx =$$



232/643

CHECK OUR WORK

If we are correct that $\int \sin^{10} x \cos^3 x \, dx =$ _____, then it should be true that $\frac{d}{dx} \left\{ \right\} = \sin^{10} x \cos^3 x$.

233/643

INTEGRATING PRODUCTS OF SINE AND COSINE

$$\int \sin^5 x \cos^4 x \, dx =$$



234/643

CHECK OUR WORK

If we are correct that $\int \sin^5 x \cos^4 x \, dx =$ _____, then it should be true that $\frac{d}{dx} \left\{ \right\} = \sin^5 x \cos^4 x$.

235/643

GENERALIZE: $\int \sin^m x \cos^n x \, dx$

To use the substitution $u = \sin x$, $du = \cos x \, dx$:

- ▶ We need to **reserve** one $\cos x$ for the differential.
- ▶ We need to **convert** the remaining $\cos^{n-1} x$ to $\sin x$ terms.
- ▶ We convert using $\cos^2 x = 1 - \sin^2 x$. To avoid square roots, that means $n - 1$ should be **even when we convert**.
- ▶ **So, we can use this substitution when the original power of cosine, n , is ODD: one cosine goes to the differential, the rest are converted to sines.**

236/643

GENERALIZE: $\int \sin^m x \cos^n x \, dx$

To use the substitution $u = \cos x$, $du = -\sin x \, dx$:

- ▶ We need to **reserve** one $\sin x$ for the differential.
- ▶ We need to **convert** the remaining $\sin^{m-1} x$ to $\cos x$ terms.
- ▶ We convert using $\sin^2 x = 1 - \cos^2 x$. To avoid square roots, that means $m - 1$ should be **even when we convert**.
- ▶ **So, we can use this substitution when the original power of sine, m , is ODD: one sine goes to the differential, the rest are converted to cosines.**

237/643

MNEMONIC: "ODD ONE OUT"

Integrating $\int \sin^m x \cos^n x \, dx$

If you want to use $u = \sin x$, there should be an odd power of **cosine**.

If you want to use $u = \cos x$, there should be an odd power of **sine**.

238/643

Carry out a suitable substitution (but do not evaluate the resulting integral):

▶ $\int \sin^4 x \cos^7 x \, dx$

▶ $\int \sin^7 x \cos^4 x \, dx$

▶ $\int \sin^7 x \cos^7 x \, dx$



239/643

To evaluate $\int \sin^m x \cos^n x \, dx$, we use:

- ▶ $u = \sin x$ if n is odd, and/or
- ▶ $u = \cos x$ if m is odd

What if n and m are both even?

$$\cos^2 x = \frac{1 + \cos 2x}{2} \qquad \sin^2 x = \frac{1 - \cos 2x}{2}$$

240/643

$$\cos^2 x = \frac{1 + \cos 2x}{2} \quad \sin^2 x = \frac{1 - \cos 2x}{2}$$

$$\int \sin^2 x \, dx =$$



CHECK OUR WORK

We check that $\int \sin^2 x \, dx =$ by differentiating:

$$\cos^2 x = \frac{1 + \cos 2x}{2} \quad \sin^2 x = \frac{1 - \cos 2x}{2}$$

Evaluate $\int \sin^4 x \, dx$.



CHECK OUR WORK

We want to check that $\int \sin^4 x \, dx =$

Recall:

- ▶ $\frac{d}{dx}\{\tan x\} = \sec^2 x$
- ▶ $\frac{d}{dx}\{\sec x\} = \sec x \tan x$
- ▶ $\tan^2 x + 1 = \sec^2 x$

245/643

$$\int \tan x \, dx =$$

246/643

CHECK OUR WORK

Let's check that $\int \tan x \, dx =$ by differentiating.

247/643

Optional: A nifty trick – you won't be expected to come up with it. There is some motivation for the trick in Example 1.8.19 in the CLP-2 text.

$$\int \sec x \, dx = \int \sec x \left(\frac{\sec x + \tan x}{\sec x + \tan x} \right) dx$$

248/643

Useful integrals:

$$\blacktriangleright \int \tan x \, dx = \log |\sec x| + C$$

$$\blacktriangleright \int \sec x \, dx = \log |\sec x + \tan x| + C$$

$$1. \int \sec x \tan x \, dx =$$

$$2. \int \sec^2 x \, dx =$$

$$3. \int \tan x \, dx =$$

$$4. \int \sec x \, dx =$$

Evaluate using the substitution rule:

$$\int \tan^5 x \sec^2 x \, dx =$$

$$\int \sec^4 x (\sec x \tan x) \, dx =$$



CHECK OUR WORK

Let's check that $\int \tan^5 x \sec^2 x \, dx =$ by differentiating.

Evaluate using the identity $\sec^2 x = 1 + \tan^2 x$

$$\int \tan^4 x \sec^6 x \, dx =$$

$$\int \tan^3 x \sec^5 x \, dx =$$



CHECK OUR WORK

Let's check that $\int \tan^4 x \sec^6 x \, dx =$

CHOOSING A SUBSTITUTION: $\int \tan^m x \sec^n x \, dx$

Using $u = \sec x$, $du = \sec x \tan x \, dx$:

- ▶ Reserve $\sec x \tan x$ for the differential.
(m, n should each be at least 1)
- ▶ From the remaining $\tan^{m-1} x \sec^{n-1} x$, convert all tangents to secants using $\tan^2 x + 1 = \sec^2 x$.
($m - 1$ should be even, to avoid square roots)

To use the substitution $u = \sec x$, $du = \sec x \tan x \, dx$ to evaluate $\int \tan^m x \sec^n x \, dx$, n should be , and m should be .

CHOOSING A SUBSTITUTION: $\int \tan^m x \sec^n x \, dx$

Using $u = \tan x$, $du = \sec^2 x \, dx$:

- ▶ Reserve for the differential.
- ▶ From the remaining terms, convert all to using $\tan^2 x + 1 = \sec^2 x$.

To use the substitution $u = \tan x$, $du = \sec^2 x \, dx$ to evaluate $\int \tan^m x \sec^n x \, dx$, n should be .

Evaluating $\int \tan^m x \sec^n x \, dx$

To evaluate $\int \tan^m x \sec^n x \, dx$, we can use:

- ▶ $u = \sec x$ if m is odd and $n \geq 1$
- ▶ $u = \tan x$ if n is even and $n \geq 2$

Choose a substitution for the integrals below.

▶ $\int \sec^2 x \tan^3 x \, dx$

▶ $\int \sec^2 x \tan^2 x \, dx$

▶ $\int \sec^3 x \tan^3 x \, dx$



$$\int \sec^2 x \tan^2 x \, dx$$



$$\int \sec^3 x \tan^3 x \, dx$$



Evaluate $\int \tan^3 x \, dx = \int \frac{\sin^3 x}{\cos^3 x} \, dx$

CHECK OUR WORK

Let's check that $\int \tan^3 x \, dx =$ by differentiating.

261/643

Generalizing the last example:

$$\begin{aligned}\int \tan^m x \sec^n x \, dx &= \int \left(\frac{\sin x}{\cos x}\right)^m \left(\frac{1}{\cos x}\right)^n dx \\ &= \int \frac{\sin^m x}{\cos^{m+n} x} dx \\ &= \int \left(\frac{\sin^{m-1} x}{\cos^{m+n} x}\right) \sin x \, dx\end{aligned}$$

To use $u = \cos x$, $du = -\sin x \, dx$: we will convert $\sin^{m-1}(x)$ into cosines, so $m - 1$ must be even, so m must be odd.

262/643

Evaluating $\int \tan^m x \sec^n x \, dx$


To evaluate $\int \tan^m x \sec^n x \, dx$, we can use:

- ▶ $u = \sec x$ if m is odd and $n \geq 1$
- ▶ $u = \tan x$ if n is even and $n \geq 2$
- ▶ $u = \cos x$ if m is odd
- ▶ $u = \tan x$ if m is even and $n = 0$ (after using $\tan^2 x = \sec^2 x - 1$, maybe several times)

Evaluate $\int \tan^2 x \, dx$

263/643

TABLE OF CONTENTS



264/643

Evaluating $\int \tan^m x \sec^n x \, dx$

To evaluate $\int \tan^m x \sec^n x \, dx$, we can use:

- ▶ $u = \sec x$ if m is odd and $n \geq 1$
- ▶ $u = \tan x$ if n is even and $n \geq 2$
- ▶ $u = \cos x$ if m is odd
- ▶ $u = \tan x$ if m is even and $n = 0$
(after using $\tan^2 x = \sec^2 x - 1$, maybe several times)

Remaining case: n odd and m is even.

The general remaining case is known, but complicated. Instead of treating it exhaustively, we'll show examples of two methods.

$\int \sec x \, dx$

We saw a way of integrating secant with the following trick:

$$\begin{aligned} \int \sec x \, dx &= \int \sec x \left(\frac{\sec x + \tan x}{\sec x + \tan x} \right) dx = \int \frac{\sec^2 x + \sec x \tan x}{\sec x + \tan x} dx \\ &= \int \frac{1}{u} du \quad \text{with } u = \sec x + \tan x \end{aligned}$$

Another trick: this time let $u = \sin x$, $du = \cos x \, dx$:

$$\begin{aligned} \int \sec x \, dx &= \int \frac{1}{\cos x} dx = \int \frac{\cos x}{\cos^2 x} dx \\ &= \int \frac{1}{1 - \sin^2 x} \cos x \, dx = \int \frac{1}{1 - u^2} du \end{aligned}$$

The integrand $\frac{1}{1-u^2}$ is a rational function of u (i.e. a ratio of two polynomials). There is a procedure, called Partial Fractions, that can be used to evaluate all integrals of rational functions. We will learn it in Section 1.10.

$\int \sec^3 x \, dx$

We can integrate around in a circle (with integration by parts) to evaluate $\int \sec^3 x \, dx$. Let $u = \sec x$, $dv = \sec^2 x \, dx$. Then $du = \sec x \tan x \, dx$ and $v = \tan x$.


$$\begin{aligned} \int \sec^3 x \, dx &= \sec x \tan x - \int \sec x \tan^2 x \, dx \\ &= \sec x \tan x - \int \sec x (\sec^2 x - 1) \, dx \\ &= \sec x \tan x - \int \sec^3 x \, dx + \int \sec x \, dx \\ &= \sec x \tan x - \int \sec^3 x \, dx + \log |\sec x + \tan x| + C' \end{aligned}$$

$$2 \int \sec^3 x \, dx = \sec x \tan x + \log |\sec x + \tan x| + C'$$

$$\int \sec^3 x \, dx = \frac{1}{2} (\sec x \tan x + \log |\sec x + \tan x|) + C$$

with $C = C'/2$.

TABLE OF CONTENTS



WARMUP

Evaluate $\int_3^7 \frac{1}{\sqrt{x^2 + 2x + 1}} dx$.

269/643

Evaluate $\int \frac{1}{\sqrt{x^2 + 1}} dx$.

270/643

CHECK OUR WORK

Let's verify that $\int \frac{1}{\sqrt{x^2 + 1}} =$
Seems improbable, right?

271/643

METHOD (ONE STANDARD CASE)

- ▶ An integrand has the form: $\sqrt{\text{quadratic}}$, and we'd like to cancel off the square root.
- ▶ So, we need to write our quadratic expression as a perfect square. Choose a helpful substitution:
 - ▶ $x = \sin \theta$, $1 - \sin^2 \theta = \cos^2 \theta$ changes $\sqrt{1 - x^2}$ into
 - ▶ $x = \tan \theta$, $1 + \tan^2 \theta = \sec^2 \theta$ changes $\sqrt{1 + x^2}$ into
 - ▶ $x = \sec \theta$, $\sec^2 \theta - 1 = \tan^2 \theta$ changes $\sqrt{x^2 - 1}$ into
- ▶ After integrating, convert back to the original variable (possibly using a triangle—more details later)

272/643

FOCUS ON THE ALGEBRA

$$1 - \sin^2 \theta = \cos^2 \theta \quad 1 + \tan^2 \theta = \sec^2 \theta \quad \sec^2 \theta - 1 = \tan^2 \theta$$

Choose a trigonometric substitution that will allow the square root to cancel out of the following expressions:

▶ $\sqrt{x^2 - 1}$

▶ $\sqrt{x^2 + 1}$

▶ $\sqrt{1 - x^2}$

273/643

FOCUS ON THE ALGEBRA

$$1 - \sin^2 \theta = \cos^2 \theta \quad 1 + \tan^2 \theta = \sec^2 \theta \quad \sec^2 \theta - 1 = \tan^2 \theta$$

Choose a trigonometric substitution that will allow the square root to cancel out of the following expressions:

▶ $\sqrt{x^2 + 7}$

▶ $\sqrt{3 - 2x^2}$

274/643

CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

Consider the substitution $x = \sin \theta$, $dx = \cos \theta d\theta$ for the integral:

$$\int_0^1 \sqrt{1 - x^2} dx$$

When $x = 0$ (lower limit of integration), what is θ ?

When $x = 1$ (upper limit of integration), what is θ ?

275/643

CLOSER LOOK AT ABSOLUTE VALUES

▶ SKIP CLOSER LOOK

More generally, suppose a is a positive constant and we use the substitution $x = a \sin \theta$ for the term $\sqrt{a^2 - x^2}$.

276/643

CLOSER LOOK AT ABSOLUTE VALUES

▶▶ SKIP CLOSER LOOK

Now, consider the substitution $x = a \tan \theta$ for $\sqrt{a^2 + x^2}$, where a is a positive constant.

277/643

CLOSER LOOK AT ABSOLUTE VALUES

▶▶ SKIP CLOSER LOOK

Finally, consider the substitution $x = a \sec \theta$ for $\sqrt{x^2 - a^2}$, where a is a positive constant.

278/643

ABSOLUTE VALUES

From now on, we will assume:

- ▶ With the substitution $x = a \sin \theta$ for $\sqrt{a^2 - x^2}$, $|\cos \theta| = \cos \theta$
- ▶ With the substitution $x = a \tan \theta$ for $\sqrt{a^2 + x^2}$, $|\sec \theta| = \sec \theta$

279/643

Identities

$$1 - \sin^2 \theta = \cos^2 \theta$$

$$\sin(2\theta) = 2 \sin \theta \cos \theta$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$\sin^2 \theta = \frac{1 - \cos(2\theta)}{2}$$

$$\sec^2 \theta - 1 = \tan^2 \theta$$

$$\cos^2 \theta = \frac{1 + \cos(2\theta)}{2}$$

Evaluate $\int_0^1 (1 + x^2)^{-3/2} dx$

280/643

Identities

$$1 - \sin^2 \theta = \cos^2 \theta$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$\sec^2 \theta - 1 = \tan^2 \theta$$

$$\sin(2\theta) = 2 \sin \theta \cos \theta$$

$$\sin^2 \theta = \frac{1 - \cos(2\theta)}{2}$$

$$\cos^2 \theta = \frac{1 + \cos(2\theta)}{2}$$

Evaluate $\int \sqrt{1 - 4x^2} \, dx$

CHECK OUR WORK

In the last example, we computed

$$\int \sqrt{1 - 4x^2} \, dx =$$

To check, we differentiate.

Identities

$$1 - \sin^2 \theta = \cos^2 \theta$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$\sec^2 \theta - 1 = \tan^2 \theta$$

$$\sin(2\theta) = 2 \sin \theta \cos \theta$$

$$\sin^2 \theta = \frac{1 - \cos(2\theta)}{2}$$

$$\cos^2 \theta = \frac{1 + \cos(2\theta)}{2}$$

Evaluate $\int \frac{1}{\sqrt{x^2 - 1}} \, dx$

CHECK OUR WORK

Let's check our result, $\int \frac{1}{\sqrt{x^2 - 1}} \, dx =$

COMPLETING THE SQUARE

Choose a trigonometric substitution to simplify $\sqrt{3 - x^2 + 2x}$.

Identities have two “parts” that turn into one part:

- ▶ $1 - \sin^2 \theta = \cos^2 \theta$
- ▶ $1 + \tan^2 \theta = \sec^2 \theta$
- ▶ $\sec^2 \theta - 1 = \tan^2 \theta$

But our quadratic expression has *three* parts.

Fact: $3 - x^2 + 2x = 4 - (x - 1)^2$

285/643

COMPLETING THE SQUARE

$$(x + b)^2 = x^2 + 2bx + b^2$$
$$c - (x + b)^2 = (c - b^2) - x^2 - 2bx$$

Write $3 - x^2 + 2x$ in the form $c - (x + b)^2$ for constants b, c .

1. Find b :
2. Solve for c :
3. All together:

286/643

Evaluate $\int \frac{x^2 - 6x + 9}{\sqrt{6x - x^2}} dx$.

Identities have two “parts” that turn into one part:

- ▶ $1 - \sin^2 \theta = \cos^2 \theta$
- ▶ $1 + \tan^2 \theta = \sec^2 \theta$
- ▶ $\sec^2 \theta - 1 = \tan^2 \theta$

One of those parts is a constant, and one is squared.

287/643

Evaluate $\int \frac{x^2 - 6x + 9}{\sqrt{6x - x^2}} dx = \int \frac{(x - 3)^2}{\sqrt{9 - (x - 3)^2}} dx$.

288/643


CHECK OUR WORK

Let's verify that

$$\int \frac{x^2 - 6x + 9}{\sqrt{6x - x^2}} =$$

289/643

TABLE OF CONTENTS



290/643

MOTIVATION

How to integrate $\int \frac{x-2}{(x+1)(2x-1)} dx$?

Useful fact: $\frac{x-2}{(x+1)(2x-1)} = \frac{1}{x+1} - \frac{1}{2x-1}$

So:

$$\begin{aligned} \int \frac{x-2}{(x+1)(2x-1)} dx &= \int \frac{1}{x+1} dx - \int \frac{1}{2x-1} dx \\ &= \log|x+1| - \frac{1}{2} \log|2x-1| + C \end{aligned}$$

Method of Partial Fractions: **Algebraic method** to turn any rational function (i.e. ratio of two polynomials) into the sum of easier-to-integrate rational functions.

291/643

DISTINCT LINEAR FACTORS

The rational function

$$\frac{\text{numerator}}{K(x-a_1)(x-a_2)\cdots(x-a_j)}$$

can be written as

$$\frac{A_1}{x-a_1} + \frac{A_2}{x-a_2} + \cdots + \frac{A_j}{x-a_j}$$

for some constants A_1, A_2, \dots, A_j , provided

(1) the linear roots a_1, \dots, a_j are distinct, and

(2) the degree of the numerator is strictly less than the degree of the denominator.

292/643 Equation 1.10.7

DISTINCT LINEAR FACTORS

$$\frac{7x + 13}{(2x + 5)(x - 2)} =$$



293/643

DISTINCT LINEAR FACTORS

We found $7x + 13 = A(x - 2) + B(2x + 5)$ for some constants A and B .
What are A and B ?

Method 1: set x to convenient values.

Method 2: match coefficients of powers of x .

294/643

DISTINCT LINEAR FACTORS

All together:

$$\frac{7x + 13}{2x^2 + x - 10} = \frac{A}{2x + 5} + \frac{B}{x - 2}$$

$A = 1, \quad B = 3$

295/643

CHECK OUR WORK

We check that $\int \frac{7x + 13}{2x^2 + x - 10} =$ by
differentiating.

296/643

DISTINCT LINEAR FACTORS

$\frac{x^2 + 5}{2x(3x + 1)(x + 5)}$ is hard to antidifferentiate, but it can be written as $\frac{A}{2x} + \frac{B}{3x + 1} + \frac{C}{x + 5}$ for some constants A , B , and C .

Once we find A , B , and C , integration is easy:

$$\begin{aligned} \int \frac{x^2 - 24x + 5}{2x(3x + 1)(x + 5)} dx \\ &= \int \left(\frac{A}{2x} + \frac{B}{3x + 1} + \frac{C}{x + 5} \right) dx \\ &= \frac{A}{2} \log |x| + \frac{B}{3} \log |3x + 1| + C \log |x + 5| + D \end{aligned}$$



297/643

DISTINCT LINEAR FACTORS

$$\frac{x^2 + 5}{2x(3x + 1)(x + 5)} = \frac{A}{2x} + \frac{B}{3x + 1} + \frac{C}{x + 5}$$

Find constants A , B , and C .

Start: make a common denominator

298/643

CHECK OUR WORK

Let's check that

$$\frac{x^2 + 5}{2x(3x + 1)(x + 5)} =$$

299/643

DISTINCT LINEAR FACTORS

All together:

$$\frac{x^2 + 5}{2x(3x + 1)(x + 5)} = \frac{1}{2x} - \frac{23/14}{3x + 1} + \frac{3/14}{x + 5}$$

300/643

Repeated Linear Factors

A rational function $\frac{P(x)}{(x-1)^4}$, where $P(x)$ is a polynomial of degree strictly less than 4, can be written as

$$\frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{(x-1)^3} + \frac{D}{(x-1)^4}$$

for some constants $A, B, C,$ and $D.$

$$\frac{5x-11}{(x-1)^2} =$$



Set up the form of the partial fractions decomposition. (You do not have to solve for the parameters.)


$$\frac{3x+16}{(x+5)^3} =$$

$$\frac{-2x-10}{(x+1)^2(x-1)} =$$




IRREDUCIBLE QUADRATIC FACTORS

Sometimes it's not possible to factor our denominator into linear factors with real terms.



If a quadratic function has real roots a and b (possibly $a = b$, possibly $a \neq b$), then we can write it as $c(x-a)(x-b)$ for some constant $c.$



If a quadratic function has no real roots, then it can't be factored into (real) linear factors. It is **irreducible.**

IRREDUCIBLE QUADRATIC FACTORS

When the denominator has an irreducible quadratic factor $x^2 + bx + c,$ we add a term $\frac{Ax+B}{x^2+bx+c}$ to our composition. (The degree of the numerator must still be smaller than the degree of the denominator.) Write out the form of the partial fraction decomposition (but do not solve for the parameters):

$$\blacktriangleright \frac{1}{(x+1)(x^2+1)} =$$

$$\blacktriangleright \frac{3x^2-x+5}{(x^2+1)(x^2+2)} =$$



The purpose of the partial fraction decomposition is to end up with functions **that we can integrate**.

► Recall: $\int \frac{1}{x^2 + 1} dx = \arctan x + C.$

► Evaluate: $\int \frac{1}{(x + 1)^2 + 1} dx$



Evaluate $\int \frac{4}{(3x + 8)^2 + 9} dx$



CHECK OUR WORK

We found $\int \frac{4}{(3x + 8)^2 + 9} dx =$



Evaluate $\int \frac{x + 1}{x^2 + 2x + 2} dx.$

(Hint: start by completing the square.)



CHECK OUR WORK

We found $\int \frac{x+1}{x^2+2x+2} dx =$

309/643

These rules work **only** when the degree of the numerator is **less than** the degree of the denominator.

$$\int \frac{x^3}{(x-2)^2(x-3)(x-4)^2} dx \quad \int \frac{x^5}{(x-2)^2(x-3)(x-4)^2} dx$$

If the degree of the numerator is too large, we use polynomial long division.

310/643

Evaluate $\int \frac{8x^2 + 22x + 23}{2x + 3} dx$.



311/643

CHECK OUR WORK

We computed

$$\int \frac{8x^2 + 22x + 23}{2x + 3} dx =$$

312/643

Evaluate $\int \frac{3x^3 + x + 3}{x - 2} dx$.



CHECK OUR WORK

We found

$$\int \frac{3x^3 + x + 3}{x - 2} dx =$$

Evaluate $\int \frac{3x^2 + 1}{x^2 + 5x} dx$.



CHECK OUR WORK

We found $\int \frac{3x^2 + 1}{x^2 + 5x} dx =$

FACTORING

$$P(x) = x^3 + 2x^2 - 5x - 6$$


317/643

FACTORING

$$P(x) = 2x^3 - 3x^2 + 4x - 6$$

318/643

TABLE OF CONTENTS



319/643

Sometimes, integrals can't be evaluated using the fundamental theorem of calculus:


$$\int_0^1 e^{x^2} dx = ? \quad \int_0^1 \sin(x^2) dx = ?$$

Sometimes, integrals can be evaluated, but only in terms of complicated constant numbers:

$$\int_0^3 \frac{1}{1+x^2} dx = \arctan(3) = \dots ?$$


A **numerical approximation** will give us an approximate **number** for a definite integral.

320/643




We can approximate the area $\int_a^b f(x) dx$ by cutting it into slices and approximating the area of those slices with a simple geometric figure, such as a rectangle, a trapezoid, or a parabola.

The **midpoint rule** approximates $\int_a^b f(x) dx$ as its midpoint Riemann sum with n intervals.



Approximate the area under the curve $y = f(x)$ from $x = x_{j-1}$ to $x = x_j$ with a rectangle.

To make our writing cleaner, let $\bar{x}_j = \frac{x_{j-1} + x_j}{2}$




Midpoint Rule

The midpoint rule approximation is

$$\int_a^b f(x) dx \approx [f(\bar{x}_1) + f(\bar{x}_2) + \dots + f(\bar{x}_n)] \Delta x$$

where $\Delta x = \frac{b-a}{n}$ and $x_j = a + j\Delta x$

Approximate $\int_0^1 \frac{4}{1+x^2} dx$ using the midpoint rule and $n = 4$ slices. Leave your answer in calculator-ready form.



ERROR

$$\pi = \int_0^1 \frac{4}{1+x^2} dx \approx \left[\frac{4}{1+(\frac{1}{8})^2} + \frac{4}{1+(\frac{3}{8})^2} + \frac{4}{1+(\frac{5}{8})^2} + \frac{4}{1+(\frac{7}{8})^2} \right] \cdot \frac{1}{4}$$

$$\approx 3.14680$$

Error: $|\text{exact} - \text{approximate}|$

Relative error: $\left| \frac{\text{exact} - \text{approximate}}{\text{exact}} \right|$

Percent error: $100 \left| \frac{\text{exact} - \text{approximate}}{\text{exact}} \right|$

325/643

ERROR

A numerical approximation will give us an approximate value for a definite integral.

This is most useful if we know something about its accuracy.

A: approximation E: exact number

Error: $|A - E|$


Relative Error: $\left| \frac{A - E}{E} \right|$

Percent Error: $100 \left| \frac{A - E}{E} \right|$

We will discuss error more after we've learned the three approximation rules. For now, we're using error to illustrate that our methods have the potential to produce reasonable approximations without too much work.


326/643 Definition 1.11.4

The **trapezoidal rule** approximates each slice of $\int_a^b f(x) dx$ with a trapezoid.




327/643

Recall the area of a right trapezoid with base b and heights h_1 and h_2 :



328/643



Trapezoid area: $\frac{\text{base}}{2}(h_1 + h_2)$

Approximate $\int_0^1 e^{x^2} dx$ using $n = 4$ trapezoids.
Leave your answer in calculator-ready form.

Trapezoidal Rule


The trapezoidal rule approximation is

$$\int_a^b f(x) dx \approx \Delta x \left[\frac{1}{2} f(x_0) + f(x_1) + f(x_2) + \cdots + f(x_{n-1}) + \frac{1}{2} f(x_n) \right]$$

where $\Delta x = \frac{b-a}{n}$ and $x_i = a + i\Delta x$


Using $n = 3$ trapezoids, approximate $\int_1^{10} \frac{1}{x} dx$.

Simpson's rule approximates each pair of slices of $\int_a^b f(x) dx$ with a parabola.




SIMPSON'S RULE

Add up **parabolas**.



SIMPSON'S RULE DERIVATION

▶ SKIP DERIVATION OF SIMPSON'S RULE



What is the area under the parabola passing through three specified points?

333/643

Find

$$\frac{h}{3} (2Ah^2 + 6C)$$

for A , B , and C such that

$$Ah^2 - Bh + C = f(-h) \quad (E1)$$


$$C = f(0) \quad (E2)$$

$$Ah^2 + Bh + C = f(h) \quad (E3)$$

Try $(E1) + 4(E2) + (E3)$:

$$\begin{aligned} 2Ah^2 + 6C &= f(-h) + 4f(0) + f(h) \\ \text{Area} = \frac{h}{3} (2Ah^2 + 6C) &= \frac{h}{3} (f(-h) + 4f(0) + f(h)) \end{aligned}$$


334/643



Area under parabola:

$$\frac{\Delta x}{3} (f(x_1) + 4f(x_2) + f(x_3))$$

335/643



336/643

Simpson's Rule

The Simpson's rule approximation is $\int_a^b f(x) dx \approx$

$$\frac{\Delta x}{3} [f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + 2f(x_4) + \cdots + 4f(x_{n-1}) + f(x_n)]$$

where n is even, $\Delta x = \frac{b-a}{n}$, and $x_i = a + i\Delta x$

Using Simpson's rule and $n = 8$ (i.e. 4 parabolas),

approximate $\int_1^{17} \frac{1}{x} dx$. Leave your answer in calculator-ready form.

(We'll call n the number of slices; some people call $n/2$ the number of slices, because that's the number of approximating parabolas.)

The instantaneous electricity use rate (kW/hr) of a factory is measured throughout the day.

time	12:00	1:00	2:00	3:00	4:00	5:00	6:00	7:00	8:00
rate	100	200	150	400	300	300	200	100	150

Use Simpson's Rule to approximate the total amount of electricity you used from noon to 8:00.

Numerical integration errors

Assume that $|f''(x)| \leq M$ for all $a \leq x \leq b$ and $|f^{(4)}(x)| \leq L$ for all $a \leq x \leq b$. Then

- ▶ the total error introduced by the midpoint rule is bounded by $\frac{M(b-a)^3}{24n^2}$,
- ▶ the total error introduced by the trapezoidal rule is bounded by $\frac{M(b-a)^2}{12n}$, and
- ▶ the total error introduced by Simpson's rule is bounded by $\frac{L(b-a)^5}{180n^4}$

when approximating $\int_a^b f(x) dx$.

Numerical integration errors

Assume that $|f''(x)| \leq M$ for all $a \leq x \leq b$. Then the total error introduced by the midpoint rule is bounded by $\frac{M(b-a)^3}{24n^2}$ when approximating $\int_a^b f(x) dx$.

Suppose we approximate $\int_0^3 \sin(x) dx$ using the midpoint rule and $n = 6$ intervals. Give an upper bound of the resulting error.



Numerical integration errors

Assume that $|f^{(4)}(x)| \leq L$ for all $a \leq x \leq b$. Then the total error introduced by Simpson's rule is bounded by $\frac{L}{180} \frac{(b-a)^5}{n^4}$ when approximating $\int_a^b f(x) dx$.

Suppose we approximate $\int_2^3 \frac{1}{x} dx$ using Simpson's rule with $n = 10$ slices (5 parabolas). Give an upper bound of the resulting error.



Numerical integration errors

Assume that $|f^{(4)}(x)| \leq L$ for all $a \leq x \leq b$. Then the total error introduced by Simpson's rule is bounded by $\frac{L}{180} \frac{(b-a)^5}{n^4}$ when approximating $\int_a^b f(x) dx$.

We will approximate $\int_0^{1/2} e^{x^2} dx$ using Simpson's rule, and we need our error to be no more than $\frac{1}{10000}$. How many intervals will suffice?

You may use, without proof:

$$\frac{d^4}{dx^4} \{e^{x^2}\} = 4e^{x^2} (4x^4 + 12x^2 + 3) \quad \frac{25\sqrt[4]{e}}{180 \cdot 2^5} < \frac{1}{3^4}$$



Numerical integration errors

Assume that $|f^{(4)}(x)| \leq L$ for all $a \leq x \leq b$. Then the total error introduced by Simpson's rule is bounded by $\frac{L}{180} \frac{(b-a)^5}{n^4}$ when approximating $\int_a^b f(x) dx$.


It can be shown that the fourth derivative of $\frac{1}{x^2+1}$ has absolute value at most 24 for all real numbers x . Using this information, find a rational number approximating $\arctan(2)$ with an error of no more than $\frac{2^6}{3 \cdot 5^5} \approx 0.007$.



First, we'll set up our integral:



TABLE OF CONTENTS



345/643

Numerical integration errors

Assume that $|f''(x)| \leq M$ for all $a \leq x \leq b$ and $|f^{(4)}(x)| \leq L$ for all $a \leq x \leq b$. Then

- ▶ the total error introduced by the midpoint rule is bounded by $\frac{M(b-a)^3}{24n^2}$,
- ▶ the total error introduced by the trapezoidal rule is bounded by $\frac{M(b-a)^3}{12n^2}$, and
- ▶ the total error introduced by Simpson's rule is bounded by $\frac{L(b-a)^5}{180n^4}$

when approximating $\int_a^b f(x) dx$.


346/643 Theorem 1.11.12

WHY THE *second* DERIVATIVE?

The midpoint rule gives the exact area under the curve for

$$f(x) = ax + b$$

when a and b are any constants.



The first derivative can be large without causing a large error.

347/643

Numerical integration errors


Assume that $|f''(x)| \leq M$ for all $a \leq x \leq b$ and $|f^{(4)}(x)| \leq L$ for all $a \leq x \leq b$. Then

- ▶ the total error introduced by the midpoint rule is bounded by $\frac{M(b-a)^3}{24n^2}$,
- ▶ the total error introduced by the trapezoidal rule is bounded by $\frac{M(b-a)^3}{12n^2}$, and
- ▶ the total error introduced by Simpson's rule is bounded by $\frac{L(b-a)^5}{180n^4}$

when approximating $\int_a^b f(x) dx$.

348/643 Theorem 1.11.12

We'll start small: let's consider one-half of a single interval being approximated using the midpoint rule.
To avoid messiness, let's also consider a simplified location:



We want to relate the actual area of this half-slice to its approximate area:

$$\int_0^q f(x) dx \approx q \cdot f(0)$$

$$\int_0^q f(x) dx \approx q \cdot f(0)$$


If you squint just right, the right-hand side looks a bit like the "u · v'" term from integration by parts, where u = f(x) and dv = dx.

- Set u = f(x) and dv = dx, so du = f'(x) dx.
We choose v(x) = x - q, so that f(v(q)) = f(0).

$$\begin{aligned} \int_0^q f(x) dx &= [(x - q)f(x)]_0^q - \int_0^q (x - q)f'(x) dx \\ &= q \cdot f(0) - \int_0^q (x - q)f'(x) dx \end{aligned}$$

- We know something about the second derivative, not the first, so repeat: set u = f'(x), dv = (x - q) dx; du = f''(x) dx, v = $\frac{(x - q)^2}{2}$


$$\int_0^q f(x) dx = q \cdot f(0) + \frac{q^2}{2} \cdot f'(0) + \int_0^q \frac{(x - q)^2}{2} f''(x) dx$$



$$\int_0^q f(x) dx = \underbrace{q \cdot f(0)}_{\text{exact}} + \underbrace{\frac{q^2}{2} \cdot f'(0)}_{\text{approximate}} + \underbrace{\int_0^q \frac{(x - q)^2}{2} f''(x) dx}_{\pm \text{error}}$$


Repeat for the other half of the slice:

$$\begin{aligned} \int_{-q}^0 f(x) dx &= \left[\underbrace{f(x)}_u \cdot \underbrace{(x + q)}_v \right]_{-q}^0 - \int_{-q}^0 \underbrace{(x + q)}_v \cdot \underbrace{f'(x)}_{du} dx \\ &= q \cdot f(0) - \int_{-q}^0 \underbrace{f'(x)}_{\dot{u}} \cdot \underbrace{(x + q)}_{d\dot{v}} dx \\ &= q \cdot f(0) - \left[\underbrace{f'(x)}_{\dot{u}} \cdot \underbrace{\frac{(x + q)^2}{2}}_{\dot{v}} \right]_{-q}^0 + \int_{-q}^0 \underbrace{\frac{(x + q)^2}{2}}_{\dot{v}} \cdot \underbrace{f''(x)}_{d\dot{u}} dx \\ &= q \cdot f(0) - \frac{q^2}{2} f'(0) + \int_{-q}^0 \frac{(x + q)^2}{2} f''(x) dx \end{aligned}$$



$$\int_{-q}^0 f(x) dx = \underbrace{q \cdot f(0)}_{\text{approximate}} - \underbrace{\frac{q^2}{2} \cdot f'(0)}_{\pm \text{error}} + \int_{-q}^0 \frac{(x+q)^2}{2} f''(x) dx$$


exact
approximate
± error



$$\int_{-q}^0 f(x) dx = q \cdot f(0) - \frac{q^2}{2} f'(0) + \int_{-q}^0 \frac{(x+q)^2}{2} f''(x) dx$$

$$\int_0^q f(x) dx = q \cdot f(0) + \frac{q^2}{2} \cdot f'(0) + \int_0^q \frac{(x-q)^2}{2} f''(x) dx$$

$$\int_{-q}^q f(x) dx = 2q \cdot f(0) + \int_{-q}^0 \frac{(x+q)^2}{2} f''(x) dx + \int_0^q \frac{(x-q)^2}{2} f''(x) dx$$



$$\int_{-q}^q f(x) dx = \underbrace{2q \cdot f(0)}_{\text{approximate}} + \underbrace{\int_{-q}^0 \frac{(x+q)^2}{2} f''(x) dx + \int_0^q \frac{(x-q)^2}{2} f''(x) dx}_{\pm \text{error}}$$

exact
approximate
± error

We re-arrange to write the **error** as the difference between the **actual** area of one slice and its rectangular **approximation**.

$$\int_{-q}^q f(x) dx - 2q \cdot f(0) = \int_{-q}^0 \frac{(x+q)^2}{2} f''(x) dx + \int_0^q \frac{(x-q)^2}{2} f''(x) dx$$

$$\text{error} = \left| \int_{-q}^0 \frac{(x+q)^2}{2} f''(x) dx + \int_0^q \frac{(x-q)^2}{2} f''(x) dx \right|$$


$$\leq \left| \int_{-q}^0 \frac{(x+q)^2}{2} f''(x) dx \right| + \left| \int_0^q \frac{(x-q)^2}{2} f''(x) dx \right|$$

$$\leq \int_{-q}^0 \frac{(x+q)^2}{2} M dx + \int_0^q \frac{(x-q)^2}{2} M dx$$

$$= M \left[\frac{(x+q)^3}{6} \right]_{-q}^0 + M \left[\frac{(x-q)^3}{6} \right]_0^q$$


$$= \frac{M \cdot q^3}{3}$$

Now we can bound the error of a single slice:




$$\left| \int_{-q}^q f(x) dx - 2q \cdot f(0) \right| \leq \frac{M}{3} \cdot q^3$$

$$\left| \int_{x_{i-1}}^{x_i} f(x) dx - \frac{b-a}{n} \cdot f(\bar{x}_i) \right| \leq \frac{M}{3} \left(\frac{b-a}{2n} \right)^3 = \frac{M(b-a)^3}{24n^3}$$




- ▶ The error in each slice is at most $\frac{M(b-a)^3}{24n^3}$
- ▶ There are n slices
- ▶ The overall error is at most $n \cdot \frac{M(b-a)^3}{24n^3} = \frac{M(b-a)^3}{24n^2}$

TABLE OF CONTENTS




An integral is **improper** if one or both of the following happen:

- ▶ The region of integration is unbounded, e.g. $\int_1^{\infty} \frac{\sin x}{x} dx$



$$\Delta x = \frac{b-a}{n} = \frac{\infty}{n} ???$$

- ▶ The integrand is unbounded over the interval, e.g. $\int_{-1}^1 \frac{1}{x^2} dx$



$$f(0)\Delta x = ???$$

Strategy

In both cases, we eliminate the offending parts of the integral using limits.

$$\int_1^{\infty} \frac{\sin x}{x} dx =$$

$$\int_0^3 \frac{1}{x} dx =$$

If the limit doesn't exist, we say the integral **diverges**. Otherwise it **converges**.


$$\int_1^{\infty} \frac{1}{x} dx =$$

$$\int_1^{\infty} \frac{1}{x^2} dx =$$

Evaluate $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$

When an integral has multiple sources of impropriety, we break it up into integrals that have only one source each. If all of them converge, the original integral converges. If any of them diverges, the original integral diverges as well.

Evaluate $\int_0^1 \frac{1}{2\sqrt{x}} dx$



365/643

Evaluate $\int_{-2}^1 \frac{1}{x^2} dx$

366/643

Evaluate $\int_0^\infty \frac{\cos x}{1 + \sin^2 x} dx$, or show that it diverges.

367/643

WARNING: SNEAKY DIVERGENCE

If you don't realize that an integral diverges, you can generate answers that look plausible but are secretly nonsense.

For example, attempting to use the Fundamental Theorem of Calculus in the example $\int_{-2}^1 \frac{1}{x^2} dx$ gives $\left[-\frac{1}{x}\right]_{-2}^1 = -\frac{3}{2}$: a poor approximation for positive infinity!

368/643

Evaluate $\int_0^1 \frac{1}{x^p} dx$ and $\int_1^\infty \frac{1}{x^p} dx$ when p is constant.

p-test


Let p be a constant.

If $p < 1$, then $\int_0^1 \frac{1}{x^p} dx$ converges

If $p \geq 1$, then $\int_0^1 \frac{1}{x^p} dx$ diverges

If $p > 1$, then $\int_1^\infty \frac{1}{x^p} dx$ converges

If $p \leq 1$, then $\int_1^\infty \frac{1}{x^p} dx$ diverges



$\int_0^1 \frac{1}{x} dx$ diverges

$\int_1^\infty \frac{1}{x} dx$ diverges

Decide whether each integral converges or diverges.

▶ $\int_0^1 \frac{1}{x^{1/3}} dx$

▶ $\int_0^1 \frac{1}{\sqrt{x}} dx$

▶ $\int_0^1 \frac{1}{x} dx$

▶ $\int_0^1 \frac{1}{x^{1.5}} dx$

▶ $\int_1^\infty \frac{1}{x^{1/3}} dx$


▶ $\int_1^\infty \frac{1}{\sqrt{x}} dx$

▶ $\int_1^\infty \frac{1}{x} dx$

▶ $\int_1^\infty \frac{1}{x^{1.5}} dx$

Theorem 1.12.20

Let a and c be real numbers with $a < c$ and let the function $f(x)$ be continuous for all $x \geq a$. Then the improper integral $\int_a^\infty f(x) dx$ converges if and only if the improper integral $\int_c^\infty f(x) dx$ converges.



373/643

Decide whether each integral converges or diverges.

$$\blacktriangleright \int_0^9 \frac{1}{x^{0.3}} dx$$

$$\blacktriangleright \int_0^{81} \frac{1}{x^2} dx$$

$$\blacktriangleright \int_0^{\frac{1}{2}} \frac{1}{x^3} dx$$

$$\blacktriangleright \int_{15}^\infty \frac{1}{x^{0.3}} dx$$

$$\blacktriangleright \int_{0.4}^\infty \frac{1}{x^2} dx$$

$$\blacktriangleright \int_{\frac{1}{2}}^\infty \frac{1}{x^3} dx$$

374/643

It is very common to encounter integrals that are too complicated to evaluate explicitly. Numerical approximation schemes, evaluated by computer, are often used instead. You want to be sure that at least the integral converges before feeding it into a computer.



Fortunately it is usually possible to determine whether or not an improper integral converges even when you cannot evaluate it explicitly.

375/643

Comparison

Let a be a real number. Let f and g be functions that are defined and continuous for all $x \geq a$ and assume that $g(x) \geq 0$ for all $x \geq a$.

- If $|f(x)| \leq g(x)$ for all $x \geq a$ and if $\int_a^\infty g(x) dx$ converges, then $\int_a^\infty f(x) dx$ converges.
- If $f(x) \geq g(x)$ for all $x \geq a$ and if $\int_a^\infty g(x) dx$ diverges, then $\int_a^\infty f(x) dx$ diverges.



376/643 Theorem 1.12.17

Does the integral $\int_1^{\infty} e^{-x^2}$ converge or diverge?

Let functions $f(x)$ and $g(x)$ be positive and continuous for all $x \geq a$.

		$\int_a^{\infty} g(x) dx$ co
	$f(x) \leq g(x)$ for all $x \geq a$	
	$f(x) \geq g(x)$ for all $x \geq a$	

For each example below, decide whether the statement is a valid use of the comparison theorem.

▶ $\int_1^{\infty} \frac{1}{x^2} dx$ converges and $0 \leq \frac{1}{x^2} \leq \frac{2+\sin x}{x^2}$ for $x \geq 1$. So by the comparison test, $\int_1^{\infty} \frac{2+\sin x}{x^2} dx$ converges as well.

▶ $\int_1^{\infty} \frac{1}{x^2} dx$ converges and $0 \leq \frac{e^{-x}}{x^2} \leq \frac{1}{x^2}$ for $x \geq 1$. So by the comparison test, $\int_1^{\infty} \frac{e^{-x}}{x^2} dx$ converges as well.

▶ $\int_1^{\infty} \frac{1}{x^2} dx$ converges and $-\frac{1}{x} \leq \frac{1}{x^2}$ for $x \geq 1$. So by the comparison test, $\int_1^{\infty} \frac{-1}{x} dx$ converges as well.



Limiting comparison


Let $-\infty < a < \infty$. Let f and g be functions that are defined and continuous for all $x \geq a$ and assume that $g(x) \geq 0$ for all $x \geq a$. If the limit

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)}$$

exists and is nonzero, then either $\int_a^{\infty} f(x) dx$ and $\int_a^{\infty} g(x) dx$ both converge, or they both diverge.

Use limiting comparison to determine whether $\int_1^{\infty} \frac{1}{x+10} dx$ converges or diverges.

TABLE OF CONTENTS



381/643

HELPFUL UNITS

- ▶ Force is measured in units of newtons, with $1 \text{ N} = 1 \frac{\text{kg} \cdot \text{m}}{\text{s}^2}$.
- ▶ From its units, we see force looks like (mass) \times (acceleration)
- ▶ Work is measured in units of joules, with $1 \text{ J} = 1 \frac{\text{kg} \cdot \text{m}^2}{\text{s}^2}$
- ▶ From its units, we see work looks like (force) \times (distance)

382/643

Intuition

Work, in physics, is a way of quantifying the amount of energy that is required to act against a force.

For example:

- ▶ An object on the ground is subject to gravity. The force acting on the object is

$$m \cdot g$$

where m is the mass of the object (here, we're using kilograms), and g is the standard acceleration due to gravity (about $9.8 \frac{\text{kg} \cdot \text{m}}{\text{s}^2}$ on Earth).

- ▶ When you lift an object in the air, you are acting against that force. How much work you have to do depends on how strong the force is (how much mass the object has, and how strong gravity is) and also how far you lift it.

383/643

Work

The work done by a force $F(x)$ in moving an object from $x = a$ to $x = b$ is

$$W = \int_a^b F(x) \, dx$$

In particular, if the force is a constant F , the work is $F \cdot (b - a)$.


(For motivation of this definition, see Section 2.1 in the CLP-2 text.)

We saw the force of gravity on an object of mass m kg is $m \cdot g$ N. So to lift such an object a distance of y metres requires work of


$$m \cdot g \cdot y \text{ J}$$

384/643 Definition 2.1.1

A cable dangles in a hole. The cable is 10 metres long, and has a mass of 5 kg. Its density is constant. How much work is done to pull the cable out of the hole?




- ▶ A piece of the cable near the top of the hole isn't lifted very far.
- ▶ A piece of the cable near the bottom of the hole is lifted farther.
- ▶ Consider a small piece of cable starting y metres from the top.



A cylinder is filled with a liquid that we will pump out the top.


- ▶ To pump out a molecule from the top of the container, we don't have to work against gravity for very far.
- ▶ To pump out a molecule from the bottom of the container, we have to work against gravity for a longer distance.



- ▶ Every molecule at the same height has the same distance to travel to reach the top of the container. So, we'll chop up the tank into thin horizontal slices.

Hooke's Law

When a (linear) spring is stretched (or compressed) by x units beyond its natural length, it exerts a force of magnitude kx , where the constant k is the spring constant of that spring.



Suppose we want to stretch a string from a units beyond its natural length to b units beyond its natural length. The force of the spring at position x is kx , for some constant k . So, the work required is:

A spring has a natural length of 0.1 m. If a 12 N force is needed to keep it stretched to a length of 0.12 m, how much work is required to stretch it from 0.12 m to 0.15 m?






TABLE OF CONTENTS



$$\text{Average} \approx \frac{f(x_1) + \cdots + f(x_n)}{n}$$

$$\begin{aligned} \text{Average} &= \lim_{n \rightarrow \infty} \left[\frac{1}{n} \sum_{i=1}^n f(x_i) \right] = \lim_{n \rightarrow \infty} \left[\frac{(b-a)}{(b-a)n} \sum_{i=1}^n f(x_i) \right] \\ &= \lim_{n \rightarrow \infty} \left[\frac{1}{b-a} \sum_{i=1}^n f(x_i) \Delta x \right] = \frac{1}{b-a} \int_a^b f(x) dx \end{aligned}$$

Average

Let $f(x)$ be an integrable function defined on the interval $a \leq x \leq b$. The average value of f on that interval is

$$f_{\text{ave}} = \frac{1}{b-a} \int_a^b f(x) dx$$


The temperature in a certain city at time t (measured in hours past midnight) is given by

$$T(t) = t - \frac{t^2}{30}$$

What was the average temperature of one day (from $t = 0$ to $t = 24$)?




Let's check that our answer makes some intuitive sense.



Since the temperature is always between 0 and 8, we expect the average to be between 0 and 8

393/643


Let's also recall the motivation for our definition



394/643

Find the average value of the function $f(x) = \frac{e^x}{e^{2x} + 1}$ over the interval

$$\left[0, \frac{\log 3}{2}\right].$$




395/643

AVERAGE VELOCITY

Let $x(t)$ be the position at time t of a car moving along the x -axis. The velocity of the car at time t is the derivative $v(t) = x'(t)$. The average velocity of the car over the time interval $a \leq t \leq b$ is:


396/643 Example 2.2.5

TABLE OF CONTENTS




Centre of Mass

If you support a body at its centre of mass (in a uniform gravitational field) it balances perfectly. That's the definition of the centre of mass of the body.



If you support a body at its centre of mass (in a uniform gravitational field) it balances perfectly. That's the definition of the centre of mass of the body.




If the body consists of a finite number of masses m_1, \dots, m_n attached to an infinitely strong, weightless (idealized) rod with mass number i attached at position x_i , then the center of mass is at the (weighted) average value of x :

$$\bar{x} = \frac{\sum_{i=1}^n m_i x_i}{\sum_{i=1}^n m_i}$$

The denominator $m = \sum_{i=1}^n m_i$ is the total mass of the body.


An idealized (weightless, unbending) rod has small masses attached to it at the following locations:

- ▶ 1 kg at $x = 1$ metre from the left end
- ▶ 4 kg at $x = 3$ metres from the left end
- ▶ 2 kg at $x = 6$ metres from the left end
- ▶ 1 kg at $x = 7$ metres from the left end




What is the location of its centre of mass?

We can also group the masses, and treat them as single points of mass at their centres of gravity, without affecting the centre of gravity of the entire object.



401/643

Sometimes we can simplify a physical calculation by treating an object as a point particle located at its centre of mass. When we were learning about work, we found the following:



A cable dangles in a hole. The cable is 10 metres long, and has a mass of 5 kg. Its density is constant. We found that the work required to pull the cable out of the hole was

$$25g J$$


where g is the acceleration due to gravity.

402/643

Consider a metre-long rod that is denser on one end than the other, with density

$$\rho(x) = (2x + 1) \frac{\text{kg}}{\text{m}}$$

at a position x metres from its left end.



What is its centre of mass?

403/643

If a body consists of mass distributed continuously along a straight line, say with mass density $\rho(x)$ kg/m and with x running from a to b , rather than consisting of a finite number of point masses, the formula for the center of mass becomes

$$\bar{x} = \frac{\int_a^b x \rho(x) dx}{\int_a^b \rho(x) dx}$$




Think of $\rho(x) dx$ as the mass of the "almost point particle" between x and $x + dx$.

404/643 Equation 2.3.2

Centre of Mass


If you support a body at its centre of mass (in a uniform gravitational field) it balances perfectly. That's the definition of the center of mass of the body.

Centre of mass isn't just for linear solids: it applies to 2- and 3-dimensional objects as well.



405/643


Consider a flat metal plate of uniform density, whose shape is the area below curve $y = T(x)$ and above curve $y = B(x)$, from $x = a$ to $x = b$.



The centre of mass will be a point in the xy -plane, (\bar{x}, \bar{y}) . To find \bar{x} and \bar{y} , we will treat vertical slices as point particles.

406/643

Consider a flat metal plate of uniform density, whose shape is the area below curve $y = T(x)$ and above curve $y = B(x)$, from $x = a$ to $x = b$.




If ρ is the density of the plate, so that a slice of width dx and height $h = T(x) - B(x)$ has mass $\rho h dx = \rho(T(x) - B(x)) dx$, then:

$$\begin{aligned}\bar{x} &= \frac{\int_a^b \rho(T(x) - B(x)) \cdot x dx}{\int_a^b \rho(T(x) - B(x)) dx} \\ &= \frac{\int_a^b (T(x) - B(x)) \cdot x dx}{\int_a^b (T(x) - B(x)) dx}\end{aligned}$$

The centre of mass will be a point in the xy -plane, (\bar{x}, \bar{y}) . To find \bar{x} and \bar{y} , we will treat vertical slices as point particles.


407/643

To find \bar{y} , note that the y -coordinate of the centre of mass of a slice that is almost a rectangle, and has uniform density, will be halfway up the slice, at $\frac{T(x)+B(x)}{2}$.




408/643

To find \bar{y} , note that the y -coordinate of the centre of mass of a slice that is almost a rectangle, and has uniform density, will be halfway up the slice, at $\frac{T(x)+B(x)}{2}$.



Find the centre of mass (centroid) of the quarter circular unit disk $x \geq 0, y \geq 0, x^2 + y^2 \leq 1$.



Find the centre of mass (centroid) of a plate of uniform density in the shape of the finite area enclosed by the functions $y = T(x) = 2 - x$ and $y = B(x) = x^2$.





TABLE OF CONTENTS



Differential Equation

A **differential equation** is an equation for an unknown function that involves the derivative of the unknown function.

Differential equations play a central role in modelling a huge number of different phenomena. Here is a table giving a bunch of named differential equations and what they are used for. It is far from complete.

Newton's Law of Motion	describes motion of particles
Maxwell's equations	describes electromagnetic radiation
Navier-Stokes equations	describes fluid motion
Heat equation	describes heat flow
Wave equation	describes wave motion
Schrödinger equation	describes atoms, molecules and crystals
Stress-strain equations	describes elastic materials
Black-Scholes models	used for pricing financial options
Predator-prey equations	describes ecosystem populations
Einstein's equations	connects gravity and geometry
Ludwig-Jones-Holling's equation	models spruce budworm/Balsam fir ecosystem
Zeeman's model	models heart beats and nerve impulses
Sherman-Rinzel-Keizer model	for electrical activity in Pancreatic β -cells
Hodgkin-Huxley equations	models nerve action potentials

413/643

Disclaimer:

We are dipping our toes into a vast topic. Most universities offer half a dozen different undergraduate courses on various aspects of differential equations. We will just look at one special, but important, type of equation.

- ▶ We will first learn to **verify** solutions without **finding** them. (If you learned about differential equations last semester, this will be review.)
- ▶ **Then**, we will learn to solve one particular type of differential equation.

414/643

DIFFERENTIAL EQUATIONS

Definition

A **differential equation** is an equation involving the derivative of an unknown function.

Examples: $\frac{dy}{dx} = 2x$; $x\frac{dy}{dx} = 7xy + y$

Definition

If a **function** makes a differential equation true, we say it **satisfies** the differential equation, or is a solution to the differential equation.

Example: $y = x^2$ and $y = x^2 + 1$ both satisfy the first differential equation

415/643

VERIFYING SOLUTIONS

Consider the equation

$$x + 2 = x^3 - x^2$$

How would you verify whether $x = 1$ satisfies the equation?

How would you verify whether $x = 2$ satisfies the equation?

Plug x into the equation, check whether the left-hand side and the right-hand side are the same **number.**

416/643

VERIFYING SOLUTIONS

Consider the differential equation

$$\frac{dy}{dx} = 2y + 4x$$

How would you verify whether $y = e^{2x} - 2x$ satisfies the equation?

How would you verify whether $y = e^{2x} - 2x - 1$ satisfies the equation?

Replace y and $\frac{dy}{dx}$ in the equation, check whether the left-hand side and the right-hand side are the same **function**.

417/643

Differential equation:

$$x \frac{dy}{dx} = 7xy + y$$

Interpretation:

There is a function $y(x)$ that makes the left-hand side and the right-hand side into the same function.

To check whether a given function satisfies the differential equation, plug it in for everything with a “ y ”: y itself and $\frac{dy}{dx}$.

Is $y = xe^{7x+9}$ a solution to the differential equation?

418/643

Which of the following solve the differential equation $\frac{dy}{dx} = \frac{x}{y}$?

A. $y = -x$

B. $y = x + 5$

C. $y = \sqrt{x^2 + 5}$

419/643

FIRST EXAMPLE OF A SEPARABLE DE

Definition

A separable differential equation is an equation for a function $y(x)$ that can be written in the form

$$g(y) \cdot \frac{dy}{dx} = f(x)$$

(It may take some rearranging to get the equation into this form.)

For example:

$$y^2 \cdot \frac{dy}{dx} = 4x$$

420/643

GENERAL METHOD FOR SOLVING SEPARABLE DES

$$g(y) \cdot \frac{dy}{dx} = f(x)$$

421/643

GENERAL METHOD FOR SOLVING SEPARABLE DES

$$g(y) \cdot \frac{dy}{dx} = f(x)$$

Shorthand:

$$g(y) \cdot \frac{dy}{dx} = f(x)$$

$$g(y) dy = f(x) dx$$

$$\int g(y) dy = \int f(x) dx$$

422/643

$$\frac{dy}{dx} = y^2 x$$

1. "Separate" y 's from x 's.
2. Integrate.
3. Solve explicitly for y .



423/643

$$\frac{dy}{dx} = (xy)^4, \quad y(0) = \frac{1}{2}$$



424/643

$$\frac{dy}{dx} = y(4x^3 - 1) \quad y(0) = -2$$



Let a and b be any two constants. We'll now solve the family of differential equations

$$\frac{dy}{dx} = a(y - b)$$

using our mnemonic device.

Linear First-Order Differential Equations

Let a and b be constants. The differentiable function $y(x)$ obeys the differential equation

$$\frac{dy}{dx} = a(y - b)$$

if and only if

$$y(x) = \{y(0) - b\} e^{ax} + b$$

Find a function $y(x)$ with $y' = 3y + 7$ and $y(2) = 5$.



The rate at which a medicine is metabolized (broken down) in the body depends on how much of it is in the bloodstream. Suppose a certain medicine is metabolized at a rate of $\frac{1}{10}A$ $\mu\text{g/hr}$, where A is the amount of medicine in the patient. The medicine is being administered to the patient at a constant rate of 2 $\mu\text{g/hr}$. If the patient starts with no medicine in their blood at $t = 0$, give the formula for the amount of medicine in the patient at time t . What happens to the amount over time?



Linear First-Order Differential Equations

Let a and b be constants. The differentiable function $y(x)$ obeys the differential equation

$$\frac{dy}{dx} = a(y - b)$$


if and only if

$$y(x) = \{y(0) - b\} e^{ax} + b$$


$$\frac{dA}{dt} = 2 - \frac{1}{10}A = -\frac{1}{10}(A - 20) \quad A(0) = 0$$

429/643



TABLE OF CONTENTS



430/643



431/643



432/643

RADIOACTIVE DECAY

One model for radioactive decay says that the rate at which an isotope decays is proportional to the amount present. So if $Q(t)$ is the amount of a radioactive substance, then

$$\frac{dQ}{dt} = -kQ(t)$$

for some constant¹ k .

This is a first-order linear differential equation. Its explicit solutions have the form:

¹By including the negative sign, we ensure k will be positive, but of course we could also write " $\frac{dQ}{dt} = KQ(t)$ for some [negative] constant K ".

433/643

HALF-LIFE

The **half-life** of an isotope is the time required for half of that isotope to decay. If we know the half-life of a substance is $t_{1/2}$, and its quantity at time t is given by $Q(0)e^{-kt}$ we can find the constant k :

434/643

Radioactive Decay

The function $Q(t)$ satisfies the equation $\frac{dQ}{dt} = -kQ(t)$ if and only if

$$Q(t) = Q(0)e^{-kt}$$

The half-life is defined to be the time $t_{1/2}$ which obeys $Q(t_{1/2}) = \frac{1}{2}Q(0)$. The half-life is related to the constant k by $t_{1/2} = \frac{\log 2}{k}$. Then

$$Q(t) = Q(0)e^{-\frac{\log 2}{t_{1/2}}t} = Q(0) \cdot \left(\frac{1}{2}\right)^{\frac{t}{t_{1/2}}}$$

If the half-life of ^{14}C is $t_{1/2} = 5730$ years, then the quantity of carbon-14 present in a sample after t years is:

435/643 Corollary 2.4.9

A particular piece of flax parchment contains about 64% as much ^{14}C as flax plants do today. We will estimate the age of the parchment, using 5730 years as the half-life of ^{14}C .


First, a rough estimate: is the parchment older or younger than 5730 years?

436/643 Example 2.4.10

Newton's law of cooling

The rate of change of temperature of an object is proportional to the difference in temperature between the object and its surroundings.

The temperature of the surroundings is sometimes called the ambient temperature.



$$\frac{dT}{dt} =$$

437/643 Equation 2.4.4

Linear First-Order Differential Equations

Let a and b be constants. The differentiable function $y(x)$ obeys the differential equation

$$\frac{dy}{dx} = a(y - b)$$

if and only if

$$y(x) = \{y(0) - b\} e^{ax} + b$$

Find an explicit formula for functions $T(t)$ solving the differential equation $\frac{dT}{dt} = K(T(t) - A)$ for some constants K and A .

438/643

The temperature of a glass of iced tea is initially 5° . After 5 minutes, the tea has heated to 10° in a room where the air temperature is 30° . Assume the temperature of the tea as it cools follows Newton's law of cooling,

$$T(t) = (T(0) - A) e^{Kt} + A$$

- Determine the temperature as a function of time.
- When the tea will reach a temperature of 14° ?



439/643 Example 2.4.12

A glass of room-temperature water is carried out onto a balcony from an apartment where the temperature is 22°C . After one minute the water has temperature 26°C and after two minutes it has temperature 28°C . Assuming the water warms according to Newton's law of cooling, what is the outdoor temperature? Assume that the temperature of the water obeys Newton's law of cooling.



440/643 Example 2.4.14

Let P be the size of a population, and let K be the carrying capacity of its environment (i.e. the population size that can be sustainably supported).

When P is much less than K , our population has...

- A. not enough resources
- B. just enough resources
- C. extra resources

So when the P is much less than K , we expect the population to...

- A. shrink
- B. stay the same
- C. grow

Malthusian growth

The Malthusian growth model relates population growth to population size:

$$\frac{dP}{dt} = bP(t)$$

where b is a constant representing net birthrate per member of the population.

Let P be the size of a population, and let K be the carrying capacity of its environment (i.e. the population size that can be sustainably supported).

When P is **greater** than K , our population has...

- A. not enough resources
- B. just enough resources
- C. extra resources

So when the P is greater than K , we expect the population to...

- A. shrink
- B. stay the same
- C. grow

Logistic growth models population growth as:

$$\frac{dP}{dt} = b_0 \left(1 - \frac{P(t)}{K}\right) P(t)$$


- ▶ If $P \ll K$, then $\frac{dP}{dt} \approx$
- ▶ If $P \approx K$, then $\frac{dP}{dt} \approx$
- ▶ If $P > K$, then $\frac{dP}{dt} \approx$



Before we solve explicitly, let's sketch some solutions to

$$\frac{dP}{dt} = b_0 \left(1 - \frac{P(t)}{K}\right) P(t)$$

- ▶ If $P(a) = 0$:
- ▶ If $0 < P(a) < K$:
- ▶ If $P(a) = K$:
- ▶ If $K < P(0)$:



Find the explicit solutions to


$$\frac{dP}{dt} = b \left(1 - \frac{P(t)}{K} \right) P(t)$$

when b and K are constants.

445/643

At time $t = 0$, where t is measured in minutes, a large tank contains 3 litres of water in which 1 kg of salt is dissolved. Fresh water enters the tank at a rate of 2 litres per minute and the fully mixed solution leaks out of the tank at the varying rate of $2t$ litres per minute.

- Determine the volume of solution $V(t)$ in the tank at time t .
- Determine the amount of salt $Q(t)$ in solution when the amount of water in the tank is at maximum.




446/643 Example 2.4.17

SETTLING TANK

A settling tank is filled with 100,000 litres of pure water. Every hour, 1,000 litres of water, containing 3 grams of pollutants, enters the tank.

90% of the pollutants in the settling tank sink to the bottom, with the remaining 10% well-mixed into the water. The tank drains 1,000 litres of this mixed water into the sewer every hour.

In order to drain the water into the local sewer, the concentration of pollutants cannot be more than 1 gram per 1,000 litres. How long can the settling tank take dirty water until the process must be stopped?



447/643

448/643

Let $P(t)$ be the total amount (in grams) of pollutants in the tank. Pollutants are entering at a rate of 3 grams per hour. How fast are they leaving?

So, the quantity of pollutants in the tank satisfies the differential equation:

You deposit $\$P$ in a bank account at time $t = 0$, and the account pays $r\%$ interest per year, compounded n times per year. Your balance at time t is $B(t)$.

If one interest payment comes at time t , then the next interest payment will be made at time $t + \frac{1}{n}$ and will be:

$$\frac{1}{n} \times \frac{r}{100} \times B(t) = \frac{r}{100n} B(t)$$

So, calling $\frac{1}{n} = h$,

$$B(t+h) = B(t) + \frac{r}{100} B(t)h \quad \text{or} \quad \frac{B(t+h) - B(t)}{h} = \frac{r}{100} B(t)$$

If the interest is compounded continuously,

$$\begin{aligned} \frac{dB}{dt}(t) &= \lim_{h \rightarrow 0} \frac{B(t+h) - B(t)}{h} = \frac{r}{100} B(t) \\ \implies B(t) &= B(0) \cdot e^{rt/100} = P \cdot e^{rt/100} \end{aligned}$$


Continuously compounding interest

If an account with balance $B(t)$ pays a continuously compounding rate of $r\%$ per year, then:

$$\begin{aligned} \frac{dB}{dt} &= \frac{r}{100} B \\ B(t) &= B(0) \cdot e^{rt/100} \end{aligned}$$

You invest $\$200\,000$ into an account with continuously compounded interest of 5% annually. You want to withdraw from the account continuously at a rate of $\$W$ per year, for the next 20 years. How big can W be?

TABLE OF CONTENTS



453/643

We can imagine the list of numbers below carrying on forever:

$$\begin{aligned}a_1 &= 0.1 \\a_2 &= 0.01 \\a_3 &= 0.001 \\a_4 &= 0.0001 \\a_5 &= 0.00001 \\&\vdots\end{aligned}$$

A **sequence** is a list of infinitely many numbers with a specified order. It is denoted $\{a_1, a_2, \dots, a_n, \dots\}$ or $\{a_n\}_{n=1}^{\infty}$, etc.

Imagine *adding up* this sequence of numbers.

A **series** is a sum $a_1 + a_2 + \dots + a_n + \dots$ of infinitely many terms.

454/643

To handle sequences and series, we should define them more carefully. A good definition should allow us to answer some basic questions, such as:

- ▶ What does it mean to add up infinitely many things?
- ▶ Should infinitely many things add up to an infinitely large number?
- ▶ Does the order in which the numbers are added matter?
- ▶ Can we add up infinitely many functions, instead of just infinitely many numbers?

455/643

Sequence

A **sequence** is a list of infinitely many numbers with a specified order.

Some examples of sequences:

- ▶ $\{1, 2, 3, 4, 5, 6, 7, 8, \dots\}$ (natural numbers)
- ▶ $\{3, 1, 4, 1, 5, 9, 2, 6, \dots\}$ (digits of π)
- ▶ $\{1, -1, 1, -1, 1, \dots\}$ (powers of -1 : $(-1)^0, (-1)^1, (-1)^2$, etc.)

456/643

SEQUENCE NOTATION

Sequence

A **sequence** is a list of infinitely many numbers with a specified order. It is denoted $\{a_1, a_2, a_3, \dots, a_n, \dots\}$ or $\{a_n\}$ or $\{a_n\}_{n=1}^{\infty}$, etc.

$$\{a_n\}_{n=1}^{\infty} = \left\{ \frac{1}{n} \right\}_{n=1}^{\infty}$$

- ▶ $n = 1$: this is the index of the first term of our sequence. Sometimes it's 0, sometimes something else, for example a year.
- ▶ ∞ : there is no end to our sequence.
- ▶ $\frac{1}{n}$: this tells us the value of a_n .
- ▶ Often we omit the limits and even the brackets, writing $a_n = \frac{1}{n}$.

457/643

For convenience, we write a_1 for the first term of a sequence, a_2 for the second term, etc.

In the sequence $1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots$, a_3 is another name for

Sometimes we can find a rule for a sequence. In the above sequence, $a_n =$

We can write $\{a_n\}_{n=1}^{\infty} =$

458/643

Our primary concern with sequences will be the behaviour of a_n as n tends to infinity and, in particular, whether or not a_n "settles down" to some value as n tends to infinity.

Convergence

A sequence $\{a_n\}_{n=1}^{\infty}$ is said to **converge** to the limit A if a_n approaches A as n tends to infinity. If so, we write

$$\lim_{n \rightarrow \infty} a_n = A \quad \text{or} \quad a_n \rightarrow A \text{ as } n \rightarrow \infty$$

A sequence is said to converge if it converges to some limit. Otherwise it is said to diverge.

459/643 Definition 3.1.3

Convergence

A sequence $\{a_n\}_{n=1}^{\infty}$ is said to **converge** to the limit A if a_n approaches A as n tends to infinity. If so, we write

$$\lim_{n \rightarrow \infty} a_n = A \quad \text{or} \quad a_n \rightarrow A \text{ as } n \rightarrow \infty$$


A sequence is said to converge if it converges to some limit. Otherwise it is said to diverge.

- ▶ $\{1, 2, 3, 4, 5, 6, 7, 8, \dots\}$ (natural numbers)
This sequence
- ▶ $\{3, 1, 4, 1, 5, 9, 2, 6, \dots\}$ (digits of π)
This sequence
- ▶ $\{1, -1, 1, -1, 1, \dots\}$ (powers of -1 : $(-1)^0, (-1)^1, (-1)^2$, etc.)
This sequence

460/643

Does the sequence $a_n = \frac{n}{2n+1}$ converge or diverge?

Consider the sequence $a_n = \frac{1}{3^n + 1}$. $\lim_{n \rightarrow \infty} a_n =$



Theorem 3.1.6


If $\lim_{x \rightarrow \infty} f(x) = L$
and if $a_n = f(n)$ for all positive integers n , then

$$\lim_{n \rightarrow \infty} a_n = L$$

CAUTIONARY TALE


Consider the sequence $b_n = \sin(\pi n) =$

$$\lim_{n \rightarrow \infty} b_n = \lim_{x \rightarrow \infty} f(x)$$



Theorem

If $\lim_{x \rightarrow \infty} f(x) = L$ and if $a_n = f(n)$ for all natural n , then $\lim_{n \rightarrow \infty} a_n = L$.



Arithmetic of Limits

Let A , B and C be real numbers and let the two sequences $\{a_n\}_{n=1}^{\infty}$ and $\{b_n\}_{n=1}^{\infty}$ converge to A and B respectively. That is, assume that

$$\lim_{n \rightarrow \infty} a_n = A \qquad \lim_{n \rightarrow \infty} b_n = B$$

Then the following limits hold.

- (a) $\lim_{n \rightarrow \infty} [a_n + b_n] = A + B$
- (b) $\lim_{n \rightarrow \infty} [a_n - b_n] = A - B$
- (c) $\lim_{n \rightarrow \infty} C a_n = C A$.
- (d) $\lim_{n \rightarrow \infty} a_n b_n = A B$
- (e) If $B \neq 0$, then $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{A}{B}$

Evaluate the following limits:

► $\lim_{n \rightarrow \infty} e^{-n} =$

► $\lim_{n \rightarrow \infty} \frac{1+n}{n} =$

► $\lim_{n \rightarrow \infty} \frac{1}{n^2} =$

► $\lim_{n \rightarrow \infty} 2n^2 =$

► $\lim_{n \rightarrow \infty} \left(\frac{1}{n^2}\right) (2n^2) =$

Continuous functions of limits

If $\lim_{n \rightarrow \infty} a_n = L$ and if the function $g(x)$ is continuous at L , then

$$\lim_{n \rightarrow \infty} g(a_n) = g(L)$$

Evaluate $\lim_{n \rightarrow \infty} \left[\sin \left(\frac{\pi n}{2n+1} \right) \right]$


Squeeze Theorem

If $a_n \leq c_n \leq b_n$ for all sufficiently large natural numbers n , and if

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = L$$

then

$$\lim_{n \rightarrow \infty} c_n = L$$




Evaluate

$$\lim_{n \rightarrow \infty} \left(\frac{2n + \cos n}{n + 1} \right)$$

Let $a_n = (-n)^{-n}$. Evaluate $\lim_{n \rightarrow \infty} a_n$.

TABLE OF CONTENTS



SEQUENCES AND SERIES

A **sequence** is a list of numbers
A **series** is the sum of such a list.


SEQUENCES AND SERIES

Sequence

List of numbers,
approaching

Series

Sum of numbers,
approaching



Square of Area 1

QUICK REVIEW: SIGMA NOTATION

Recall:

$$\sum_{n=1}^5 \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2}$$

We informally interpret:

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2} + \frac{1}{6^2} + \frac{1}{7^2} + \frac{1}{8^2} + \frac{1}{9^2} + \frac{1}{10^2} + \dots$$

(a more rigorous definition will be discussed soon)

Let a_n and b_n be sequences, and let C be a constant.

$$\sum_{n=1}^{\infty} (C \cdot a_n) =$$

A. $\sum_{n=1}^{\infty} C \cdot \sum_{n=1}^{\infty} a_n$

B. $\sum_{n=1}^{\infty} C + \sum_{n=1}^{\infty} a_n$

C. $C \sum_{n=1}^{\infty} a_n$

D. $a_n \sum_{n=1}^{\infty} C$

E. none of the above



Let a_n and b_n be sequences, and let C be a constant.

$$\sum_{n=1}^{\infty} (a_n + b_n) =$$

A. $\sum_{n=1}^{\infty} a_n \cdot \sum_{n=1}^{\infty} b_n$

B. $\sum_{n=1}^{\infty} a_n + \sum_{n=1}^{\infty} b_n$

C. $a_n + \sum_{n=1}^{\infty} b_n$

D. $a_n \sum_{n=1}^{\infty} b_n$

E. none of the above



Let a_n and b_n be sequences, and let C be a constant.

$$\sum_{n=1}^{\infty} (a_n)^C =$$

A. $\sum_{n=1}^{\infty} C \cdot \sum_{n=1}^{\infty} a_n$

B. $\left(\sum_{n=1}^{\infty} a_n \right)^C$

C. $C^n \sum_{n=1}^{\infty} a_n$

D. $\sum_{n=1}^{\infty} C(a_n)^{C-1}$

E. none of the above



SERIES PHILOSOPHY


What does it really mean to add up infinitely many things?

$$1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + \dots$$

We need an unambiguous definition.

HOW CAN WE ADD UP INFINITELY MANY THINGS?

SEQUENCE OF PARTIAL SUMS




$$S_1 = 0.2000$$

$$S_2 = 0.2400$$

$$S_3 = 0.2480$$

$$S_4 = 0.2496$$

$$S_5 = 0.2499$$



Partial sums let us think about series (sums) using the tools we've developed for sequences (lists).

$$a_1 = \frac{1}{5} = 0.2$$

$$S_1 = 0.2$$

$$a_2 = \frac{1}{5^2} = 0.04$$

$$S_2 = 0.24$$

$$a_3 = \frac{1}{5^3} = 0.008$$

$$S_3 = 0.248$$

$$a_4 = \frac{1}{5^4} = 0.0016$$

$$S_4 = 0.2496$$

$$a_5 = \frac{1}{5^5} = 0.00032$$

$$S_5 = 0.24992$$

We define $\sum_{n=1}^{\infty} a_n = \lim_{N \rightarrow \infty} \sum_{n=1}^N a_n = \lim_{N \rightarrow \infty} S_N$.


$a_1 = \frac{1}{5} = 0.2$	$S_1 = 0.2$	$a_5 = \frac{1}{5^5} = 0.00032$	$S_5 = 0.24992$
$a_2 = \frac{1}{5^2} = 0.04$	$S_2 = 0.24$	$a_6 = \frac{1}{5^6} = 0.000064$	$S_6 = 0.249984$
$a_3 = \frac{1}{5^3} = 0.008$	$S_3 = 0.248$	$a_7 = \frac{1}{5^7} = 0.0000128$	$S_7 = 0.2499968$
$a_4 = \frac{1}{5^4} = 0.0016$	$S_4 = 0.2496$	$a_8 = \frac{1}{5^8} = 0.00000256$	$S_8 = 0.24999936$

From the sequence of partial sums, we guess

$$\sum_{n=1}^{\infty} a_n = \lim_{N \rightarrow \infty} S_N =$$

481/643

NOTATION: $S_N = \sum_{n=1}^N a_n$



$$S_1 = a_1$$


$$S_2 = a_1 + a_2$$

$$S_3 = a_1 + a_2 + a_3$$

$$S_4 = a_1 + \dots + a_4$$

$$S_5 = a_1 + \dots + a_5$$

$$S_6 = a_1 + \dots + a_6$$



482/643

NOTATION PRACTICE

Suppose $\sum_{n=1}^{\infty} a_n$ has partial sums $S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$.


► Evaluate $\sum_{n=1}^{100} a_n$.

► Evaluate $\sum_{n=1}^{\infty} a_n$.



483/643

NOTATION PRACTICE




Andrew brings a plate of cookies to the professor's desk. When he puts them down, there are 10 cookies on the desk.

Then, Joel brings a plate of cookies. When he puts them down, there are 19 cookies on the desk.

How many cookies did each person bring?

484/643

NOTATION PRACTICE



Andrew brings a plate of cookies to the professor's desk. When he puts them down, there are 10 cookies on the desk.

Then, Joel brings a plate of cookies. When he puts them down, there are 19 cookies on the desk.

How many cookies did each person bring?

485/643

NOTATION PRACTICE

Suppose $\sum_{n=1}^{\infty} a_n$ has partial sums $S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$.


- ▶ Find a_1 .

- ▶ Give an explicit expression for a_n , when $n > 1$.



486/643

$$S_N = \sum_{n=1}^N a_n = \frac{N}{N+1}$$



$$S_1 = 1/(1+1)$$

$$S_2 = 2/(2+1)$$

$$S_3 = 3/(3+1)$$


$$S_4 = 4/(4+1)$$

$$S_5 = 5/(5+1)$$

$$S_6 = 6/(6+1)$$

$$S_7 = 7/(7+1)$$

$$S_8 = 8/(8+1)$$



487/643

Definition

The N^{th} partial sum of the series $\sum_{n=1}^{\infty} a_n$ is the sum of its first N terms

$$S_N = \sum_{n=1}^N a_n.$$

The partial sums form a sequence $\{S_N\}_{N=1}^{\infty}$. If this sequence of partial sums converges $S_N \rightarrow S$ as $N \rightarrow \infty$ then we say that the series $\sum_{n=1}^{\infty} a_n$ converges to S and we write

$$\sum_{n=1}^{\infty} a_n = S$$

If the sequence of partial sums diverges, we say that the series diverges.

488/643 Definition 3.2.3

Geometric Series

Let a and r be two fixed real numbers with $a \neq 0$. The series

$$a + ar + ar^2 + ar^3 + \dots$$

is called the **geometric series** with first term a and ratio r .

We call r the *ratio* because it is the quotient of consecutive terms:

$$\frac{ar^{n+1}}{ar^n} = r$$

Another useful way of identifying geometric series is to determine whether all pairs of consecutive terms have the same ratio.

- ▶ Geometric: $1 + \frac{1}{5} + \frac{1}{5^2} + \frac{1}{5^3} + \frac{1}{5^4} + \dots$
- ▶ Geometric: $\sum_{n=0}^{\infty} \frac{1}{2^n}$
- ▶ Not geometric: $1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \frac{1}{6} + \dots$

489/643

Consider the partial sum S_N of a geometric series:

$$S_N = a + ar + ar^2 + ar^3 + \dots + ar^N$$

$$rS_N =$$

$$rS_N - S_N =$$

$$S_N(r - 1) = ar^{N+1} - a$$

If $r \neq 1$, then

$$S_N = \frac{ar^{N+1} - a}{r - 1} = a \frac{r^{N+1} - 1}{r - 1}$$

490/643


Geometric Series and Partial Sums

Let a and r be constants with $a \neq 0$, and let N be a natural number.

- ▶ If $r \neq 1$, then $a + ar + ar^2 + ar^3 + \dots + ar^N = a \frac{r^{N+1} - 1}{r - 1}$.
- ▶ If $r = 1$, then $a + ar + ar^2 + ar^3 + \dots + ar^N =$
- ▶ If $|r| < 1$, then $\sum_{n=0}^{\infty} ar^n =$
- ▶ If $r = 1$, then $\sum_{n=0}^{\infty} ar^n =$
- ▶ If $r = -1$, then $\sum_{n=0}^{\infty} ar^n =$
- ▶ If $|r| > 1$, then $\sum_{n=0}^{\infty} ar^n =$

491/643 Example 3.2.4

$$\sum_{n=0}^{\infty} ar^n, r = 1, a \neq 0$$



$$S_0 = a$$


$$S_1 = 2a$$

$$S_2 = 3a$$

$$S_3 = 4a$$


$$S_4 = 5a$$

$$S_5 = 6a$$



492/643


$$\sum_{n=0}^{\infty} ar^n, r = -1, a \neq 0$$



$$\begin{aligned} S_0 &= a \\ S_1 &= 0 \\ S_2 &= a \\ S_3 &= 0 \\ S_4 &= a \\ S_5 &= 0 \end{aligned}$$

493/643


$$\sum_{n=0}^{\infty} ar^n, r > 1, a \neq 0$$



$$\begin{aligned} S_0 &= a \\ S_1 &= a \frac{r^2-1}{r-1} \\ S_2 &= a \frac{r^3-1}{r-1} \\ S_3 &= a \frac{r^4-1}{r-1} \\ S_4 &= a \frac{r^5-1}{r-1} \end{aligned}$$

494/643

$$\sum_{n=0}^{\infty} ar^n, r < -1, a \neq 0$$



$$\begin{aligned} S_0 &= a \\ S_1 &= a \frac{r^2-1}{r-1} \\ S_2 &= a \frac{r^3-1}{r-1} \\ S_3 &= a \frac{r^4-1}{r-1} \\ S_4 &= a \frac{r^5-1}{r-1} \\ S_5 &= a \frac{r^6-1}{r-1} \end{aligned}$$

495/643

GEOMETRIC SERIES

New bitcoins are produced when a particular type of computational problem is solved. Every time 210,000 solutions are found, the number of bitcoins each solution can produce is cut in half.


- ▶ Each of the first 210,000 solutions can produce 50 bitcoins.
- ▶ Each of the next 210,000 solutions can produce $\frac{50}{2}$ bitcoins.
- ▶ Each of the next 210,000 solutions can produce $\frac{50}{2^2}$ bitcoins.
- ▶ Each of the next 210,000 solutions can produce $\frac{50}{2^3}$ bitcoins.

Assume that this continues forever, and that bitcoins are infinitely divisible.² How many bitcoins can possibly be produced?

²Actually the smallest allowed division of a bitcoin is 10^{-8} .

496/643

$$\sum_{n=0}^{\infty} 210\,000 \left(\frac{50}{2^n}\right) = 21\,000\,000$$




$$S_0 = 10\,500\,000$$

$$S_1 = 15\,750\,000$$

$$S_2 = 18\,375\,000$$

$$S_3 = 19\,687\,500$$

$$S_4 = 20\,343\,750$$



Arithmetic of Series

Let S , T , and C be real numbers. Let the two series $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ converge to S and T respectively. Then

$$\sum_{n=1}^{\infty} [a_n + b_n] = S + T$$

$$\sum_{n=1}^{\infty} [a_n - b_n] = S - T$$

$$\sum_{n=1}^{\infty} [Ca_n] = CS$$

Geometric Series and Partial Sums

Let a and r be fixed numbers, and let N be a positive integer. Then

$$\sum_{n=0}^N ar^n = \begin{cases} a \cdot \frac{1-r^{N+1}}{1-r} & \text{if } r \neq 1 \\ a(N+1) & \text{if } r = 1 \end{cases}$$

so

$$\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r} \text{ provided } |r| < 1$$

Evaluate $\sum_{n=0}^{\infty} \left(\frac{2}{3^n} + \frac{4}{5^n}\right)$



Geometric Series and Partial Sums

Let a and r be fixed numbers, and let N be a positive integer. Then

$$\sum_{n=0}^N ar^n = \begin{cases} a \cdot \frac{1-r^{N+1}}{1-r} & \text{if } r \neq 1 \\ a(N+1) & \text{if } r = 1 \end{cases}$$

so

$$\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r} \text{ provided } |r| < 1$$

Evaluate $\sum_{n=6}^{\infty} \left(\frac{3^{n-1}}{5^{2n}}\right)$



Geometric Series and Partial Sums

Let a and r be fixed numbers, and let N be a positive integer. Then

$$\sum_{n=0}^N ar^n = \begin{cases} a \cdot \frac{1-r^{N+1}}{1-r} & \text{if } r \neq 1 \\ a(N+1) & \text{if } r = 1 \end{cases}$$

so

$$\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r} \text{ provided } |r| < 1$$

Evaluate $\sum_{n=0}^{\infty} \left(\frac{2^{2n}}{3^n}\right)$



TELESCOPING SUMS

Evaluate $\sum_{n=1}^{800} \left(\frac{1}{n} - \frac{1}{n+1}\right)$.

Evaluate $\sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n+1}\right)$.




Evaluate $\sum_{n=1}^{1000} \log\left(\frac{n+1}{n}\right)$.

Evaluate $\sum_{n=1}^{\infty} \log\left(\frac{n+1}{n}\right)$.



TABLE OF CONTENTS




For a convergent geometric or telescoping series, we can easily determine what the series converges to.

For other types of series, finding out what the series converges to can be very difficult. It is often necessary to resort to approximating the full sum by, for example, using a computer to find the sum of the first N terms, for some large N . But before we even try to do that, we should at least know *whether or not the series converges*.


Suppose a series $\sum_{n=1}^{\infty} a_n$ converges to a limit L . Let $S_N = \sum_{n=1}^N a_n$.

$$\begin{aligned} \lim_{N \rightarrow \infty} S_N &= \\ \lim_{N \rightarrow \infty} S_{N-1} &= \\ \lim_{N \rightarrow \infty} [S_N - S_{N-1}] &= \\ \lim_{N \rightarrow \infty} a_N &= \end{aligned}$$



Suppose a series $\sum_{n=1}^{\infty} a_n$ converges to a limit L . Let $S_N = \sum_{n=1}^N a_n$.

$$\begin{aligned} \lim_{N \rightarrow \infty} S_N &= \\ \lim_{N \rightarrow \infty} S_{N-1} &= \\ \lim_{N \rightarrow \infty} [S_N - S_{N-1}] &= \\ \lim_{N \rightarrow \infty} a_N &= \end{aligned}$$



Every convergent series has its N^{th} term, a_N , tending to 0 as $N \rightarrow \infty$.


Divergence Test

If the sequence $\{a_n\}_{n=c}^{\infty}$ fails to converge to zero as $n \rightarrow \infty$, then the series $\sum_{n=c}^{\infty} a_n$ diverges.

Do the following series diverge?


- ▶ $\sum_{n=0}^{\infty} (-1)^n$
- ▶ $\sum_{n=10}^{\infty} \left(\frac{1}{10} + \frac{1}{2^n} \right)$
- ▶ $\sum_{n=15}^{\infty} \frac{e^n}{2e^n - 1}$
- ▶ $\sum_{n=15}^{\infty} \frac{1}{n}$

USING THE DIVERGENCE TEST FOR $\sum a_n$




509/643 Warning 3.3.3

HARMONIC SERIES: $\sum_{n=1}^{\infty} \frac{1}{n}$



510/643 Example 3.3.4

$\sum_{n=1}^{\infty} \frac{1}{n}$ DIVERGES




$$S_1 = 1.0000$$

$$S_2 = 1.5000$$

$$S_3 = 1.8333$$


$$S_4 = 2.0833$$

$$S_5 = 2.2833$$



511/643


$\sum_{n=1}^{\infty} \frac{1}{n^2+1}$



512/643

$$\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$$

CONVERGES




$$S_1 = 0.5000$$

$$S_2 = 0.7000$$

$$S_3 = 0.8000$$

$$S_4 = 0.8588$$

$$S_5 = 0.8973$$




513/643

Integral Test

Let N_0 be any natural number. If $f(x)$ is a function which is defined and continuous for all $x \geq N_0$ and which obeys

- (i) $f(x) \geq 0$ for all $x \geq N_0$ and
- (ii) $f(x)$ decreases as x increases and
- (iii) $f(n) = a_n$ for all $n \geq N_0$.



Then

$$\sum_{n=1}^{\infty} a_n \text{ converges} \iff \int_{N_0}^{\infty} f(x) \, dx \text{ converges}$$


Furthermore, when the series converges, the truncation error satisfies

$$0 \leq \sum_{n=1}^{\infty} a_n - \sum_{n=1}^N a_n \leq \int_N^{\infty} f(x) \, dx \quad \text{for all } N \geq N_0$$

514/643 Theorem 3.3.5

Does the series $\sum_{n=10}^{\infty} \frac{1}{n \log n}$ converge or diverge?

Does the series $\sum_{n=10}^{\infty} \frac{1}{n \log n}$ converge or diverge?



$$\int_{10}^{\infty} \frac{1}{x \log x} \, dx = \infty$$


515/643

516/643

Integral Test, abridged

... When the series converges, the truncation error satisfies

$$0 \leq \sum_{n=1}^{\infty} a_n - \sum_{n=1}^N a_n \leq \int_N^{\infty} f(x) \, dx$$



517/643

Integral Test, abridged

When the series converges, the truncation error satisfies

$$0 \leq \sum_{n=1}^{\infty} a_n - \sum_{n=1}^N a_n \leq \int_N^{\infty} f(x) \, dx$$

We already decided that the series $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$ converges.

Suppose we had a computer add up the terms $n = 1$ through $n = 100$.

Use the integral test to bound the error, $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1} - \sum_{n=1}^{100} \frac{1}{n^2 + 1}$.

518/643

By computer, $\sum_{n=1}^{100} \frac{1}{n^2 + 1} \approx 1.0667$. Using the truncation error of about

0.01, give a (small) range of possible values for $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$.

$$0 \leq \sum_{n=1}^{\infty} \frac{1}{n^2 + 1} - \sum_{n=1}^{100} \frac{1}{n^2 + 1} \leq \int_{100}^{\infty} \frac{1}{x^2 + 1} \, dx$$

519/643

p -TEST

Let p be a positive constant. When we talked about improper integrals, we showed:

$$\int_1^{\infty} \frac{1}{x^p} \, dx \quad \left\{ \begin{array}{l} \text{converges if } p > 1 \\ \text{diverges if } p \leq 1 \end{array} \right.$$

Set $f(x) = \frac{1}{x^p}$.

- (i) $f(x) \geq 0$ for all $x \geq 1$, and
- (ii) $f(x)$ decreases as x increases

$$\sum_{n=1}^{\infty} \frac{1}{n^p} \, dx \quad \left\{ \right.$$

520/643 Example 3.3.6

Consider the series

$$\sum_{n=1}^{\infty} \frac{1}{n^3}$$

By the p -test, we know this series

How many terms should we add up to approximate the series to within an error of no more than 0.02?

$$\sum_{n=1}^{\infty} \frac{1}{n^3} \text{ converges to within } 0.02 \text{ of } \sum_{n=1}^5 \frac{1}{n^3}$$



$$S_1 = 1.0000$$

$$S_2 = 1.1250$$

$$S_3 = 1.1620$$

$$S_4 = 1.1776$$


$$S_5 = 1.1856$$



Observation


In a series with **positive** terms, the series either **converges**, or **diverges to infinity**.

If terms are "too big," series will diverge.




$$\sum \frac{1}{n^2} \text{ converges}$$

$$\sum \frac{1}{n^2 + n} \text{ converges, too}$$



Terms are "small enough" for sum to converge



Terms are also "small enough" for sum to converge

The Comparison Test

Let N_0 be a natural number and let $K > 0$.

- (a) If $|a_n| \leq Kc_n$ for all $n \geq N_0$ and $\sum_{n=0}^{\infty} c_n$ converges, then $\sum_{n=0}^{\infty} a_n$ converges.
- (b) If $a_n \geq Kd_n \geq 0$ for all $n \geq N_0$ and $\sum_{n=0}^{\infty} d_n$ diverges, then $\sum_{n=0}^{\infty} a_n$ diverges.

Consider $\sum_{n=1}^{\infty} \frac{1}{n-0.1}$.

- ▶ We know $0 < \frac{1}{n} < \frac{1}{n-0.1}$
- ▶ We know $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges (harmonic series)
- ▶ So, by the comparison test, $\sum_{n=1}^{\infty} \frac{1}{n-0.1}$ diverges as well.

Does the series $\sum_{n=1}^{\infty} \frac{n + \cos n}{n^3 - 1/3}$ converge or diverge?

Step 1: Intuition.

When n is very large, we expect:

- ▶ $n + \cos n \approx$
- ▶ $n^3 + \frac{1}{3} \approx$
- ▶ So, we expect $\frac{n + \cos n}{n^3 - 1/3} \approx$

Since $\sum_{n=1}^{\infty} \frac{1}{n^2} \dots$

we expect $\sum_{n=1}^{\infty} \frac{n + \cos n}{n^3 - 1/3}$ to also ...

Does the series $\sum_{n=1}^{\infty} \frac{n + \cos n}{n^3 - 1/3}$ converge or diverge?

Step 2: Choose comparison series.

The Comparison Test, abridged

Let N_0 be a natural number and let $K > 0$.

If $|a_n| \leq Kc_n$ for all $n \geq N_0$ and $\sum_{n=0}^{\infty} c_n$ converges, then $\sum_{n=0}^{\infty} a_n$ converges.

To show that original series **converges**, we should find a comparison series that also **converges** and whose terms (times some positive constant) are **larger** than the original terms. *There are many possibilities.* For $n \geq 1$,

- ▶ $n + \cos n <$
- ▶ $n^3 - \frac{1}{3} >$
- ▶ So $\frac{n + \cos n}{n^3 - 1/3} <$

Does the series $\sum_{n=1}^{\infty} \frac{n + \cos n}{n^3 - 1/3}$ converge or diverge?


Step 3: Verify.

The Comparison Test, abridged

Let N_0 be a natural number and let $K > 0$.


If $|a_n| \leq Kc_n$ for all $n \geq N_0$ and $\sum_{n=0}^{\infty} c_n$ converges, then $\sum_{n=0}^{\infty} a_n$ converges.

For the comparison test as we have seen it so far, to conclude that a given series **diverges**, we have to find a divergent comparison series whose terms are **smaller** than (a positive multiple of) those of our original series .



529/643

For the comparison test as we've seen it so far, to conclude that a given series **converges**, we have to find a convergent comparison series whose terms are **larger** than (a positive multiple of) those of our original series .



530/643

Limit Comparison Theorem

Let $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ be two series with $b_n > 0$ for all n . Assume that

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = L$$

exists.

- (a) If $\sum_{n=1}^{\infty} b_n$ converges, then $\sum_{n=1}^{\infty} a_n$ converges too.
- (b) If $L \neq 0$ and $\sum_{n=1}^{\infty} b_n$ diverges, then $\sum_{n=1}^{\infty} a_n$ diverges too.

In particular, if $L \neq 0$, then $\sum_{n=1}^{\infty} a_n$ converges if and only if $\sum_{n=1}^{\infty} b_n$ converges.

- ▶ For large n , $a_n \approx L \cdot b_n$;
- ▶ so we expect $\sum a_n$ to behave roughly like $\sum (L \cdot b_n)$;
- ▶ and since $L \neq 0$, we expect $\sum (L \cdot b_n)$ to converge if and only if $\sum b_n$ converges.

531/643 Theorem 3.3.11, with a very rough justification

By the p -test, $\sum_{n=1}^{\infty} \frac{1}{n^{3/2}}$ converges.

Can we conclude that $\sum_{n=1}^{\infty} \frac{1}{n^{3/2} - n + 1}$ also converges?

532/643

Does the series $\sum_{n=1}^{\infty} \frac{\sqrt{n+1}}{n^2-2n+3}$ converge or diverge?

Step 1: Intuition

For large n ,

533/643

Does the series $\sum_{n=1}^{\infty} \frac{\sqrt{n+1}}{n^2-2n+3}$ converge or diverge?

Step 2: Verify Intuition

Let $a_n = \frac{\sqrt{n+1}}{n^2-2n+3}$ and $b_n = \frac{1}{n^{3/2}}$.

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} =$$

534/643

COMPARISON STRATEGIES

- ▶ Before you can use either comparison test, you need to guess a series to compare.
- ▶ The series you guess should be easy to deal with.
 - ▶ p -series
 - ▶ geometric series
- ▶ Common guess (especially if monotone): consider “largest” piece of numerator and denominator
(constant) < (logarithm) < (polynomial) < (exponential)
- ▶ After you guess a comparison series, **show it works** by finding the correct inequality (comparison test), or computing the limit of the ratio (limit comparison test).

535/643

CHOOSE A SERIES TO COMPARE

$$\sum_{n=1}^{\infty} \frac{3n}{n^2+1}$$


$$\sum_{n=1}^{\infty} \frac{n^2+n+1}{n^5-n}$$

$$\sum_{k=1}^{\infty} \frac{k(2+\sin k)}{k\sqrt{2}}$$

$$\sum_{m=1}^{\infty} \frac{3m+\sin\sqrt{m}}{m^2}$$

536/643

TABLE OF CONTENTS



537/643

REVIEW

$$\text{Let } S_N = \sum_{n=1}^N a_n.$$

Simplify: $S_N - S_{N-1}$.

(This will come in handy soon.)

538/643

ALTERNATING SERIES

Alternating Series

The series

$$A_1 - A_2 + A_3 - A_4 + \dots = \sum_{n=1}^{\infty} (-1)^{n-1} A_n$$

is alternating if every $A_n \geq 0$.

Alternating series:

▶ $1 - 2 + 3 - 4 + 5 - 6 + 7 - 8 + \dots$


▶ $1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \dots$

Not alternating:

▶ $\cos(1) + \cos(2) + \cos(3) + \dots$

▶ $1 - \left(-\frac{1}{2}\right) + \frac{1}{3} - \left(-\frac{1}{4}\right) + \dots$

539/643



$$S_1 = 6.0000$$


$$S_2 = 1.0000$$

$$S_3 = 5.0000$$


$$S_4 = 2.0000$$

$$S_5 = 4.0000$$

$$S_6 = 3.0000$$




540/643




541/643

Consider an alternating series $a_1 - a_2 + a_3 - a_4 + \dots$, where $\{a_n\}$ is a sequence with positive, **decreasing** terms and with $\lim_{n \rightarrow \infty} a_n = 0$.



Since $a_2 > a_3$, we have $a_1 - (a_2 - a_3) < a_1$, so $S_3 < S_1$.

542/643



- ▶ For all $n \geq 2$, S_n lies between S_1 and S_2 .
- ▶ For all $n \geq 3$, S_n lies between S_2 and S_3 .
- ▶ For all $n \geq 4$, S_n lies between S_3 and S_4 .
- ▶ For all $n \geq 5$, S_n lies between S_4 and S_5 .

The difference between consecutive sums S_n and S_{n-1} is:

543/643

Alternating Series Test

Let $\{a_n\}_{n=1}^{\infty}$ be a sequence of real numbers that obeys

- (i) $a_n \geq 0$ for all $n \geq 1$;
- (ii) $a_{n+1} \leq a_n$ for all $n \geq 1$ (i.e. the sequence is monotone decreasing);
- (iii) and $\lim_{n \rightarrow \infty} a_n = 0$.

Then

$$a_1 - a_2 + a_3 - a_4 + \dots = \sum_{n=1}^{\infty} (-1)^{n-1} a_n = S$$

converges and, for each natural number N , $S - S_N$ is between 0 and (the first dropped term) $(-1)^N a_{N+1}$. Here S_N is, as previously, the N^{th} partial sum $\sum_{n=1}^N (-1)^{n-1} a_n$.

544/643 Theorem 3.3.14

Alternating Series Test (abridged)

Let $\{a_n\}_{n=1}^{\infty}$ be a sequence of real numbers that obeys

- (i) $a_n \geq 0$ for all $n \geq 1$;
- (ii) $a_{n+1} \leq a_n$ for all $n \geq 1$ (i.e. the sequence is monotone decreasing);
- (iii) and $\lim_{n \rightarrow \infty} a_n = 0$.

Then


$$a_1 - a_2 + a_3 - a_4 + \cdots = \sum_{n=1}^{\infty} (-1)^{n-1} a_n$$

converges.

- ▶ True or false: the harmonic series $\sum_{n=1}^{\infty} \frac{1}{n}$ converges.
- ▶ True or false: the alternating harmonic series $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ converges.

545/643

DIVERGENCE TEST + ALTERNATING SERIES TEST



546/643

Warning 3.3.3

Alternating Series Test

Let $\{a_n\}_{n=1}^{\infty}$ be a sequence of real numbers that obeys $a_n \geq 0$ for all $n \geq 1$; $a_{n+1} \leq a_n$ for all $n \geq 1$; and $\lim_{n \rightarrow \infty} a_n = 0$. Then $\sum_{n=1}^{\infty} (-1)^{n-1} a_n = S$ converges and $S - S_N$ is between 0 and $(-1)^N a_{N+1}$.

Using a computer, you find $\sum_{n=1}^{99} \frac{(-1)^{n-1}}{n} \approx 0.698$.

How close is that to the value $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}$?

547/643

Alternating Series Test

Let $\{a_n\}_{n=1}^{\infty}$ be a sequence of real numbers that obeys $a_n \geq 0$ for all $n \geq 1$; $a_{n+1} \leq a_n$ for all $n \geq 1$; and $\lim_{n \rightarrow \infty} a_n = 0$. Then $\sum_{n=1}^{\infty} (-1)^{n-1} a_n = S$ converges and $S - S_N$ is between 0 and $(-1)^N a_{N+1}$.

Using a computer, you find $\sum_{n=1}^{19} (-1)^{n-1} \frac{n^2}{n^2+1} \approx 0.6347$.

How close is that to the value $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{n^2}{n^2+1}$?

548/643

Recall for a geometric series, the **ratios of consecutive terms** is constant.

$$\begin{array}{ccccccc} & \times \frac{1}{2} & & \times \frac{1}{2} & & \times \frac{1}{2} & & \times \frac{1}{2} \\ & \swarrow & \searrow & \swarrow & \searrow & \swarrow & \searrow & \swarrow \\ \frac{1}{2} & + & \frac{1}{4} & + & \frac{1}{8} & + & \frac{1}{16} & + & \frac{1}{32} & + \dots \end{array}$$

$$\underbrace{\frac{1/4}{1/2}} = \underbrace{\frac{1/8}{1/4}} = \underbrace{\frac{1/16}{1/8}} = \underbrace{\frac{1/32}{1/16}} = \frac{1}{2}$$

If that ratio has magnitude **less than one**, then the series converges.
If the ratio has magnitude **greater than one**, the series diverges.

549/643

For series convergence, we are concerned with what happens to terms a_n when n is sufficiently large.
Suppose for a sequence a_n , $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = L$ for some constant L .

$$\begin{array}{cccccccc} a_n & + & a_{n+1} & + & a_{n+2} & + & a_{n+3} & + & a_{n+4} & + \dots \\ \underbrace{} & \underbrace{\phantom{a_{n+1}}} & \underbrace{\phantom{a_{n+2}}} & \underbrace{\phantom{a_{n+3}}} & \underbrace{\phantom{a_{n+4}}} & \underbrace{\phantom{a_{n+5}}} & \underbrace{\phantom{a_{n+6}}} & \underbrace{\phantom{a_{n+7}}} & \underbrace{\phantom{a_{n+8}}} & \underbrace{\phantom{a_{n+9}}} \\ \frac{a_{n+1}}{a_n} & \approx & \frac{a_{n+2}}{a_{n+1}} & \approx & \frac{a_{n+3}}{a_{n+2}} & \approx & \frac{a_{n+4}}{a_{n+3}} & \approx & \frac{a_{n+5}}{a_{n+4}} & \approx & L \end{array}$$

Like in a geometric series:

If L has magnitude **less than one**, then the series converges.
If L has magnitude **greater than one**, the series diverges.

550/643

Ratio Test

Let N be any positive integer and assume that $a_n \neq 0$ for all $n \geq N$.

- (a) If $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L < 1$, then $\sum_{n=1}^{\infty} a_n$ converges.
(b) If $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L > 1$, or $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \infty$, then $\sum_{n=1}^{\infty} a_n$ diverges.

551/643 Theorem 3.3.18

Ratio Test

Let N be any positive integer and assume that $a_n \neq 0$ for all $n \geq N$.

- (a) If $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L < 1$, then $\sum_{n=1}^{\infty} a_n$ converges.
(b) If $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L > 1$, or $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \infty$, then $\sum_{n=1}^{\infty} a_n$ diverges.

Use the ratio test to determine whether the series

$$\sum_{n=1}^{\infty} \frac{n}{3^n}$$

converges or diverges.

552/643



REMARK

The series we just considered, $\sum_{n=1}^{\infty} \frac{n}{3^n}$, looks similar to a geometric series, but it is not exactly a geometric series. That's a good indicator that the ratio test will be helpful!

We could have used other tests, but ratio was probably the easiest.

- ▶ Integral test: $\int \frac{x}{3^x} dx$ can be evaluated using integration by parts.
- ▶ Comparison test:
 - ▶ $\sum \frac{1}{3^n}$ is not a valid comparison series, nor is $\sum n$.
 - ▶ Because $n < 2^n$ for all $n \geq 1$, the series $\sum (\frac{2}{3})^n$ will work.
- ▶ The divergence test is inconclusive, and the alternating series test does not apply. Our series is not geometric, and not obviously telescoping.

553/643

Ratio Test

Let N be any positive integer and assume that $a_n \neq 0$ for all $n \geq N$.

- (a) If $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L < 1$, then $\sum_{n=1}^{\infty} a_n$ converges.
- (b) If $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L > 1$, or $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \infty$, then $\sum_{n=1}^{\infty} a_n$ diverges.

Let a and x be nonzero constants. Use the ratio test to determine whether

$$\sum_{n=1}^{\infty} anx^{n-1}$$

converges or diverges. (This may depend on the values of a and x .)



554/643

Let x be a constant. Use the ratio test to determine whether

$$\sum_{n=1}^{\infty} \frac{(-3)^n \sqrt{n+1}}{2n+3} x^n$$

converges or diverges. (This may depend on the value of x .)



555/643 Example 3.3.23

FILL IN THE BLANKS

Divergence Test

If the sequence $\{a_n\}_{n=c}^{\infty}$ then the series $\sum_{n=c}^{\infty} a_n$ diverges.

Ratio Test

Let N be any positive integer and assume that $a_n \neq 0$ for all $n \geq N$.


- (a) If $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|$, then $\sum_{n=1}^{\infty} a_n$ converges.
- (b) If $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|$, or $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \infty$, then $\sum_{n=1}^{\infty} a_n$ diverges.

556/643

Integral Test

Let N_0 be any natural number. If $f(x)$ is a function which is defined and continuous for all $x \geq N_0$ and which obeys

- (i) and
 (ii) and
 (iii) $f(n) = a_n$ for all $n \geq N_0$.



Then

$$\sum_{n=1}^{\infty} a_n \text{ converges} \iff \int_{N_0}^{\infty} f(x) \, dx \text{ converges}$$

Furthermore, when the series converges, the truncation error satisfies

$$0 \leq \sum_{n=1}^{\infty} a_n - \sum_{n=1}^N a_n \leq \int_N^{\infty} f(x) \, dx \quad \text{for all } N \geq N_0$$

557/643

FILL IN IN THE BLANKS

The Comparison Test

Let N_0 be a natural number and let $K > 0$.

- (a) If $|a_n| \leq Kc_n$ for all $n \geq N_0$ and $\sum_{n=0}^{\infty} c_n$ converges, then $\sum_{n=0}^{\infty} a_n$ converges.
 (b) If $a_n \geq Kd_n \geq 0$ for all $n \geq N_0$ and $\sum_{n=0}^{\infty} d_n$ diverges, then $\sum_{n=0}^{\infty} a_n$ diverges.

558/643

FILL IN IN THE BLANKS

Limit Comparison Theorem

Let $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ be two series with $b_n > 0$ for all n . Assume that

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = L$$

exists.

- (a) If $\sum_{n=1}^{\infty} b_n$ converges, then $\sum_{n=1}^{\infty} a_n$ converges too.
 (b) If $L \neq 0$ and $\sum_{n=1}^{\infty} b_n$ diverges, then $\sum_{n=1}^{\infty} a_n$ diverges too.

In particular, if , then $\sum_{n=1}^{\infty} a_n$ converges if and only if $\sum_{n=1}^{\infty} b_n$ converges.

559/643

Alternating Series Test

Let $\{a_n\}_{n=1}^{\infty}$ be a sequence of real numbers that obeys

- (i)
 (ii) $a_{n+1} \leq a_n$ for all $n \geq 1$ (i.e. the sequence is monotone decreasing);
 (iii) and

Then

$$a_1 - a_2 + a_3 - a_4 + \dots = \sum_{n=1}^{\infty} (-1)^{n-1} a_n = S$$

converges and, for each natural number N , $S - S_N$ is between 0 and (the first dropped term) $(-1)^N a_{N+1}$. Here S_N is, as previously, the N^{th} partial sum $\sum_{n=1}^N (-1)^{n-1} a_n$.

560/643

LIST OF CONVERGENCE TESTS

Divergence Test

When the n^{th} term in the series *fails* to converge to zero as n tends to infinity.

This is a good first thing to check: if it works, it's quick, but it doesn't always work.

Alternating Series Test

- ▶ successive terms in the series alternate in sign
- ▶ don't forget to check that successive terms decrease in magnitude and tend to zero as n tends to infinity

Integral Test

- ▶ works well when, if you substitute x for n in the n^{th} term you get a function, $f(x)$, that you can easily integrate
- ▶ don't forget to check that $f(x) \geq 0$ and that $f(x)$ decreases as x increases

561/643

LIST OF CONVERGENCE TESTS

Ratio Test

- ▶ works well when $\frac{a_{n+1}}{a_n}$ simplifies enough that you can easily compute $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L$
- ▶ this often happens when a_n contains powers, like 7^n , or factorials, like $n!$
- ▶ don't forget that $L = 1$ tells you nothing about the convergence/divergence of the series

Comparison Test and Limit Comparison Test

- ▶ Comparison test lets you ignore pieces of a function that feel extraneous (like replacing $n^2 + 1$ with n^2) *but* there is a test to make sure the comparison is still valid. Either the limit of a ratio is the right thing, or an inequality goes the right way.
- ▶ Limit comparison works well when, for very large n , the n^{th} term a_n is approximately the same as a simpler, nonnegative term b_n

562/643

- ▶ The integral test gave us the p -test. When you're looking for comparison series, p -series $\sum \frac{1}{n^p}$ are often good choices, because their convergence or divergence is so easy to ascertain.

- ▶ Geometric series have the form $\sum a \cdot r^n$ for some nonzero constants a and r . The magnitude of r is all you need to know to decide whether they converge or diverge, so these are also common comparison series.

- ▶ Telescoping series have partial sums that are easy to find because successive terms cancel out. These are less obvious, and are less common choices for comparison series.

563/643

Test List

- ▶ divergence
- ▶ integral
- ▶ alternating series
- ▶ ratio
- ▶ comparison
- ▶ limit comparison

Determine whether the series $\sum_{n=1}^{\infty} \frac{\cos n}{2^n}$ converges or diverges.

564/643



Test List

- ▶ divergence
- ▶ integral
- ▶ alternating series
- ▶ ratio
- ▶ comparison
- ▶ limit comparison

Determine whether the series $\sum_{n=1}^{\infty} \frac{2^n \cdot n^2}{(n+5)^5}$ converges or diverges.



Test List


- ▶ divergence
- ▶ integral
- ▶ alternating series
- ▶ ratio
- ▶ comparison
- ▶ limit comparison

Determine whether the series $\sum_{n=1}^{\infty} \frac{1}{n} \sin\left(\frac{1}{n}\right)$ converges or diverges.

Hint: If $\theta \geq 0$ then $\sin \theta \leq \theta$.



TABLE OF CONTENTS



FOUR SERIES

Let $a_n = \left(-\frac{2}{3}\right)^n$. Do the following series converge or diverge?

$$\sum_{n=0}^{\infty} a_n \qquad \sum_{n=0}^{\infty} |a_n|$$

Let $b_n = \frac{(-1)^n}{n}$. Do the following series converge or diverge?

$$\sum_{n=1}^{\infty} b_n \qquad \sum_{n=1}^{\infty} |b_n|$$

The series

$$\sum_{n=0}^{\infty} \left(-\frac{2}{3}\right)^n$$

is called **absolutely convergent**, because the series converges and if we replace the terms being added by their absolute values, that series *still* converges.

The series

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{n}$$

is called **conditionally convergent**, because the series converges, but if we replace the terms being added by their absolute values, that series *diverges*.

Absolute and conditional convergence

(a) A series $\sum_{n=1}^{\infty} a_n$ is said to **converge absolutely** if the series

$$\sum_{n=1}^{\infty} |a_n| \text{ converges.}$$

(b) If $\sum_{n=1}^{\infty} a_n$ converges but $\sum_{n=1}^{\infty} |a_n|$ diverges we say that

$$\sum_{n=1}^{\infty} a_n \text{ is **conditionally convergent**.}$$

Theorem

If the series $\sum_{n=1}^{\infty} |a_n|$ converges then the series $\sum_{n=1}^{\infty} a_n$ also converges.

That is, absolute convergence implies convergence.

If $\sum a_n \dots$	and $\sum a_n \dots$	then we say $\sum a_n$ is ...
converges	converges	
converges	diverges	
diverges	diverges	
diverges	converges	

Does the series

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n^2}$$

converge or diverge?




Does the series

$$\sum_{n=1}^{\infty} \frac{\sin(n)}{n^2}$$

converge or diverge?



TABLE OF CONTENTS



Finite addition is commutative

$$1 + 2 + 3 + 4 = 4 + 1 + 3 + 2$$

What happens if we re-arrange the terms in a series?

We'll illustrate some possibilities, but first we need to establish some preliminary results.

PRELIMINARY RESULTS

Split up the alternating harmonic series into two series: one with the positive terms, and one with the negative terms.

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \frac{1}{7} - \frac{1}{8} \dots$$

$$\begin{aligned} & -\frac{1}{2} - \frac{1}{4} - \frac{1}{6} - \frac{1}{8} - \dots \\ &= -\frac{1}{2} \left(1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots \right) \\ &= -\frac{1}{2} \sum_{n=1}^{\infty} \frac{1}{n} \end{aligned}$$

So, we can make an arbitrarily large negative number by adding up these terms.

$$\begin{aligned} & 1 + \frac{1}{3} + \frac{1}{5} + \frac{1}{7} + \dots \\ & \geq \frac{1}{2} + \frac{1}{4} + \frac{1}{6} + \frac{1}{8} + \dots \end{aligned}$$

So, we can make an arbitrarily large positive number by adding up these terms.

PRELIMINARY RESULTS



We've shown that the alternating harmonic series converges. We don't have the tools to do it just yet, but later we'll be able to compute what it converges to:

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} = \log 2$$

Surprising fact: if we reorder the terms of the series carefully, we can make a new series adding up to any number we want.

577/643



Rearrange the alternating harmonic series to sum to 0.



- ▶ Add positive terms until the partial sum is greater than 0.
- ▶ Add negative terms until the partial sum is less than 0.
- ▶ Repeat.

578/643


Rearrange the alternating harmonic series to sum to 0.



- ▶ Add positive terms until the partial sum is greater than 0.
- ▶ Add negative terms until the partial sum is less than 0.
- ▶ Repeat.

579/643


Rearrange the alternating harmonic series to sum to 2.



- ▶ Add positive terms until the partial sum is greater than 2.
- ▶ Add negative terms until the partial sum is less than 2.
- ▶ Repeat.

580/643

Rearrange the alternating harmonic series to sum to 2.




- ▶ Add positive terms until the partial sum is greater than 2.
- ▶ Add negative terms until the partial sum is less than 2.
- ▶ Repeat.

In fact: you can reorder *any* conditionally convergent series to

- ▶ add up to *any* number, or
- ▶ diverge to infinity, or
- ▶ diverge to negative infinity.

This doesn't work with absolutely convergent series.


Let's try to rearrange the terms of $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^2}$ to add up to 0:



- ▶ Add positive terms until the partial sum is greater than 0.
- ▶ Add negative terms (those with $n = 2m, m = 1, 2, 3, \dots$) until the partial sum is less than 0.
- ▶ Repeat.

This doesn't work with absolutely convergent series.

Let's try to rearrange the terms of $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^2}$ to add up to 0:




- ▶ Add positive terms until the partial sum is greater than 0.
- ▶ Add negative terms (those with $n = 2m, m = 1, 2, 3, \dots$) until the partial sum is less than 0.
- ▶ Repeat.

In fact: you can reorder *any* conditionally convergent series to

- ▶ add up to *any* number, or
- ▶ diverge to infinity, or
- ▶ diverge to negative infinity.

Changing the order of terms in an absolutely convergent series does not change its value.

TABLE OF CONTENTS



Recall the geometric series: for a constant r , with $|r| < 1$:


$$\sum_{n=0}^{\infty} r^n = \frac{1}{1-r}$$

We can think of this as a function. If we set

$$f(x) = \sum_{n=0}^{\infty} x^n$$

and restrict our domain to $-1 < x < 1$, then

$$f(x) = \sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$$



Why would we ever prefer to write $\sum_{n=0}^{\infty} x^n$ instead of $\frac{1}{1-x}$?

The function

$$\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + x^4 + \dots$$

isn't a polynomial, but in certain ways it behaves like one. For $|x| < 1$:

$$\frac{d}{dx} \left\{ \frac{1}{1-x} \right\} = \frac{d}{dx} \sum_{n=0}^{\infty} x^n = \sum_{n=0}^{\infty} \left(\frac{d}{dx} \{x^n\} \right) = \sum_{n=0}^{\infty} nx^{n-1}$$

$$\int \frac{1}{1-x} dx = \int \left(\sum_{n=0}^{\infty} x^n \right) dx = \sum_{n=0}^{\infty} \left(\int x^n dx \right) = \sum_{n=0}^{\infty} \frac{x^{n+1}}{n+1}$$

Definition

A series of the form

$$\sum_{n=0}^{\infty} A_n(x-c)^n = A_0 + A_1(x-c) + A_2(x-c)^2 + A_3(x-c)^3 + \dots$$

is called a *power series in $(x-c)$* or a *power series centered on c* . The numbers A_n are called the coefficients of the power series.

One often considers power series centered on $c = 0$ and then the series reduces to

$$A_0 + A_1x + A_2x^2 + A_3x^3 + \dots = \sum_{n=0}^{\infty} A_nx^n$$

$$\sum_{n=0}^{\infty} A_n(x-c)^n = A_0 + A_1(x-c) + A_2(x-c)^2 + A_3(x-c)^3 + \dots$$

In a power series, we think of the coefficients A_n as fixed constants, and we think of x as the variable of a function.

Evaluate the power series $\sum_{n=0}^{\infty} A_n(x-c)^n$ when $x = c$:

A fundamental question we want to ask when we see a series is whether it converges or diverges. So, let's find all values of x for which the power series

$$\sum_{n=1}^{\infty} \frac{x^n}{n} = x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots$$

converges.



A fundamental question we want to ask when we see a series is whether it converges or diverges. So, let's find all values of x for which the power series

$$\sum_{n=1}^{\infty} \frac{x^n}{n} = x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots$$

converges.

Definition

Consider the power series

$$\sum_{n=0}^{\infty} A_n(x - c)^n.$$

The set of real x -values for which it converges is called the interval of convergence of the series.



Find the interval of convergence of the power series

$$\sum_{n=0}^{\infty} 2^n(x - 1)^n = 1 + 2(x - 1) + 2^2(x - 1)^2 + 2^3(x - 1)^3 + \dots$$



What happens if we apply the ratio test to a generic power series,


$$\sum_{n=0}^{\infty} A_n(x - c)^n?$$

$$\lim_{n \rightarrow \infty} \left| \frac{A_{n+1}(x - c)^{n+1}}{A_n(x - c)^n} \right| = \lim_{n \rightarrow \infty} \left| \frac{A_{n+1}}{A_n}(x - c) \right| = |x - c| \lim_{n \rightarrow \infty} \left| \frac{A_{n+1}}{A_n} \right|$$


- ▶ If $\left| \frac{A_{n+1}}{A_n} \right|$ does not approach a limit as $n \rightarrow \infty$, the ratio test tells us nothing. (We should try other tests.)
- ▶ If $\lim_{n \rightarrow \infty} \left| \frac{A_{n+1}}{A_n} \right| = 0$, then
- ▶ If $\lim_{n \rightarrow \infty} \left| \frac{A_{n+1}}{A_n} \right| = \infty$, then
- ▶ If $\lim_{n \rightarrow \infty} \left| \frac{A_{n+1}}{A_n} \right| = A$ for some real number A , then

Definition: Radius of Convergence


- (a) Let $0 < R < \infty$. If $\sum_{n=0}^{\infty} A_n(x - c)^n$ converges for $|x - c| < R$, and diverges for $|x - c| > R$, then we say that the series has radius of convergence R .




- (b) If $\sum_{n=0}^{\infty} A_n(x - c)^n$ converges for every number x , we say that the series has an infinite radius of convergence.




- (c) If $\sum_{n=0}^{\infty} A_n(x - c)^n$ diverges for every $x \neq c$, we say that the series has radius of convergence zero.




- ▶ We saw that $\sum_{n=0}^{\infty} x^n$ converges when $|x| < 1$ and diverges when $|x| > 1$, so this series has radius of convergence $R =$



- ▶ We saw that $\sum_{n=1}^{\infty} \frac{x^n}{n}$ converges when $|x| < 1$ and diverges when $|x| > 1$, so this series also has radius of convergence $R =$



- ▶ We saw that $\sum_{n=1}^{\infty} 2^n(x-1)^n$ converges when $|x-1| < \frac{1}{2}$ and diverges when $|x-1| > \frac{1}{2}$, so this series has radius of convergence $R =$



What is the radius of convergence for the series $\sum_{n=0}^{\infty} \frac{x^n}{n!}$?

Recall: $n! = (n)(n-1)(n-2) \cdots (2)(1)$.




What is the radius of convergence for the series $\sum_{n=0}^{\infty} n! \cdot (x-3)^n$?


Theorem

Given a power series (say with centre c), one of the following holds.

- (a) The power series converges for every number x . In this case we say that the radius of convergence is ∞ .



- (b) There is a number $0 < R < \infty$ such that the series converges for $|x - c| < R$ and diverges for $|x - c| > R$. Then R is called the radius of convergence.



- (c) The series converges for $x = c$ and diverges for all $x \neq c$. In this case, we say that the radius of convergence is 0.



We are told that a certain power series with centre $c = 3$ converges at $x = 4$ and diverges at $x = 1$. What else can we say about the convergence or divergence of the series for other values of x ?

Operations on Power Series

Assume that the functions $f(x)$ and $g(x)$ are given by the power series

$$f(x) = \sum_{n=0}^{\infty} A_n(x-c)^n \quad g(x) = \sum_{n=0}^{\infty} B_n(x-c)^n$$

for all x obeying $|x-c| < R$. Let K be a constant. Then:

$$f(x) + g(x) = \sum_{n=0}^{\infty} [A_n + B_n] (x-c)^n$$

$$Kf(x) = \sum_{n=0}^{\infty} K A_n (x-c)^n$$

for all x obeying $|x-c| < R$.

Operations on Power Series

Assume that the functions $f(x)$ and $g(x)$ are given by the power series

$$f(x) = \sum_{n=0}^{\infty} A_n(x-c)^n \quad g(x) = \sum_{n=0}^{\infty} B_n(x-c)^n$$

for all x obeying $|x-c| < R$. Let K be a constant. Then:

$$\begin{aligned} (x-c)^N f(x) &= \sum_{n=0}^{\infty} A_n (x-c)^{n+N} \quad \text{for any integer } N \geq 1 \\ &= \sum_{k=N}^{\infty} A_{k-N} (x-c)^k \quad \text{where } k = n + N \end{aligned}$$

for all x obeying $|x-c| < R$.

Operations on Power Series

Assume that the functions $f(x)$ and $g(x)$ are given by the power series

$$f(x) = \sum_{n=0}^{\infty} A_n(x-c)^n \quad g(x) = \sum_{n=0}^{\infty} B_n(x-c)^n$$

for all x obeying $|x-c| < R$. Let K be a constant. Then:

$$f'(x) = \sum_{n=0}^{\infty} A_n n (x-c)^{n-1} = \sum_{n=1}^{\infty} A_n n (x-c)^{n-1}$$

$$\int_c^x f(t) dt = \sum_{n=0}^{\infty} A_n \frac{(x-c)^{n+1}}{n+1}$$

$$\int f(x) dx = \left[\sum_{n=0}^{\infty} A_n \frac{(x-c)^{n+1}}{n+1} \right] + C \quad \text{with } C \text{ an arbitrary constant}$$

for all x obeying $|x-c| < R$.

Operations on Power Series

Assume that the functions $f(x)$ and $g(x)$ are given by the power series

$$f(x) = \sum_{n=0}^{\infty} A_n(x-c)^n \quad g(x) = \sum_{n=0}^{\infty} B_n(x-c)^n$$

for all x obeying $|x-c| < R$. Let K be a constant. Then:

for all x obeying $|x-c| < R$.

Differentiating, antiderivating, multiplying by a nonzero constant, and multiplying by a positive power of $(x-c)$ do not change the radius of convergence of $f(x)$ (although they may change the interval of convergence).

Given that $\frac{d}{dx} \left\{ \frac{1}{1-x} \right\} = \frac{1}{(1-x)^2}$, find a power series representation for $\frac{1}{(1-x)^2}$ when $|x| < 1$.



Find a power series representation for $\log(1+x)$ when $|x| < 1$.



Find a power series representation for $\arctan(x)$ when $|x| < 1$.



Substituting in a Power Series

Assume that the function $f(x)$ is given by the power series

$$f(x) = \sum_{n=0}^{\infty} A_n x^n$$

for all x in the interval I . Also let K and k be real constants. Then


$$f(Kx^k) = \sum_{n=0}^{\infty} A_n K^n x^{kn}$$

whenever Kx^k is in I . In particular, if $\sum_{n=0}^{\infty} A_n x^n$ has radius of convergence R , K is nonzero and k is a natural number, then $\sum_{n=0}^{\infty} A_n K^n x^{kn}$ has radius of convergence $\sqrt[k]{R/|K|}$.

Find a power series representation for $\frac{1}{5-x}$ with centre 3.



TABLE OF CONTENTS



Taylor polynomial

Let a be a constant and let n be a non-negative integer. The n^{th} order Taylor polynomial for $f(x)$ about $x = a$ is

$$T_n(x) = \sum_{k=0}^n \frac{1}{k!} f^{(k)}(a) \cdot (x - a)^k.$$

Taylor series

The Taylor series for the function $f(x)$ expanded around a is the power series

$$\sum_{n=0}^{\infty} \frac{1}{n!} f^{(n)}(a) (x - a)^n.$$

When $a = 0$ it is also called the Maclaurin series of $f(x)$.

Let's compute some Taylor series, using the definition.

The method is nearly identical to finding Taylor *polynomials*, which is covered in CLP-1.



Find the Maclaurin series for $f(x) = \sin x$.



Find the Maclaurin series for $f(x) = \cos x$.

The Maclaurin series for $f(x) = e^x$ is:




Let $T_n(x)$ be the n -th order Taylor polynomial of the function $f(x)$, centred at a .

When we introduced Taylor polynomials in CLP-1, we framed $T_n(x)$ as an approximation of $f(x)$.

Let's see how those approximations look in two cases:

TAYLOR POLYNOMIALS FOR e^x



It seems like high-order Taylor polynomials do a pretty good job of approximating the function e^x , at least when x is near enough to 0.


TAYLOR POLYNOMIALS FOR A DIFFERENT FUNCTION

But that is not the case for all functions. Define

$$f(x) = \begin{cases} e^{-\frac{1}{x}} & x > 0 \\ 0 & x \leq 0 \end{cases}$$

Using the definition of the derivative and l'Hôpital's rule, one can show that $f^{(n)}(0) = 0$ for all natural numbers n .

TAYLOR POLYNOMIALS FOR A DIFFERENT FUNCTION



Taylor polynomial approximations don't **always** get better as their orders increase – it depends on the function being approximated.

INVESTIGATION

- ▶ We found the Maclaurin series for $f(x) = e^x$ is $\sum_{n=0}^{\infty} \frac{x^n}{n!}$.
- ▶ But, it's not immediately clear whether $e^x \stackrel{?}{=} \sum_{n=0}^{\infty} \frac{x^n}{n!}$.
- ▶ We're going to demonstrate that e^x is in fact equal to $\sum_{n=0}^{\infty} \frac{x^n}{n!}$. The proof involves a particular limit: $\lim_{n \rightarrow \infty} \frac{|x|^n}{n!}$. We'll talk about that limit first, so that it doesn't distract us later.

621/643

Intermediate result: $\lim_{n \rightarrow \infty} \frac{|x|^n}{n!}$, when x is some fixed number.

For large n , we can think of $\frac{|x|^n}{n!}$ as a long multiplication, with decreasing terms. At some point, those terms are all decreasing *and less than 1*.

$$\frac{|x|^n}{n!} = \frac{|x| \cdot |x| \cdot |x| \cdot |x| \cdot |x| \cdot |x| \cdot \dots \cdot |x|}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot \dots \cdot n}$$

$$=$$

622/643 Convenient notation: $\lceil x \rceil$ is the number you get when you round x up to the nearest whole number.

Intermediate result: $\lim_{n \rightarrow \infty} \frac{|x|^n}{n!}$, when x is some fixed number.

We're multiplying terms that are closer and closer to 0, so it seems quite reasonable that this sequence should converge to 0.

For a more formal proof, we can use the squeeze theorem to compare this sequence to a geometric sequence.

INVESTIGATION

- ▶ We found the Maclaurin series for $f(x) = e^x$ is $\sum_{n=0}^{\infty} \frac{x^n}{n!}$.

- ▶ But, it's not immediately clear whether $e^x \stackrel{?}{=} \sum_{n=0}^{\infty} \frac{x^n}{n!}$.
How could we determine this?

▶

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

$$\iff 0 = e^x - \sum_{n=0}^{\infty} \frac{x^n}{n} = e^x - \lim_{n \rightarrow \infty} \underbrace{\sum_{k=0}^n \frac{x^k}{k!}}_{T_n(x)} = \lim_{n \rightarrow \infty} \underbrace{[e^x - T_n(x)]}_{E_n(x)}$$

$$\iff 0 = \lim_{n \rightarrow \infty} E_n(x) \quad (\text{for all } x)$$


623/643 Convenient notation: $\lceil x \rceil$ is the number you get when you round x up to the nearest whole number.

624/643

TAYLOR POLYNOMIAL ERROR FOR $f(x) = e^x$

If $\lim_{n \rightarrow \infty} E_n(x) = 0$ for all x , then $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$ for all x .

It *looks* plausible, especially when x is close to 0. Let's try to prove it.



625/643

Equation 3.6.1-b

Let $T_n(x)$ be the n -th order Taylor approximation of a function $f(x)$, centred at a . Then $E_n(x) = f(x) - T_n(x)$ is the error in the n -th order Taylor approximation.

For some c strictly between x and a ,

$$E_n(x) = \frac{1}{(n+1)!} f^{(n+1)}(c) \cdot (x-a)^{n+1}$$

When $f(x) = e^x$,


$$E_n(x) = e^c \frac{x^{n+1}}{(n+1)!}$$

for some c between 0 and x .

626/643 CLP-1 Equation 3.4.33, CLP-2 Equation 3.6.1-b

$$\begin{aligned} E_n(x) &= e^x - T_n(x) \\ &= e^c \frac{x^{n+1}}{(n+1)!} \quad \text{for some } c \text{ between } 0 \text{ and } x \end{aligned}$$

We found $0 \leq |E_n(x)| < e^{|x|} \frac{|x|^{n+1}}{(n+1)!}$ for large n , hence $\lim_{n \rightarrow \infty} |E_n(x)| = 0$.



For a particular value of x :

$$\text{We saw } 0 = \lim_{n \rightarrow \infty} \frac{|x|^{n+1}}{(n+1)!}$$

$$\text{so } 0 = \lim_{n \rightarrow \infty} E_n(x)$$

$$\text{That is, } 0 = \lim_{n \rightarrow \infty} [e^x - T_n(x)]$$

$$\text{So, } e^x = \lim_{n \rightarrow \infty} T_n(x)$$

$$= \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

627/643

628/643

TAYLOR POLYNOMIAL ERROR FOR SINE AND COSINE

Equation 3.6.1-b

Let $T_n(x)$ be the n -th order Taylor approximation of a function $f(x)$, centred at a . Then $E_n(x) = f(x) - T_n(x)$ is the error in the n -th order Taylor approximation.


For some c strictly between x and a ,

$$E_n(x) = \frac{1}{(n+1)!} f^{(n+1)}(c) \cdot (x-a)^{n+1}$$


Suppose $f(x)$ is either $\sin x$ or $\cos x$. Is $f(x)$ equal to its Maclaurin series?

$$|E_n(x)| = \frac{1}{(n+1)!} |f^{(n+1)}(c)| |x|^{n+1}$$

TAYLOR POLYNOMIALS FOR $\sin(x)$



TAYLOR POLYNOMIALS FOR $\cos(x)$



Selected Taylor series that equal their functions

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} \quad \text{for all } -\infty < x < \infty$$

$$\sin(x) = \sum_{n=0}^{\infty} (-1)^n \frac{1}{(2n+1)!} x^{2n+1} \quad \text{for all } -\infty < x < \infty$$

$$\cos(x) = \sum_{n=0}^{\infty} (-1)^n \frac{1}{(2n)!} x^{2n} \quad \text{for all } -\infty < x < \infty$$

$$\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n \quad \text{for all } -1 < x < 1$$

$$\log(1+x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{n+1}}{n+1} \quad \text{for all } -1 < x \leq 1$$

$$\arctan x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1} \quad \text{for all } -1 \leq x \leq 1$$

COMPUTING π

Use the fact that $\arctan 1 = \frac{\pi}{4}$ to find a series converging to π whose terms are rational numbers.




ERROR FUNCTION

The *error function*

$$\operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt$$

is used in computing “bell curve” probabilities.



ERROR FUNCTION

The *error function*

$$\operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt$$

is used in computing “bell curve” probabilities.

The indefinite integral of the integrand e^{-t^2} cannot be expressed in terms of standard functions. But we can still evaluate the integral to within any desired degree of accuracy by using the Taylor expansion of the exponential.

For example, evaluate $\operatorname{erf}\left(\frac{1}{\sqrt{2}}\right)$.



EVALUATING A CONVERGENT SERIES

Evaluate $\sum_{n=1}^{\infty} \frac{1}{n \cdot 3^n}$



FINDING A HIGH-ORDER DERIVATIVE

Let $f(x) = \sin(2x^3)$. Find $f^{(15)}(0)$, the fifteenth derivative of f at $x = 0$.



Given that $\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \cdots$, we have a new way of evaluating the familiar limit












$$\lim_{x \rightarrow 0} \frac{\sin x}{x} :$$











Evaluate $\lim_{x \rightarrow 0} \frac{\arctan x - x}{\sin x - x}$.



Included Work

-  Vector illustration of role play game map icon for an arch is in the Public Domain (accessed Jan 8, 2021), 533
-  balancing by [Olena Panasovska](#) is licensed under [CC BY 3.0](#) (accessed 10 January 2023), 409
-  'Balloon' by [Simon Farkas](#) is licensed under [CC-BY](#) (accessed November 2022, edited), 497, 541
-  Elephant eating by [Paulami Roychoudhury](#) is licensed under [CC BY 3.0](#) (accessed 10 January 2023), 433
-  Firewood by [Aline Escobar](#) is licensed under [CC BY 3.0](#) (accessed 10 January 2023), 433
-  HAND GRAB by [Oleksandr Panasovskyi](#) is licensed under [CC BY 3.0](#) (accessed 10 January 2023), 389, 393
-  Hippopotamus vector image is in the Public Domain (accessed January 2021), 525, 533
-  kettle bell by [Made](#) is licensed under [CC BY 3.0](#) (accessed 10 January 2023), 401, 405
-  Roasting Marshmallows by [Caitlin George](#) is licensed under [CC BY 3.0](#) (accessed 28 August 2022; colour modified), 441
-  pull by [Pavel N](#) is licensed under [CC BY 3.0](#) (accessed 10 January 2023, modified), 389, 405
-  sardine by [Jaime Serrais](#) licensed under [CC BY 3.0](#) (accessed 28 August 2022) , 445

-  'Waage/Libra' by [B. Lachner](#) is in the public domain (accessed April 2021, edited), 481, 485, 489, 493, 497, 501, 513, 517, 525, 541
-  Tree by [Felipe Alvarado](#) is licensed under [CC BY 3.0](#) (accessed 10 January 2023), 433
-  'Weight' by [Kris Brauer](#) is licensed under [CC-BY](#) (accessed May 2021), 481, 485, 489, 493, 497, 501, 513, 517, 525, 541
-  'boy' by [Xinh Studio](#) is licensed under [CC BY 3.0](#) (accessed 6 June 2023), 485, 489
-  'cookies' by [Azam Ishaq](#) is licensed under [CC BY 3.0](#) (accessed 6 June 2023), 485, 489
-  'cookies' by [Vectors Point](#) is licensed under [CC BY 3.0](#) (accessed 6 June 2023), 485, 489
-  'office desk' by [Abdul Baasith](#) is licensed under [CC BY 3.0](#) (accessed 6 June 2023), 485, 489
-  'Man' by [Xinh Studio](#) is licensed under [CC BY 3.0](#) (accessed 6 June 2023), 485, 489