



TABLE OF CONTENTS




INVERTIBILITY GAME



I'm thinking of a number x . Your hint: $\sin(x) = 0$. What number am I thinking of?

I'm thinking of a number x , and x is between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$. Your hint: $\sin(x) = 0$. What number am I thinking of?

ARCSINE



$\arcsin(x)$ is the inverse of $\sin x$ restricted to $[-\frac{\pi}{2}, \frac{\pi}{2}]$

$\arcsin x$ is the (unique) number θ such that:

- ▶ $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$, and
- ▶ $\sin \theta = x$

ARCSINE

Reference Angles:

θ	$\sin \theta$
0	0
$\frac{\pi}{6}$	$\frac{1}{2}$
$\frac{\pi}{4}$	$\frac{1}{\sqrt{2}}$
$\frac{\pi}{3}$	$\frac{\sqrt{3}}{2}$
$\frac{\pi}{2}$	1

▶ $\arcsin(0)$


▶ $\arcsin\left(\frac{1}{\sqrt{2}}\right)$

▶ $\arcsin\left(-\frac{1}{\sqrt{2}}\right)$

▶ $\arcsin\left(\frac{\pi}{2}\right)$

▶ $\arcsin\left(\frac{\pi}{4}\right)$

ARCCOSINE




$\arccos(x)$ is the inverse of $\cos x$ restricted to $[0, \pi]$.

$\arccos(x)$ is the (unique) number θ such that:

- ▶ $\cos(\theta) = x$ and
- ▶ $0 \leq \theta \leq \pi$

ARCTANGENT



$\arctan(x) = \theta$ means:

- (1) $\tan(\theta) = x$ and
- (2) $-\pi/2 < \theta < \pi/2$

ARCSECANT, ARCSINE, AND ARCCOTANGENT

$$\operatorname{arcsec}(x) =$$

ARCSECANT, ARCSINE, AND ARCCOTANGENT

$$\operatorname{arccsc}(x) = \arcsin\left(\frac{1}{x}\right)$$

$$\operatorname{arccsc}(x) = y$$

$$\csc y = x$$

$$\frac{1}{\sin y} = x$$

$$\sin y = \frac{1}{x}$$

$$y = \arcsin\left(\frac{1}{x}\right)$$

$$\operatorname{arccsc}(x) = \arcsin\left(\frac{1}{x}\right)$$

$$\operatorname{arccot}(x) = \arctan\left(\frac{1}{x}\right)$$

$$\operatorname{arccot}(x) = y$$

$$\cot y = x$$

$$\frac{1}{\tan y} = x$$


$$\tan y = \frac{1}{x}$$

$$y = \arctan\left(\frac{1}{x}\right)$$

$$\operatorname{arccot}(x) = \arctan\left(\frac{1}{x}\right)$$


$$\operatorname{arcsec}(x) = \arccos\left(\frac{1}{x}\right)$$

The domain of $\arccos(y)$ is $-1 \leq y \leq 1$, so the domain of $\operatorname{arcsec}(y)$ is




$$\operatorname{arccsc}(x) = \arcsin\left(\frac{1}{x}\right)$$

Domain of $\arcsin(y)$ is $-1 \leq y \leq 1$, so the domain of $\operatorname{arccsc}(x)$ is



$$\operatorname{arccot}(x) = \arctan\left(\frac{1}{x}\right)$$

Domain of $\arctan(x)$ is all real numbers, so the domain of $\operatorname{arccot}(x)$ is



$$y = \arcsin x$$

Find $\frac{dy}{dx}$.

$$y = \arctan x$$

Find $\frac{dy}{dx}$.

$$y = \arccos x$$

Find $\frac{dy}{dx}$.

To differentiate arcsecant, arccosecant, and arccotangent, you can use the chain rule!

$$\frac{d}{dx} [\operatorname{arccsc}(x)] = \frac{d}{dx} \left[\arcsin \left(\frac{1}{x} \right) \right] = \frac{d}{dx} [\arcsin (x^{-1})]$$

Derivatives of Inverse Trigonometric Functions – Theorem 2.12.7

Memorize:

$$\frac{d}{dx} [\arcsin x] = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} [\arccos x] = -\frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} [\arctan x] = \frac{1}{1+x^2}$$

Be able to derive:

$$\frac{d}{dx} [\operatorname{arccsc} x] = -\frac{1}{|x|\sqrt{x^2-1}}$$

$$\frac{d}{dx} [\operatorname{arcsec} x] = \frac{1}{|x|\sqrt{x^2-1}}$$

$$\frac{d}{dx} [\operatorname{arccot} x] = -\frac{1}{1+x^2}$$