

# PLP - 5

## TOPIC 5 —MODUS PONENS AND PROOFS

Demirbaş & Rechnitzer

# MODUS PONENS — TOWARDS PROOFS

# MODUS PONENS

Once we know an implication  $P \implies Q$  is true, we want to use it.

- If we also know that  $P$  is true
- Then (by our truth table) we must have that  $Q$  is true

This deduction very useful and is precisely why we want to prove implications.

## DEFINITION: (MODUS PONENS).

The deduction:

$(P \implies Q)$  and  $P$  are true so  $Q$  must be true

is called **modus ponens**.

# MODUS PONENS AND PROOFS

Say we need to prove  $P \implies Q$  but cannot directly. Instead show that

- $P \implies P_1$  is true
- $P_1 \implies P_2$  is true
- $P_2 \implies P_3$  is true
- ...
- $P_n \implies Q$  is true

Consider what happens when  $P$  is true (we can ignore  $P$  is false).

- When  $P$  is true, we can use *modus ponens*:
  - Since  $P$  is true and  $P \implies P_1$  is true, we know  $P_1$  is true
  - Since  $P_1$  is true, we know that  $P_2$  is true
  - ...
  - Since  $P_n$  is true, we know that  $Q$  is true
- Hence  $Q$  cannot be false, so the implication is true!

# TYPICAL PROOF STRUCTURE

## PROPOSITION:

If  $P$  then  $Q$

## PROOF.

- Assume that the hypothesis  $P$  is true
- We see that  $P$  implies  $P_1$
- From this we know that  $P_1$  implies  $P_2$
- $\vdots$
- From  $P_n$  we know that  $Q$  as required.