

Note on the equivalence of parabolic Harnack inequalities and heat kernel estimates

Martin T. Barlow, Richard F. Bass and Takashi Kumagai

Version 0.9 (sn8.tex), 8 April 2005

Abstract. The aim of this note is to prove the equivalence of parabolic Harnack inequalities and sub-Gaussian heat kernel estimates in a general metric measure space with a local regular Dirichlet form.

1. Framework and the main theorem.

Let (X, d) be a connected locally compact complete separable metric space. We assume that the metric d is geodesic: for each $x, y \in X$ there exists a (not necessarily unique) geodesic path $\gamma(x, y)$ such that for each $z \in \gamma(x, y)$, we have $d(x, z) + d(z, y) = d(x, y)$. Let μ be a Borel measure on X such that $0 < \mu(B) < \infty$ for every ball B in X . We write $B(x, r) = \{y : d(x, y) < r\}$, and $V(x, r) = \mu(B(x, r))$. Note that under the assumptions above, the closure of $B(x, r)$ is compact for all $x \in X$ and $0 < r < \infty$. For simplicity in what follows, we will also assume that X has infinite diameter, but similar results (with obvious modifications to the statements and the proofs) hold when the diameter of X is finite. We will call such a space a *metric measure space*, or a MM space.

Now let $(\mathcal{E}, \mathcal{F})$ be a regular, strong local Dirichlet form on $L^2(X, \mu)$: see [FOT] for details. We denote by Δ the corresponding self-adjoint operator; that is, we say h is in the domain of Δ and $\Delta h = f$ if $h \in \mathcal{F}$ and $\mathcal{E}(h, g) = -\int fg d\mu$ for every $g \in \mathcal{F}$. Let $\{P_t\}$ be the corresponding semigroup. $(\mathcal{E}, \mathcal{F})$ is called *conservative* (or *stochastically complete*) if $P_t 1 = 1$ for all $t > 0$. Throughout the paper, we assume that $(\mathcal{E}, \mathcal{F})$ is conservative. Since \mathcal{E} is regular, $\mathcal{E}(f, g)$ can be written in terms of a signed measure $\Gamma(f, g)$. To be more precise, for $f \in \mathcal{F}_b$ (the collection \mathcal{F}_b is the set of functions in \mathcal{F} that are essentially bounded) $\Gamma(f, f)$ is the unique smooth Borel measure (called the energy measure) on X satisfying

$$\int_X \tilde{g} d\Gamma(f, f) = 2\mathcal{E}(f, fg) - \mathcal{E}(f^2, g), \quad g \in \mathcal{F}_b, \quad (1.0)$$

where \tilde{g} is the quasi-continuous modification of $g \in \mathcal{F}$. (Recall that $u : X \rightarrow \mathbb{R}$ is called quasi-continuous if for any $\varepsilon > 0$, there exists an open set $G \subset X$ such that $\text{Cap}(G) < \varepsilon$ and $u|_{X \setminus G}$ is continuous. It is known that each $u \in \mathcal{F}$ admits a quasi-continuous modification \tilde{u} – see [FOT], Theorem 2.1.3.) Throughout the paper, we will abuse notation and take the quasi-continuous modification of $g \in \mathcal{F}_b$ without writing \tilde{g} . $\Gamma(f, g)$ is defined by

$$\Gamma(f, g) = \frac{1}{2}(\Gamma(f + g, f + g) - \Gamma(f, f) - \Gamma(g, g)), \quad f, g \in \mathcal{F}.$$

$\Gamma(f, g)$ is also local, linear in f and g , and satisfies the Leibniz and chain rules – see [FOT], p. 115-116. That is, if f_1, \dots, f_m, g , and $\varphi(f_1, \dots, f_m)$ are in \mathcal{F}_b , and φ_i denotes the partial derivative of φ in the i^{th} direction, we have:

$$d\Gamma(fg, h) = f d\Gamma(g, h) + g d\Gamma(f, h),$$

$$d\Gamma(\varphi(f_1, \dots, f_m), g) = \sum_{i=1}^m \varphi_i(f_1, \dots, f_m) d\Gamma(f_i, g).$$

We call (X, d, μ, \mathcal{E}) a metric measure Dirichlet space, or a MMD space.

Let $Y = (Y_t, t \geq 0, \mathbb{P}^x, x \in X)$ be the Hunt process associated with the Dirichlet form \mathcal{E} on $L^2(X, \mu)$ – see [FOT, Theorem 7.2.1]. Since \mathcal{E} is strongly local, by [FOT, Theorem 7.2.2] Y is a diffusion.

Throughout the note, we let $\beta \geq 1$.

Definition 1.1. (a) We say a function u is *harmonic* on a domain D if $u \in \mathcal{F}_{loc}$ and $\mathcal{E}(u, g) = 0$ for all $g \in \mathcal{F}$ with support in D . Here $u \in \mathcal{F}_{loc}$ if and only if for any relatively compact open set G , there exists a function $w \in \mathcal{F}$ such that $u = w$ μ -a.e. on G . See page 117 in [FOT] for the definition of $\mathcal{E}(u, g)$ for $u \in \mathcal{F}_{loc}$ when $(\mathcal{E}, \mathcal{F})$ is a regular, strong local Dirichlet form. Functions in \mathcal{F} are only defined up to quasi-everywhere equivalence; we use a quasi-continuous modification of u . X satisfies the *elliptic Harnack inequality* EHI if there exists a constant c_1 such that, for any ball $B(x, R)$, whenever u is a non-negative harmonic function on $B(x, R)$ then there is a quasi-continuous modification \tilde{u} of u that satisfies

$$\sup_{B(x, R/2)} \tilde{u} \leq c_1 \inf_{B(x, R/2)} \tilde{u}. \quad (\text{EHI})$$

Note that by a standard argument (see, e.g., [M], p. 571) EHI implies that \tilde{u} is Hölder continuous.

(b) Let $Q = Q(x_0, T, R) = (0, 4T) \times B(x_0, 2R) =: I \times B_{2R}$. Let $u(t, x) : Q \rightarrow \mathbb{R}$.

- We define $u_t = \frac{\partial u}{\partial t} \in L^2(dt \times \mu)$ as the derivative in the Schwartz' distribution sense. That is, we define u_t to be the function f in $L^2(dt \times \mu)$ so that for any function $g : Q \rightarrow \mathbb{R}$ such that $g(x, \cdot) \in C_K^\infty(0, 4T)$ for each $x \in B(x_0, 2R)$, $g_t = \frac{\partial g}{\partial t} \in L^2(dt \times \mu)$ for each t , then

$$\int_Q (f(x, t)g(x, t) + u(x, t)g_t(x, t)) dt \mu(dx) = 0.$$

- Let $H(I \rightarrow \mathcal{F}^*)$ be the space of functions $u \in L^2(I \rightarrow \mathcal{F}^*)$ with the distributional time derivative $u_t \in L^2(I \rightarrow \mathcal{F}^*)$ equipped with the norm

$$\left(\int_I \|u(t, \cdot)\|_{\mathcal{F}^*}^2 + \|u_t(t, \cdot)\|_{\mathcal{F}^*}^2 dt \right)^{1/2}.$$

Here we identify $L^2(X, \mu)$ with its own dual and denote the dual of \mathcal{F} by \mathcal{F}^* . So, $\mathcal{F} \subset L^2(X, \mu) \subset \mathcal{F}^*$ with continuous and dense embeddings.

Let $\mathcal{F}(I \times X) = L^2(I \rightarrow \mathcal{F}) \cap H(I \rightarrow \mathcal{F}^*)$ be a Hilbert space with norm

$$\|u\|_{\mathcal{F}(I \times X)} = \left(\int_I \|u(t, \cdot)\|_{\mathcal{F}}^2 + \|u_t(t, \cdot)\|_{\mathcal{F}^*}^2 dt \right)^{1/2}.$$

- We define $\mathcal{F}_{loc}(Q)$ to be the set of $dt \otimes d\mu$ -measurable functions on Q such that for every relatively compact open set $D \subset\subset B_{2R}$ and every open interval $I' \subset\subset I$, there exists a function $u' \in \mathcal{F}(I \times X)$ with $u = u'$ on $I' \times D$. We define

$$\mathcal{F}_c(Q) := \{u \in \mathcal{F}(I \times X) : u(t, \cdot) \text{ has compact support in } B_{2R} \text{ for a.e. } t \in I\}.$$

We say a function $u(t, x) : Q \rightarrow \mathbb{R}$ is a solution of the heat equation in Q if $u \in \mathcal{F}_{loc}(Q)$ and

$$\int_J \left[\int f(t, x) u_t(t, x) \mu(dx) + \mathcal{E}(f(t, \cdot), u(t, \cdot)) \right] dt = 0, \quad \forall J \subset\subset I, \quad \forall f \in \mathcal{F}_c(Q). \quad (1.1)$$

X satisfies the *parabolic Harnack inequality of order β* , $\text{PHI}(\beta)$, if there exists a constant c_2 such that the following holds. Let $x_0 \in X$, $R > 0$, $T = R^\beta$, and $u = u(t, x)$ be a non-negative solution of the heat equation in $Q(x_0, T, R)$. Write $Q_- = (T, 2T) \times B(x_0, R)$ and $Q_+ = (3T, 4T) \times B(x_0, R)$; then there exists $\tilde{u} = \tilde{u}(t, x)$ such that $\tilde{u}(t, \cdot)$ is a quasi-continuous modification of $u(t, \cdot)$ for each t and

$$\sup_{Q_-} \tilde{u} \leq c_2 \inf_{Q_+} \tilde{u}. \quad (\text{PHI}(\beta))$$

Given this PHI, a standard oscillation argument implies that \tilde{u} is jointly continuous.

Let

$$h_\beta(r, t) = \exp \left(- \left(\frac{r^\beta}{t} \right)^{1/(\beta-1)} \right). \quad (1.2)$$

Definition 1.2. We say X satisfies $\text{HK}(\beta)$ if there exists a version of the heat kernel $p_t(x, y)$ on X which satisfies

$$\frac{c_1 h_\beta(c_2 d(x, y), t)}{V(x, t^{1/\beta})} \leq p_t(x, y) \leq \frac{c_3 h_\beta(c_4 d(x, y), t)}{V(x, t^{1/\beta})}, \quad (1.3)$$

for $x, y \in X$ and $t \in (0, \infty)$.

Theorem 1.3. *The following are equivalent:*

- (a) X satisfies $\text{PHI}(\beta)$.
- (b) X satisfies $\text{HK}(\beta)$.

Remark. 1) This equivalence is well-known for manifolds when $\beta = 2$. For MMD with $\beta = 2$, it is indirectly proved in [St2]. (There (a) \iff VD + PI and (b) \iff VD + PI are proved.) For MMD with general time scaling, [HSC] proves the equivalence assuming apriori that solutions to the heat equation are sufficiently regular. (See also [GT2] for the case of an infinite connected weighted graph.) We will prove the equivalence without assuming any apriori condition for solutions to the heat equation.

2) In this note, we only discuss the case when the time scaling exponent is $\beta \geq 1$, but a similar argument gives the equivalence for more general time scalings, for example, when the time scaling function Ψ is $\Psi(R) = R^{\beta_1}$ for $R \leq 1$ and $\Psi(R) = R^{\beta_2}$ for $R \geq 1$ where $\beta_1, \beta_2 \geq 1$.

We give some more definitions for later use.

Definition 1.4. (a) X satisfies volume doubling, VD, if there exists a constant c_1 such that

$$V(x, 2R) \leq c_1 V(x, R) \quad \text{for all } x \in X, R \geq 0. \quad (\text{VD})$$

(b) X satisfies the *Poincaré inequality*, PI(β), if there exists a constant c_2 such that for any ball $B = B(x, R) \subset X$ and $f \in \mathcal{F}$,

$$\int_B (f(x) - \bar{f}_B)^2 d\mu(x) \leq c_2 R^\beta \int_B d\Gamma(f, f). \quad (\text{PI}(\beta))$$

Here $\bar{f}_B = \mu(B)^{-1} \int_B f(x) d\mu(x)$.

(c) X satisfies the condition RES(β) if there exist constants c_3, c_4 such that for any $x_0 \in X$, $R \geq 0$,

$$c_3 \frac{R^\beta}{V(x_0, R)} \leq R_{\text{eff}}(B(x_0, R), B(x_0, 2R)^c) \leq c_4 \frac{R^\beta}{V(x_0, R)}. \quad (\text{RES}(\beta))$$

Here, for A, B which are disjoint subsets of X , we define the effective resistance $R_{\text{eff}}(A, B)$ by

$$R_{\text{eff}}(A, B)^{-1} = \inf \left\{ \int_X d\Gamma(f, f) : f = 0 \text{ on } A \text{ and } f = 1 \text{ on } B, f \in \mathcal{F} \right\}.$$

(d) X satisfies the condition E(β) if for any $x_0 \in X$, $R \geq 0$,

$$c_5 R^\beta \leq \mathbb{E}^{x_0} [\tau_{B(x_0, R)}] \leq c_6 R^\beta, \quad (\text{E}(\beta))$$

where $\tau_A = \inf\{t \geq 0 : Y_t \notin A\}$, Y_t is the strong Markov process associated to the Dirichlet form $(\mathcal{E}, \mathcal{F})$, and \mathbb{E}^{x_0} denotes the expectation starting from the point x_0 .

2. Proof of (b) \Rightarrow (a).

Throughout this section, we assume that (b) holds. Fix $x_0 \in X$ and for $R > 0$, let $B_R := B(x_0, R)$. Let $\mathcal{F}_{B_R} = \{u \in L^2(X, \mu) : u = 0 \text{ } \mu\text{-a.e. on } B_R^c\}$ and consider the part of the Dirichlet form $(\mathcal{E}, \mathcal{F}_{B_R})$ (see [FOT], Section 4.4). Let $\{P_t^{B_R}\}$ be the corresponding semigroup.

Lemma 2.1. *There exists a version of the heat kernel $p_t^{B_R}(x, y)$ for $\{P_t^{B_R}\}$ and, for each $\varepsilon_1, \varepsilon_2 \in (0, 1)$, there exists $c_{\varepsilon_1, \varepsilon_2} > 0$ such that*

$$p_t^{B_R}(x, y) \geq \frac{c_{\varepsilon_1, \varepsilon_2}}{V(x_0, \varepsilon_1 R)},$$

for all $x, y \in B(x_0, \varepsilon_1 R)$ and $\varepsilon_2 R^\beta < t < R^\beta$.

Proof. First, define

$$p_t^{B_R}(x, y) := p_t(x, y) - \mathbb{E}^x[p_{t-\tau_{B_R}}(Y_{\tau_{B_R}}, y), \tau_{B_R} \leq t], \quad (2.1)$$

where Y_t is the diffusion process corresponding to $(\mathcal{E}, \mathcal{F})$ and $\tau_{B_R} = \inf\{t \geq 0 : Y_t \notin B(x_0, R)\}$. Then, it is easy to check, using the strong Markov property, that $p_t^{B_R}(x, y)$ is a version of the heat kernel for $\{P_t^{B_R}\}$. The proof of (2.1) is now a standard argument (see, for example, Lemma 5.1 in [FS]). \square

Let $d\nu = dt \otimes d\mu$, $\mathcal{H} = L^2(\mathbb{R}^1 \times X, d\nu)$ and $\tilde{\mathcal{F}} = \{u : \mathbb{R}^1 \rightarrow \mathcal{F} : \mathcal{A}(u, u) + \|u\|_{\mathcal{H}}^2 < \infty\}$ where $\mathcal{A}(u, u) = \int_{\mathbb{R}^1} \mathcal{E}(u(t, \cdot), u(t, \cdot)) dt$. Let $\tilde{\mathcal{F}}^* = \{u : \mathbb{R}^1 \rightarrow \mathcal{F}^* : \int_{\mathbb{R}^1} \|u(t, \cdot)\|_{\mathcal{F}^*}^2 dt + \|u\|_{\mathcal{H}}^2 < \infty\}$, where \mathcal{F}^* is the dual of \mathcal{F} in the sense $\mathcal{F} \subset L^2(X, \mu) \subset \mathcal{F}^*$. Note that $\tilde{\mathcal{F}} \subset \mathcal{H} = \mathcal{H}^* \subset \tilde{\mathcal{F}}^*$. Let

$$\tilde{\mathcal{W}} = \{u \in \tilde{\mathcal{F}} : \frac{\partial u}{\partial t} \in \tilde{\mathcal{F}}^*\}$$

$$\tilde{\mathcal{E}}(u, v) = (u, \frac{\partial v}{\partial t})_{\nu} + \mathcal{A}(u, v) \quad \text{if } u \in \tilde{\mathcal{F}}, v \in \tilde{\mathcal{W}},$$

where $(u, v)_{\nu} = \int_{\mathbb{R}^1} \int_X uv \, d\mu \, dt$. Let $\{Y_t(x)\}$ be the diffusion process corresponding to $(\mathcal{E}, \mathcal{F})$. Then the semigroup corresponding to $\tilde{\mathcal{E}}$ can be written as $P_t u(t_0, x_0) = \mathbb{E}[u(t_0 + t, Y_t(x_0))]$ so that the corresponding generator is $\frac{\partial}{\partial t} + \mathcal{L}$ (the corresponding diffusion is $Z_t = (t, Y_t)$), whereas the dual semigroup $\{\hat{P}_t\}$ can be written as $\hat{P}_t u(t_0, x_0) = \mathbb{E}[u(t_0 - t, Y_t(x_0))]$ and the corresponding generator is $-\frac{\partial}{\partial t} + \mathcal{L}$. (See [O] for details.)

Lemma 2.2. *Let u be a non-negative solution of the heat equation on $Q := I \times G$, where $I = (a, b)$ and G is an open connected subset of X . Then $u(t, x) \geq \int p_{t-s}^B(x, y) u(s, y) d\mu(y)$ μ -a.e. x and all $0 < s < t$ where $B \subset\subset G$.*

Proof. The claim is equivalent to $(u - \hat{P}_{t-s}^Q u)(t, x) \geq 0$ for all $(t, x) \in Q$ and all $0 < s < t$.

Let $\alpha > 0$. Then, $\tilde{\mathcal{E}}_{\alpha}(u, g) \geq 0$ for all non-negative $g \in \tilde{\mathcal{F}}_Q$. So, for any non-negative α -excessive function (w.r.t. $(\tilde{\mathcal{E}}, \tilde{\mathcal{F}}_Q)$ –see [O], Section 4.3, for a discussion of excessive functions in the parabolic case) $v \in \mathcal{F}_Q$, we have

$$\begin{aligned} (u - e^{-\alpha s} \hat{P}_s^Q u, v)_{\nu} &= (u, v - e^{-\alpha s} P_s^Q v)_{\nu} \\ &= (u_Q, v - e^{-\alpha s} P_s^Q v)_{\nu} + (H_Q^{\alpha} u, v - e^{-\alpha s} P_s^Q v)_{\nu} \\ &\geq (u_Q, v - e^{-\alpha s} P_s^Q v)_{\nu} = \tilde{\mathcal{E}}_{\alpha}(u, G_{\alpha}^Q v - e^{-\alpha s} P_s^Q G_{\alpha}^Q v)_{\nu} =: I_1, \end{aligned}$$

where $u = u_Q + H_Q^{\alpha} u$ is the orthogonal decomposition of u into $\mathcal{F}_Q \oplus \mathcal{H}_{Q^c}^{\alpha}$ (see p. 149 of [FOT] –the same proof works for the parabolic case). Here the inequality in the third line is because $H_Q^{\alpha} u(x) = E^x(e^{-\alpha \sigma_{Q^c}} u(Z_{\sigma_{Q^c}})) \geq 0$ (due to Lemma 5.1.3 in p. 105 of [O]) and the fact that v is α -excessive (the definition of excessive functions in [O] is different from that in [FOT], but the proof of Theorem 2.2.1 in [FOT] also establishes equivalent conditions for the parabolic case, too). Since $G_{\alpha}^Q v - e^{-\alpha s} P_s^Q G_{\alpha}^Q v = \int_0^s e^{-\alpha l} P_l^Q v \, dl \in \mathcal{F}_Q$ is non-negative on Q , $I_1 \geq 0$. Thus $u - e^{-\alpha s} \hat{P}_s^Q u \geq 0$ on Q . Since this holds for all $\alpha > 0$, we have $u \geq \hat{P}_s^Q u$ on Q . \square

Once these properties are established, then proving (a) is standard; prove the oscillation inequality first and then use the inequality to establish PHI. Indeed, the proof of Lemma 5.2 and Theorem 5.4 in [FS] work line by line, with suitable changes of the scaling exponents.

3. Proof of (a) \Rightarrow (b).

There is a standard argument, given in [SC1] and Section 5.5 of [SC2] which proves that $\text{PHI}(\beta)$ implies VD , $\text{PI}(\beta)$, and $\text{HK}(\beta)$. See also [HSC] for the case $\beta \neq 2$. However, as this argument uses existence and regularity of caloric and harmonic functions, we will give more complete details of the initial stages of this argument.

First, if $f \in L^2(X, \mu)$ we have that $P_t f \in \mathcal{D}(\mathcal{L})$, and $v(t, x) = P_t f(x)$ is a solution to the heat equation in $X \times (0, \infty)$. Let $x \in X$, $t > 0$, $r = t^\beta$ and $f \geq 0$ with $\int f = 1$. Then applying $\text{PHI}(\beta)$ in $Q = (0, 4t) \times B(x, 2r)$ we obtain

$$\sup_{Q_-} \tilde{v} \leq C \inf_{Q_+} \tilde{v}.$$

Hence if $B = B(x, r)$ then since $\int P_s f = 1$

$$\mu(B) \sup_{Q_-} \tilde{v} \leq C \int_B v(2t, y) \mu(dy) \leq C.$$

Thus for each $x \in X$ we have

$$\widetilde{P_t f}(x) \leq c(t) \|f\|_1. \quad (3.1)$$

Given (3.1), we can use the same arguments as in p. 52 of [B] (using the results in [Y]) to deduce the existence of a transition density $p_t(x, y)$.

Lemma 3.1. *There exist an exceptional set N and a jointly measurable transition density $p_t(x, y)$, $t > 0$, $x, y \in (X \setminus N) \times (X \setminus N)$, such that*

$$\begin{aligned} P_t(x, A) &= \int_A p_t(x, y) \mu(dy) \text{ for } x \in X \setminus N, \quad t > 0, A \in \mathcal{B}(X \setminus N), \\ p_t(x, y) &= p_t(y, x) \text{ for all } x, y, t, \\ p_{t+s}(x, z) &= \int p_s(x, y) p_t(y, z) \mu(dy) \quad \text{for all } x, z, t, s. \end{aligned}$$

Since $p_t(x, y) = P_{t/2} p_{t/2}(\cdot, y)(x)$ it follows that $p_t(\cdot, y)$ is a solution of the heat equation. Now take a quasi continuous modification $\tilde{p}_t(x, y)$ w.r.t. x and use it in the procedure of (4) in [Y]. Then, by Theorem 1 in [Y], there exists $p_t(x, y)$ which is quasi continuous and satisfies the three equalities in Lemma 3.1. (In fact, the uniqueness criteria in Theorem 1 in [Y] shows that this $p_t(x, y)$ is the same as the original one.) Thus it satisfies the PHI, and so can be extended to $(0, \infty) \times X \times X$ as a jointly continuous function.

We now sketch the argument that $\text{PHI}(\beta)$ implies VD , $\text{PI}(\beta)$, and $\text{HK}(\beta)$. We begin with VD , which also gives a key lower bound on the transition density for the killed process.

Applying the PHI to the function $u(t, x) = p_t(x_0, x)$ in the region $Q(x_0, 0, R)$ we obtain (writing $T = R^\beta$)

$$p_{2T}(x_0, x_0) \leq c p_{4T}(x_0, y), \quad y \in B(x_0, R).$$

Integrating over $B(x_0, R)$ gives

$$p_{2T}(x_0, x_0)V(x_0, R) \leq c \int_{B(x_0, R)} u(4T, y) \leq c, \quad (3.2)$$

which gives an upper bound on $p_{2T}(x_0, x_0)$ in terms of the volume of balls.

To obtain a lower bound, write $B_\lambda = B(x_0, \lambda R)$, and let $\varphi \in \mathcal{F}$ be a cut-off function for $B_{5/2} \subset B_3$. Let $p_t^0(x, y)$ be the heat kernel for the process Y killed on exiting B_4 . Define

$$u(t, x) = \begin{cases} \varphi(x), & x \in B_2, 0 < t \leq 2T, \\ \int_{B_3} p_{t-2T}^0(x, y)\varphi(y)\mu(dy), & x \in B_2, 2T < t \leq 4T. \end{cases}$$

Lemma 3.2. *u is a solution of the heat equation in $Q(x_0, T, R)$.*

Proof. The function $u_t(x, t) = \frac{\partial u}{\partial t}$ exists for $t > 2T$, and is zero for $t < 2T$. Since $u(x, t)$ is continuous at $t = 2T$ for $x \in B$, it is straightforward to check that u_t is the derivative of u in the Schwartz' distribution sense.

Since we have $u(t, \cdot) \in \mathcal{D}(\mathcal{L})$ for all $t > 2T$, we have for $f \in \mathcal{F} \cap C(X)$ with support in B_2 that

$$\int f u_t d\mu = -\mathcal{E}(f, u(t, \cdot)), \quad t > 2T. \quad (3.3)$$

If $t < 2T$ then since $u = 1$ on B_2 (3.3) also holds for $t < 2T$. Thus it follows that (1.1) holds. \square

We can now, as in [SC1], [SC2], [HSC], use $\text{PHI}(\beta)$ in $Q(x_0, 0, R)$ to obtain

$$1 = u(y, 2T) \leq cu(x_0, 4T) \leq c \int_{B_3} p_{2T}^0(x_0, y), \quad y \in B(x_0, R). \quad (3.4)$$

Using the PHI in a chain of regions $Q(y_i, t_i, r) \subset [0, 4T] \times B(x_0, 4R)$ we obtain

$$p_{2T}^0(x_0, y') \leq cp_{4T}^0(x_0, y), \quad y' \in B(x_0, 3R), y \in B(x_0, R). \quad (3.5)$$

Integrating (3.5) over $y' \in B_3$ gives

$$\int_{B_3} p_{2T}^0(x_0, y')\mu(dy') \leq cp_{4T}^0(x_0, x_0)V(x_0, 3R), \quad (3.6)$$

and combining (3.4) and (3.6) we deduce that

$$V(x_0, 3R)^{-1} \leq cp_{4T}^0(x_0, y), \quad y \in B(x_0, R). \quad (3.7)$$

The inequalities (3.2) and (3.7) control $p_t(x_0, x_0)$ from above and below in terms of the volume of balls, and since $t \rightarrow p_t(x_0, x_0)$ is decreasing one easily deduces, by the same arguments as in [SC2], that volume doubling holds.

Given the lower bound (3.7), the proof of $\text{HK}(\beta)$ now follows as in Section 5 of [HSC]. For the global lower bound one uses (3.7) and a standard chaining argument. (3.7) gives uniform control of the probability that Y exits a ball radius r before time $t = r^\beta$, and using this the upper bounds on $p_t(x, y)$ follow as in p. 1472–1475 of [HSC].

We remark that (3.7) also gives a lower bound on the transition density of the process Y reflected at ∂B (see [Ch]). Using this the argument of [SC1] can be used to obtain $\text{PI}(\beta)$.

4. Appendix: Proof of $\mathbf{VD} + \mathbf{EHI} + \mathbf{RES}(\beta) \Rightarrow \mathbf{VD} + \mathbf{EHI} + \mathbf{E}(\beta)$.

In this appendix, we modify the proof in [GT2] and prove $\mathbf{VD} + \mathbf{EHI} + \mathbf{RES}(\beta) \Rightarrow \mathbf{E}(\beta)$ in a general MMD framework. This fact is used in [BBK] Theorem 2.15.

Recall from [FOT, Section 1.6] the definition of invariant sets and an irreducible Dirichlet form.

Lemma 4.1. *Let X satisfy EHI. Then \mathcal{E} is irreducible.*

Proof. Let A be an irreducible set, and suppose both $\mu(A) > 0$ and $\mu(A^c) > 0$. Then there exists a ball $B = B(x, R)$ with $\mu(A \cap B') > 0$ and $\mu(A^c \cap B') > 0$, where $B' = B(x, R/2)$. Since $P_t 1_A = 1_A$ it follows that $u = 1_A$ and $v = 1_{A^c}$ are harmonic on B . So by EHI we have

$$\tilde{u}(x) \leq C\tilde{u}(y), \quad x, y \in B'.$$

Since $u > 0$ on a set of positive measure, we have that there exists $x \in B'$ with $\tilde{u}(x) > 0$; hence by the EHI $\tilde{u} > 0$ on B' . But as $\tilde{u} = 1_A$ μ -a.e., we deduce that $\mu(A^c \cap B') = 0$, a contradiction. \square

Proposition 4.2. *Let X satisfy EHI, and $B = B(x, R)$. Then $Gg < \infty$ on B if $g \in L_+^1(B)$.*

Proof. (sketch). Consider the Dirichlet form \mathcal{E}_B with domain $\mathcal{F}_B = \{f \in \mathcal{F} : f|_{B^c} = 0\}$. Let $A = B(x, R/2)$ and $h(x) = P^x(T_A < \tau_B)$. Then h is excessive with respect to \mathcal{E}_B . If h were constant on B then we would have $h = 1$ on B , and the set B would be an invariant set for \mathcal{E} . Thus h is non-constant.

So by [BG, Ex. (4.22), p. 89] we deduce that the killed semigroup P_t^B is transient. Hence (see [FOT, Section 1.6]) we have $Gg < \infty$ for any $g \in L_+^1(B, \mu)$. \square

Lemma 4.3. *Let D be a bounded domain in X . Then EHI implies that there exists the Green density $g^D(\cdot, \cdot)$ which is continuous on $(X \times X) \setminus \Delta$ and $g_D(x, y) = g_D(y, x)$ for all $x, y \in (X \times X) \setminus \Delta$, where Δ is the diagonal. Further, there exists $C > 0$ such that for any $r > 0$, if $y_0, y_1 \in X$ satisfy $d(y_0, y_1) \geq 2r$, then*

$$g_D(y_0, x) \leq Cg_D(y_0, y) \quad \forall x, y \in B(y_1, r). \quad (4.1)$$

Proof. Let $x_0, x_1 \in D$, Choose $r > 0$ such that $B(x_i, 2r) \subset D$, $B(x_0, 2r) \cap B(x_1, 2r) = \emptyset$. Write $B_i = B(x_i, 2r)$, $B'_i = B(x_i, r)$. Let $f, g \in \mathcal{F}$ with supports in B'_0 and B'_1 , and $\int f = \int g = 1$. Let G_D be the Green operator for the process Y killed on exiting D . By Proposition 4.2 we have $G_D f < \infty$, $G_D g < \infty$.

Then if $u \in \mathcal{F}$ with $\text{Supp } u \subset B(x_1, 2r)$,

$$\mathcal{E}(G_D f, u) = (f, u) = 0, \quad (4.2)$$

so $G_D f$ is harmonic on B_1 . Similarly $G_D g$ is harmonic on B_0 . By the EHI if $x \in B'_1$ then

$$G_D f(x) \leq CG_D f(y), \quad y \in B'_1. \quad (4.3)$$

Similarly

$$G_D g(x) \leq C G_D g(x_0), \quad x \in B'_0.$$

So

$$G_D f(x_1) \leq C(g, G_D f) = C(G_D g, f) \leq C^2 G_D g(x_0).$$

Now fix g such that $C_1 = G_D g(x_0) < \infty$ – such a g exists by choosing $g \leq ch_0$. Then we have $G_D f(x_1) \leq c' \|f\|_1$ for all f with support in B'_0 . Therefore the kernel $G_D(x_1, dx)$ has a density $g_D(x_1, y)$ on B'_0 . Since $(f, G_D g) = (G_D f, g)$ for $f, g \in L^2$, it follows that $g_D(x, y) = g_D(y, x)$ $\mu \times \mu$ -a.e.

Now, take $y_0, y_1 \in X$ that satisfy $d(y_0, y_1) \geq 2r$. For any $\epsilon > 0$ and $f \in L^2$ with support in $B(y_0, \epsilon r)$, similarly to (4.2) we can show that $G_D f$ is harmonic on $B(y_1, (2 - \epsilon)r)$. Thus, by the same way as (4.3), we have

$$G_D f(x) \leq C G_D f(y), \quad x, y \in B(y_1, r). \quad (4.4)$$

Now let $f_n(z) = V(y_0, r_n)^{-1} 1_{B(y_0, r_n)}(z)$ where $\epsilon r \geq r_n \downarrow 0$. Applying (4.4) to f_n and take $n \rightarrow \infty$, we obtain (4.1) for μ -a.e. y_0 . By the usual oscillation argument, we can deduce that $g_D(x, y)$ is continuous on $(X \times X) \setminus \Delta$. Especially, $g_D(x, y) = g_D(y, x)$ for all $x, y \in (X \times X) \setminus \Delta$. We thus obtain (4.1) for all $y_0 \in X$. \square

Now let $M \geq 2$ be fixed. (In fact, we can take $M=2$.)

Definition 4.4. $(\mathcal{E}, \mathcal{F})$ satisfies (HG) if there exists a constant $c_1 > 0$ such that for any ball $B(x_0, R)$, there exists the Green kernel $g^{B_R}(x_0, y)$ and for any $0 < r \leq R/M$, we have

$$\sup_{y \notin B(x_0, r)} g^{B_R}(x_0, y) \leq c_1 \inf_{y \in B(x_0, r)} g^{B_R}(x_0, y). \quad (HG)$$

Lemma 4.5. $(EHI) \Rightarrow (HG)$.

Proof. Given Lemma 4.3 this is the same argument as in [B]. We prove that if $d(x_0, x) = d(x_0, y) = R$, and $B(x_0, 2R) \subset D$ then

$$C_1^{-1} g_D(x_0, y) \leq g_D(x_0, x) \leq C_1 g_D(x_0, y). \quad (4.5)$$

Once (4.5) is proved, then (HG) holds by the maximum principle (which holds for $G_D f$ and so for g_D as well). By symmetry it is enough to prove the right hand inequality of (4.5).

Let x', y' be the midpoints of $\gamma(x_0, x)$, and $\gamma(x_0, y)$. Thus $d(x_0, x') = d(x_0, y') = R/2$. Clearly we have $d(x', y) \geq R/2$ and $d(x, y') \geq R/2$.

We now consider two cases.

Case 1. $d(x', y') \leq R/3$. Let z be the midpoint of $\gamma(x', y')$. Then $d(z, x') \leq R/6 \leq R/4$. So applying (4.1) to $g_D(x_0, \cdot)$ in $B(x', R/4) \subset B(x', R/2)$, we deduce that

$$C_2^{-1} g_D(x_0, x') \leq g_D(x_0, z) \leq C_2 g_D(x_0, x'). \quad (4.6)$$

Now apply (4.1) to $g_D(x_0, \cdot)$ in $B(x, R/2) \subset B(x, R)$, to deduce that

$$C_2^{-1} g_D(x_0, x) \leq g_D(x_0, x') \leq C_2 g_D(x_0, x).$$

Combining these inequalities we deduce that

$$C_2^{-2} g_D(x_0, x) \leq g_D(x_0, z) \leq C_2^2 g_D(x_0, x),$$

and this, with a similar inequality for $g_D(x_0, y)$, proves (4.5).

Case 2. $d(x', y') > R/3$. Apply (4.1) to $g_D(y, \cdot)$ in $B(x_0, R/2) \subset B(x_0, R)$, to deduce that

$$C_2^{-1} g_D(y, x') \leq g_D(y, x_0) \leq C_2 g_D(y, x'). \quad (4.7)$$

Now look at $g_D(x', \cdot)$. If z' is on $\gamma(y', y)$ with $d(y', z') = s \in [0, R/2]$ then as $d(x', y') > R/3$ and $d(x', y) \geq R/2$ we have $d(x', z') \geq \max(R/3 - s, s)$. Hence we deduce $d(x', z') \geq R/6$. So applying (4.1) repeatedly to $g_D(x', \cdot)$ for a chain of balls $B(z', R/12) \subset B(z', R/6)$ we deduce that

$$C_2^{-6} g_D(x', y') \leq g_D(x', y) \leq C_2^6 g_D(x', y'). \quad (4.8)$$

So, we obtain from (4.7) and (4.8),

$$g_D(y, x_0) \leq C_2 g_D(y, x') \leq C_2^7 g_D(x', y'), \quad g_D(x', y') \leq C_2^6 g_D(y, x') \leq C_2^7 g_D(y, x_0).$$

We have similar inequalities relating $g_D(x, x_0)$ and $g_D(x', y')$, which proves (4.5). \square

Lemma 4.6. Assume that $(\mathcal{E}, \mathcal{F})$ satisfies (HG).

1) For any ball $B(x_0, R)$ and for any $0 < r \leq R/M$, we have

$$\sup_{y \notin B(x_0, r)} g^{B_R}(x_0, y) \asymp R(B_r, B_R^c) \asymp \inf_{y \in B(x_0, r)} g^{B_R}(x_0, y). \quad (4.9)$$

2) Let $B_k = B(x_0, M^k r)$ for $k = 0, 1, \dots$. Then, for any integers $0 \leq m < n$,

$$\sup_{y \notin B_m} g^{B_n}(x_0, y) \asymp \sum_{k=m}^{n-1} R(B_k, B_{k+1}^c) \asymp \inf_{y \in B_m} g^{B_n}(x_0, y). \quad (4.10)$$

Proof. For 1), first the following is standard (see for example (4.7) in [GT2]).

$$\sup_{y \notin B(x_0, r)} g^{B_R}(x_0, y) \geq R(B_r, B_R^c) \geq \inf_{y \in B(x_0, r)} g^{B_R}(x_0, y).$$

Thus, using (HG), we obtain (4.9).

For 2), note first that the following holds by the definition of resistance

$$\sum_{k=m}^{n-1} R(B_k, B_{k+1}^c) \leq R(B_m, B_n^c).$$

This and (4.9) implies the lower bound for $\inf g^{B_n}$ in (4.10). Next, by the reproducing property of g^{B_k} , we know that $g^{B_{k+1}}(x, \cdot) - g^{B_k}(x, \cdot)$ is a harmonic function in B_k . Thus,

$$g^{B_{k+1}}(x, y) - g^{B_k}(x, y) \leq \sup_{z \notin B_k} g^{B_{k+1}}(x, z) \leq cR(B_k, B_{k+1}), \quad \forall y \in X, \quad (4.11)$$

where the first inequality is by the maximum principle and the second inequality is by (4.9). For $y \notin B_m$, by (4.9)

$$g^{B_{m+1}}(x, y) \leq c'R(B_m, B_{m+1}). \quad (4.12)$$

For such y , adding up (4.12) with (4.11) for $m < k < n$, we obtain the upper bound of $\sup g^{B_n}$ in (4.10). \square

Proof of $VD + EHI + RES(\beta) \Rightarrow E(\beta)$.

$$\begin{aligned} E^{x_0}[\tau_{B_R}] &= \int g^{B_R}(x_0, y) d\mu(y) \geq \int_{B(x_0, r)} g^{B_R}(x_0, y) d\mu(y) \\ &\geq cR(B_r, B_R^c)V(x_0, r) \geq cR^\beta, \end{aligned}$$

where we used Lemma 4.6 1) in the second inequality and $VD + RES(\beta)$ in the last inequality.

Now, for each $k \in \mathbb{Z}$, let $r_k = M^k$, $B_k = B(x_0, r_k)$ and let n_0 be the minimum number such that $R < r_{n_0}$. Then

$$\begin{aligned} E^{x_0}[\tau_{B_R}] &\leq E^{x_0}[\tau_{B(x_0, r_{n_0})}] = \int_{B_{n_0}} g^{B_{n_0}}(x_0, y) d\mu(y) \\ &= \sum_{m=-\infty}^{n_0-1} \int_{B_{m+1} \setminus B_m} g^{B_m}(x_0, y) d\mu(y) \leq c \sum_{m=-\infty}^{n_0-1} \left(\sum_{k=m}^{n_0-1} R(B_k, B_{k+1}^c) \right) \mu(B_{m+1} \setminus B_m) \\ &= c \sum_{k=-\infty}^{n_0-1} \left(\sum_{m=-\infty}^k \mu(B_{m+1} \setminus B_m) \right) R(B_k, B_{k+1}^c) = c \sum_{k=-\infty}^{n_0-1} \mu(B_{k+1}) R(B_k, B_{k+1}^c) \\ &\leq c' \sum_{k=-\infty}^{n_0-1} r_{k+1}^\beta \leq c'' R^\beta, \end{aligned}$$

where we used Lemma 4.6 2) in the second inequality and $VD + RES(\beta)$ in the third inequality. We thus obtain $E(\beta)$. \square

Acknowledgment. The authors thank Z.Q. Chen, K.T. Sturm and M. Takeda for valuable comments.

References

- [B] M.T. Barlow. Some remarks on the elliptic Harnack inequality, *Bull. London Math. Soc.*, to appear.
- [BBK] M.T. Barlow, R.F. Bass, T. Kumagai. Stability of parabolic Harnack inequalities on metric measure spaces, Preprint.
- [BG] R.M. Blumenthal, R.K. Gettoor. *Markov processes and potential theory*. Pure and Applied Mathematics, Vol. 29 Academic Press, New York-London 1968.
- [Ch] Z.Q. Chen. On reflected Dirichlet spaces. *Probab. Theory Rel. Fields* **94** (1992), no. 2, 135–162.
- [FS] E.B. Fabes, D.W. Stroock. A new proof of Moser’s parabolic Harnack inequality using the old ideas of Nash. *Arch. Mech. Rat. Anal.* **96** (1986), 327–338.
- [FOT] M. Fukushima, Y. Oshima, M. Takeda. *Dirichlet Forms and Symmetric Markov Processes*. de Gruyter, Berlin, 1994.
- [GT2] A. Grigor’yan, A. Telcs. Harnack inequalities and sub-Gaussian estimates for random walks. *Math. Annalen* **324** (2002), 521–556.
- [HSC] W. Hebisch, L. Saloff-Coste. On the relation between elliptic and parabolic Harnack inequalities. *Ann. Inst. Fourier (Grenoble)* **51** (2001), 1437–1481.
- [M] J. Moser, On Harnack’s theorem for elliptic differential equations. *Comm. Pure Appl. Math.* **14** (1961), 577–591.
- [O] Y. Oshima. Lecture notes on Dirichlet forms. Unpublished lecture notes.
- [SC1] L. Saloff-Coste. A note on Poincaré, Sobolev, and Harnack inequalities. *Inter. Math. Res. Notices* **2** (1992), 27–38.
- [SC2] L. Saloff-Coste. *Aspects of Sobolev-Type inequalities*. Lond. Math. Soc. Lect. Notes **289**, Cambridge Univ. Press, 2002.
- [St2] K.-T. Sturm. Analysis on local Dirichlet spaces -III. The parabolic Harnack inequality. *J. Math. Pures Appl.* **75** (1996), no. 9, 273–297.
- [Y1] J.-A. Yan. A formula for densities of transition functions. *Sem. Prob. XXII*, 92–100. Lect Notes Math. **1321**.

MTB: Department of Mathematics, University of British Columbia, Vancouver V6T 1Z2, Canada

RFB: Department of Mathematics, University of Connecticut, Storrs, CT 06269-3009, USA

TK: Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606-8502, Japan