

---

**Math 405: Numerical Methods for Differential Equations 2016 W1**  
**Topic 4b: Composite Quadrature**

---

See Chapter 7 of Süli and Mayers.

**Motivation:** we've seen oscillations in polynomial interpolation—the Runge phenomenon—for high-degree polynomials.

**Idea:** split a required integration interval  $[a, b] = [x_0, x_n]$  into  $n$  equal intervals  $[x_{i-1}, x_i]$  for  $i = 1, \dots, n$ . Then use a **composite rule**:

$$\int_a^b f(x) dx = \int_{x_0}^{x_n} f(x) dx = \sum_{i=1}^n \int_{x_{i-1}}^{x_i} f(x) dx$$

in which each  $\int_{x_{i-1}}^{x_i} f(x) dx$  is approximated by quadrature.

Thus rather than increasing the degree of the polynomials to attain high accuracy, instead increase the number of intervals.

**Trapezium Rule:**

$$\int_{x_{i-1}}^{x_i} f(x) dx = \frac{h}{2}[f(x_{i-1}) + f(x_i)] - \frac{h^3}{12}f''(\xi_i), \quad \text{for some } \xi_i \in (x_{i-1}, x_i).$$

**Composite Trapezium Rule:**

$$\begin{aligned} \int_{x_0}^{x_n} f(x) dx &= \sum_{i=1}^n \left[ \frac{h}{2}[f(x_{i-1}) + f(x_i)] - \frac{h^3}{12}f''(\xi_i) \right] \\ &= \frac{h}{2}[f(x_0) + 2f(x_1) + 2f(x_2) + \dots + 2f(x_{n-1}) + f(x_n)] + e_h^T \end{aligned}$$

where  $\xi_i \in (x_{i-1}, x_i)$  and  $h = x_i - x_{i-1} = (x_n - x_0)/n = (b - a)/n$ , and the error  $e_h^T$  is given by

$$e_h^T = -\frac{h^3}{12} \sum_{i=1}^n f''(\xi_i) = -\frac{nh^3}{12} f''(\xi) = -(b - a) \frac{h^2}{12} f''(\xi)$$

for some  $\xi \in (a, b)$ , using the Intermediate-Value Theorem  $n$  times. Note that if we halve the stepsize  $h$  by introducing a new point halfway between each current pair  $(x_{i-1}, x_i)$ , the factor  $h^2$  in the error should decrease by four.

**Another composite rule:** if  $[a, b] = [x_0, x_{2n}]$ ,

$$\int_a^b f(x) dx = \int_{x_0}^{x_{2n}} f(x) dx = \sum_{i=1}^n \int_{x_{2i-2}}^{x_{2i}} f(x) dx$$

in which each  $\int_{x_{2i-2}}^{x_{2i}} f(x) dx$  is approximated by quadrature.

**Simpson's Rule:**

$$\int_{x_{2i-2}}^{x_{2i}} f(x) dx = \frac{h}{3}[f(x_{2i-2}) + 4f(x_{2i-1}) + f(x_{2i})] - \frac{(2h)^5}{2880}f''''(\xi_i), \quad \text{for some } \xi_i \in (x_{2i-2}, x_{2i}).$$

---

### Composite Simpson's Rule:

$$\begin{aligned}\int_{x_0}^{x_{2n}} f(x) dx &= \sum_{i=1}^n \left[ \frac{h}{3} [f(x_{2i-2}) + 4f(x_{2i-1}) + f(x_{2i})] - \frac{(2h)^5}{2880} f''''(\xi_i) \right] \\ &= \frac{h}{3} [f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + 2f(x_4) + \cdots \\ &\quad + 2f(x_{2n-2}) + 4f(x_{2n-1}) + f(x_{2n})] + e_h^S\end{aligned}$$

where  $\xi_i \in (x_{2i-2}, x_{2i})$  and  $h = x_i - x_{i-1} = (x_{2n} - x_0)/2n = (b - a)/2n$ , and the error  $e_h^S$  is given by

$$e_h^S = -\frac{(2h)^5}{2880} \sum_{i=1}^n f''''(\xi_i) = -\frac{n(2h)^5}{2880} f''''(\xi) = -(b-a) \frac{h^4}{180} f''''(\xi)$$

for some  $\xi \in (a, b)$ , using the Intermediate-Value Theorem  $n$  times. Note that if we halve the stepsize  $h$  by introducing a new point half way between each current pair  $(x_{i-1}, x_i)$ , the factor  $h^4$  in the error should decrease by sixteen (assuming  $f$  is smooth enough).

**Adaptive (or automatic) procedure:** if  $S_h$  is the value given by Simpson's rule with a stepsize  $h$ , then

$$S_h - S_{\frac{1}{2}h} \approx -\frac{15}{16} e_h^S.$$

This suggests that if we wish to compute  $\int_a^b f(x) dx$  with an absolute error  $\varepsilon$ , we should compute the sequence  $S_h, S_{\frac{1}{2}h}, S_{\frac{1}{4}h}, \dots$  and stop when the difference, in absolute value, between two consecutive values is smaller than  $\frac{16}{15}\varepsilon$ . That will ensure that (approximately)  $|e_h^S| \leq \varepsilon$ .

Often spatially-varying adaptivity is used in practice: refine only in regions where a local estimate is large.

### Comments:

Sometimes much better accuracy may be obtained using the Trapezoidal Rule: for example, as might happen when computing Fourier coefficients, if  $f$  is periodic with period  $b - a$  so that  $f(a + x) = f(b + x)$  for all  $x$ .