

## Homework 2

*Like the first one, you can work on this alone or with others as you prefer. We ask you to copy a few of your answers into VISTA; this will count towards your participation for the week only and our purpose in doing so is to have you focus on understanding this week's topic and give us feedback that we can use to change the content of Friday's class.*

In this worksheet, you asked to use an interactive figure for linear approximation to see how the formula relates to the graph and tangent line without drawing a new picture for each point you want to estimate. If you are not sure how to get started, ask on the Vista Discussion Board or in the Math Learning Centre.

Visit the website <http://calculusapplets.com/linearapprox.html> and make sure the interactive graph is visible (you may have to "allow" the applet to run in your browser; you probably do not need to update Java even if it asks for this). You can adjust the viewable area of the graph using the controls on the right side, and you can adjust the values of  $a$  and  $x$  separately.

The interactive Figure 4.63 through MyMathLab has a similar picture, and you can get started on the questions with any figure that looks like Figure 4.63 in your book and confirm later with the applet if you like.

**Some of these are questions asked on VISTA. Copy your answers there when done.**

1. What happens to the tangent line when you vary  $a$  in the figure? What happens to the estimate?
2. What happens to the tangent line when you vary  $x$  in the picture? What happens to the estimate?
3. For  $f(x) = x^2$  (the default), do you see underestimates or overestimates using linear approximation? How is this different from the square root function we looked at in class, and why?
4. Switch to the sine function: there is a drop-down menu at the top of the applet with this as a choice, or you can enter in  $\sin(x)$  for  $f$  directly just under the picture.

We introduced the error bound term  $\frac{M}{2}(x-a)^2$  for the linear approximation  $L_a(x) = f(a) + f'(a)(x-a)$

where  $M$  is a strange kind of constant; if  $M$  is "big enough" then we have a guarantee that

$$|\text{error}| < \frac{M}{2}(x-a)^2$$

One  $M$  that is "big enough" is the maximum absolute value of the second derivative for all the  $x$  we care

about: if  $|f''(z)| < M$  is true for all  $z$  near  $a$  that we care about (our estimate at  $x$  would be included in that interval of values). For example,  $f(x) = \sin(x)$  has  $f''(x) = -\sin(x)$ , and since the sine function varies from -1 to 1,  $-1 < f''(x) < 1$  or  $|f''(z)| < 1$ . So in this case,  $M = 1$  is "big enough", and we can conclude that  $|error| < \frac{1}{2}(x-a)^2$ . Try setting  $a = 0$  and vary  $x$ . Why does the error get so large as  $x$  moves away from  $a$ ?

5. If you want to estimate values between  $\sqrt{3}$  and  $\sqrt{5}$  (the square roots of numbers like 3, 3.01, 3.6, 4.2, 4.345) using linear approximation to  $f(x) = \sqrt{x}$ :

- What is a good choice for  $a$  for this case?
- If I tell you that  $-0.05 < f''(x) < -0.01$  for  $x$  between 3 and 5, what is a value of  $M$  that works in the error bound formula?
- Sketch a picture (you may have room below) of this situation that would help you explain it to somebody else and bring to Friday's class.

**Don't forget to enter your answers on Vista! That will help for Friday's class.**