

# Disjoint Cycles and Equitable Colorings

H. Kierstead   A. Kostochka   T. Molla   E. Yeager

*yeager2@illinois.edu*

Eau Claire, Wisconsin  
Special Session on Graph and Hypergraph Theory  
20 September 2014

# Disjoint Cycles

# Corrádi-Hajnal Theorem

Corrádi-Hajnal, 1963

If  $G$  is a graph on  $n$  vertices with  $n \geq 3k$  and  $\delta(G) \geq 2k$ , then  $G$  contains  $k$  disjoint cycles.

# Corrádi-Hajnal Theorem

Corrádi-Hajnal, 1963

If  $G$  is a graph on  $n$  vertices with  $n \geq 3k$  and  $\delta(G) \geq 2k$ , then  $G$  contains  $k$  disjoint cycles.

Examples:

- $k = 1$

# Corrádi-Hajnal Theorem

Corrádi-Hajnal, 1963

If  $G$  is a graph on  $n$  vertices with  $n \geq 3k$  and  $\delta(G) \geq 2k$ , then  $G$  contains  $k$  disjoint cycles.

Examples:

- $k = 1$ : easy


# Corrádi-Hajnal Theorem

Corrádi-Hajnal, 1963

If  $G$  is a graph on  $n$  vertices with  $n \geq 3k$  and  $\delta(G) \geq 2k$ , then  $G$  contains  $k$  disjoint cycles.

Examples:

- $k = 1$ : easy
- Sharpness:





# Corrádi-Hajnal Theorem

Corrádi-Hajnal, 1963

If  $G$  is a graph on  $n$  vertices with  $n \geq 3k$  and  $\delta(G) \geq 2k$ , then  $G$  contains  $k$  disjoint cycles.

Examples:

- $k = 1$ : easy
- Sharpness:



Corrádi-Hajnal, 1963

If  $G$  is a graph on  $n$  vertices with  $n \geq 3k$  and  $\delta(G) \geq 2k$ , then  $G$  contains  $k$  disjoint cycles.

## Corrádi-Hajnal, 1963

If  $G$  is a graph on  $n$  vertices with  $n \geq 3k$  and  $\delta(G) \geq 2k$ , then  $G$  contains  $k$  disjoint cycles.

$$\sigma_2(G) := \min\{d(x) + d(y) : xy \notin E(G)\}$$

## Corrádi-Hajnal, 1963

If  $G$  is a graph on  $n$  vertices with  $n \geq 3k$  and  $\delta(G) \geq 2k$ , then  $G$  contains  $k$  disjoint cycles.

$$\sigma_2(G) := \min\{d(x) + d(y) : xy \notin E(G)\}$$

## Enomoto 1998, Wang 1999

If  $G$  is a graph on  $n$  vertices with  $n \geq 3k$  and  $\sigma_2(G) \geq 4k - 1$ , then  $G$  contains  $k$  disjoint cycles.

## Corrádi-Hajnal, 1963

If  $G$  is a graph on  $n$  vertices with  $n \geq 3k$  and  $\delta(G) \geq 2k$ , then  $G$  contains  $k$  disjoint cycles.

$$\sigma_2(G) := \min\{d(x) + d(y) : xy \notin E(G)\}$$

## Enomoto 1998, Wang 1999

If  $G$  is a graph on  $n$  vertices with  $n \geq 3k$  and  $\sigma_2(G) \geq 4k - 1$ , then  $G$  contains  $k$  disjoint cycles.

Implies Corrádi-Hajnal


Enomoto 1998, Wang 1999

If  $G$  is a graph on  $n$  vertices with  $n \geq 3k$  and  $\sigma_2(G) \geq 4k - 1$ , then  $G$  contains  $k$  disjoint cycles.

Enomoto 1998, Wang 1999

If  $G$  is a graph on  $n$  vertices with  $n \geq 3k$  and  $\sigma_2(G) \geq 4k - 1$ , then  $G$  contains  $k$  disjoint cycles.

Sharpness:



Independence Number:

Independence Number:


Observation:

$\alpha(G) \geq n - 2k + 1 \Rightarrow G$  has no  $k$  cycles

Independence Number:

Observation:



$\alpha(G) \geq n - 2k + 1 \Rightarrow G$  has no  $k$  cycles



Independence Number:

Observation:

$\alpha(G) \geq n - 2k + 1 \Rightarrow G$  has no  $k$  cycles



Independence Number:

Observation:

$\alpha(G) \geq n - 2k + 1 \Rightarrow G$  has no  $k$  cycles

Enomoto 1998, Wang 1999

If  $G$  is a graph on  $n$  vertices with  $n \geq 3k$  and  $\sigma_2(G) \geq 4k - 1$ , then  $G$  contains  $k$  disjoint cycles.

# Kierstead-Kostochka-Y, 2014+

Independence Number:

Observation:

$\alpha(G) \geq n - 2k + 1 \Rightarrow G$  has no  $k$  cycles

Enomoto 1998, Wang 1999

If  $G$  is a graph on  $n$  vertices with  $n \geq 3k$  and  $\sigma_2(G) \geq 4k - 1$ , then  $G$  contains  $k$  disjoint cycles.

KKY, 2014+

For  $k \geq 4$ , if  $G$  is a graph on  $n$  vertices with  $n \geq 3k + 1$  and  $\sigma_2(G) \geq 4k - 3$ , then  $G$  contains  $k$  disjoint cycles if and only if  $\alpha(G) \leq n - 2k$ .



KKY, 2014+

For  $k \geq 4$ , if  $G$  is a graph on  $n$  vertices with  $n \geq 3k + 1$  and  $\sigma_2(G) \geq 4k - 3$ , then  $G$  contains  $k$  disjoint cycles if and only if  $\alpha(G) \leq n - 2k$ .

KKY, 2014+

For  $k \geq 4$ , if  $G$  is a graph on  $n$  vertices with  $n \geq 3k + 1$  and  $\sigma_2(G) \geq 4k - 3$ , then  $G$  contains  $k$  disjoint cycles if and only if  $\alpha(G) \leq n - 2k$ .


$$n \geq 3k + 1$$



KKY, 2014+

For  $k \geq 4$ , if  $G$  is a graph on  $n$  vertices with  $n \geq 3k + 1$  and  $\sigma_2(G) \geq 4k - 3$ , then  $G$  contains  $k$  disjoint cycles if and only if  $\alpha(G) \leq n - 2k$ .


$k = 1$ :



## KKY, 2014+

For  $k \geq 4$ , if  $G$  is a graph on  $n$  vertices with  $n \geq 3k + 1$  and  $\sigma_2(G) \geq 4k - 3$ , then  $G$  contains  $k$  disjoint cycles if and only if  $\alpha(G) \leq n - 2k$ .



$k = 2$ :



KKY, 2014+

For  $k \geq 4$ , if  $G$  is a graph on  $n$  vertices with  $n \geq 3k + 1$  and  $\sigma_2(G) \geq 4k - 3$ , then  $G$  contains  $k$  disjoint cycles if and only if  $\alpha(G) \leq n - 2k$ .



$k = 3$ :



KKY, 2014+

For  $k \geq 4$ , if  $G$  is a graph on  $n$  vertices with  $n \geq 3k + 1$  and  $\sigma_2(G) \geq 4k - 3$ , then  $G$  contains  $k$  disjoint cycles if and only if  $\alpha(G) \leq n - 2k$ .

$$\sigma_2 = 4k - 4:$$



Dirac:  $(2k - 1)$ -connected without  $k$  disjoint cycles

Dirac, 1963

What  $(2k - 1)$ -connected graphs do not have  $k$  disjoint cycles?

# Dirac: $(2k - 1)$ -connected without $k$ disjoint cycles

Dirac, 1963

What  $(2k - 1)$ -connected graphs do not have  $k$  disjoint cycles?

Observation:

$G$  is  $(2k - 1)$  connected  $\Rightarrow$

# Dirac: $(2k - 1)$ -connected without $k$ disjoint cycles

Dirac, 1963

What  $(2k - 1)$ -connected graphs do not have  $k$  disjoint cycles?

Observation:

$G$  is  $(2k - 1)$  connected  $\Rightarrow \delta(G) \geq 2k - 1 \Rightarrow$

# Dirac: $(2k - 1)$ -connected without $k$ disjoint cycles

Dirac, 1963

What  $(2k - 1)$ -connected graphs do not have  $k$  disjoint cycles?

Observation:

$G$  is  $(2k - 1)$  connected  $\Rightarrow \delta(G) \geq 2k - 1 \Rightarrow \sigma_2(G) \geq 4k - 2$

# Dirac: $(2k - 1)$ -connected without $k$ disjoint cycles

Dirac, 1963

What  $(2k - 1)$ -connected graphs do not have  $k$  disjoint cycles?

Observation:

$G$  is  $(2k - 1)$  connected  $\Rightarrow \delta(G) \geq 2k - 1 \Rightarrow \sigma_2(G) \geq 4k - 2$

KKY, 2014+

For  $k \geq 4$ , if  $G$  is a graph on  $n$  vertices with  $n \geq 3k + 1$  and  $\sigma_2(G) \geq 4k - 3$ , then  $G$  contains  $k$  disjoint cycles if and only if  $\alpha(G) \leq n - 2k$ .

## Dirac: $(2k - 1)$ -connected without $k$ disjoint cycles

KKY, 2014+

For  $k \geq 4$ , if  $G$  is a graph on  $n$  vertices with  $n \geq 3k + 1$  and  $\sigma_2(G) \geq 4k - 3$ , then  $G$  contains  $k$  disjoint cycles if and only if  $\alpha(G) \leq n - 2k$ .

### Answer to Dirac's Question for Simple Graphs

Let  $k \geq 2$ . Every graph  $G$  with (i)  $|G| \geq 3k$  and (ii)  $\delta(G) \geq 2k - 1$  contains  $k$  disjoint cycles if and only if

- $\alpha(G) \leq |G| - 2k$ , and
- if  $k$  is odd and  $|G| = 3k$ , then  $G \neq 2K_k \vee \overline{K_k}$ , and
- if  $k = 2$  then  $G$  is not a wheel.

## Dirac: $(2k - 1)$ -connected without $k$ disjoint cycles

KKY, 2014+

For  $k \geq 4$ , if  $G$  is a graph on  $n$  vertices with  $n \geq 3k + 1$  and  $\sigma_2(G) \geq 4k - 3$ , then  $G$  contains  $k$  disjoint cycles if and only if  $\alpha(G) \leq n - 2k$ .

### Answer to Dirac's Question for Simple Graphs

Let  $k \geq 2$ . Every graph  $G$  with (i)  $|G| \geq 3k$  and (ii)  $\delta(G) \geq 2k - 1$  contains  $k$  disjoint cycles if and only if

- $\alpha(G) \leq |G| - 2k$ , and
- if  $k$  is odd and  $|G| = 3k$ , then  $G \neq 2K_k \vee \overline{K_k}$ , and
- if  $k = 2$  then  $G$  is not a wheel.

Further:

## Dirac: $(2k - 1)$ -connected without $k$ disjoint cycles

KKY, 2014+

For  $k \geq 4$ , if  $G$  is a graph on  $n$  vertices with  $n \geq 3k + 1$  and  $\sigma_2(G) \geq 4k - 3$ , then  $G$  contains  $k$  disjoint cycles if and only if  $\alpha(G) \leq n - 2k$ .

### Answer to Dirac's Question for Simple Graphs

Let  $k \geq 2$ . Every graph  $G$  with (i)  $|G| \geq 3k$  and (ii)  $\delta(G) \geq 2k - 1$  contains  $k$  disjoint cycles if and only if

- $\alpha(G) \leq |G| - 2k$ , and
- if  $k$  is odd and  $|G| = 3k$ , then  $G \neq 2K_k \vee \overline{K_k}$ , and
- if  $k = 2$  then  $G$  is not a wheel.

Further:

characterization for multigraphs

# $(2k - 1)$ -connected multigraphs with no $k$ disjoint cycles

## Answer to Dirac's Question for multigraphs:

Let  $k \geq 2$  and  $n \geq k$ . Let  $G$  be an  $n$ -vertex graph with simple degree at least  $2k - 1$  and no loops. Let  $F$  be the simple graph induced by the strong edges of  $G$ ,  $\alpha' = \alpha'(F)$ , and  $k' = k - \alpha'$ . Then  $G$  does not contain  $k$  disjoint cycles if and only if one of the following holds:

- $n + \alpha' < 3k$ ;
- $|F| = 2\alpha'$  (i.e.,  $F$  has a perfect matching) and either (i)  $k'$  is odd and  $G - F = Y_{k',k'}$ , or (ii)  $k' = 2 < k$  and  $G - F$  is a wheel with 5 spokes;
- $G$  is extremal and either (i) some big set is not incident to any strong edge, or (ii) for some two distinct big sets  $I_j$  and  $I_{j'}$ , all strong edges intersecting  $I_j \cup I_{j'}$  have a common vertex outside of  $I_j \cup I_{j'}$ ;
- $n = 2\alpha' + 3k'$ ,  $k'$  is odd, and  $F$  has a superstar  $S = \{v_0, \dots, v_s\}$  with center  $v_0$  such that either (i)  $G - (F - S + v_0) = Y_{k'+1,k'}$ , or (ii)  $s = 2$ ,  $v_1 v_2 \in E(G)$ ,  $G - F = Y_{k'-1,k'}$  and  $G$  has no edges between  $\{v_1, v_2\}$  and the set  $X_0$  in  $G - F$ ;
- $k = 2$  and  $G$  is a wheel, where some spokes could be strong edges;
- $k' = 2$ ,  $|F| = 2\alpha' + 1 = n - 5$ , and  $G - F = C_5$ .

# Multigraphs

## Theorem (Extension of Corrádi-Hajnal to Multigraphs)

For  $k \in \mathbb{Z}^+$ , let  $G$  be a multigraph with simple degree at least  $2k$ .  
Then  $G$  has  $k$  disjoint cycles if and only if


$$|V(G)| \geq 3k - 2\ell - \alpha' \tag{1}$$

# Multigraphs

## Theorem (Extension of Corrádi-Hajnal to Multigraphs)

For  $k \in \mathbb{Z}^+$ , let  $G$  be a multigraph with simple degree at least  $2k$ .  
Then  $G$  has  $k$  disjoint cycles if and only if

$$|V(G)| \geq 3k - 2\ell - \alpha' \quad (1)$$




# Multigraphs

## Theorem (Extension of Corrádi-Hajnal to Multigraphs)

For  $k \in \mathbb{Z}^+$ , let  $G$  be a multigraph with simple degree at least  $2k$ .  
Then  $G$  has  $k$  disjoint cycles if and only if

$$|V(G)| \geq 3k - 2\ell - \alpha' \quad (1)$$




# Multigraphs

## Theorem (Extension of Corrádi-Hajnal to Multigraphs)

For  $k \in \mathbb{Z}^+$ , let  $G$  be a multigraph with simple degree at least  $2k$ .  
Then  $G$  has  $k$  disjoint cycles if and only if

$$|V(G)| \geq 3k - 2\ell - \alpha' \quad (1)$$




# Multigraphs

## Theorem (Extension of Corrádi-Hajnal to Multigraphs)

For  $k \in \mathbb{Z}^+$ , let  $G$  be a multigraph with simple degree at least  $2k$ .  
Then  $G$  has  $k$  disjoint cycles if and only if

$$|V(G)| \geq 3k - 2\ell - \alpha' \quad (1)$$




# Multigraphs

## Theorem (Extension of Corrádi-Hajnal to Multigraphs)

For  $k \in \mathbb{Z}^+$ , let  $G$  be a multigraph with simple degree at least  $2k$ .  
Then  $G$  has  $k$  disjoint cycles if and only if

$$|V(G)| \geq 3k - 2\ell - \alpha' \quad (1)$$




# Multigraphs

## Theorem (Extension of Corrádi-Hajnal to Multigraphs)

For  $k \in \mathbb{Z}^+$ , let  $G$  be a multigraph with simple degree at least  $2k$ .  
Then  $G$  has  $k$  disjoint cycles if and only if

$$|V(G)| \geq 3k - 2\ell - \alpha' \quad (1)$$




Remaining graph:  $\min \text{ degree} \geq 2k - \ell - 2\alpha'$

# Multigraphs

## Theorem (Extension of Corrádi-Hajnal to Multigraphs)

For  $k \in \mathbb{Z}^+$ , let  $G$  be a multigraph with simple degree at least  $2k$ .  
Then  $G$  has  $k$  disjoint cycles if and only if

$$|V(G)| \geq 3k - 2\ell - \alpha' \quad (1)$$




Remaining graph:  $\min \text{ degree} \geq 2k - \ell - 2\alpha' = 2(k - \ell - \alpha') + \ell$

# Multigraphs

## Theorem (Extension of Corrádi-Hajnal to Multigraphs)

For  $k \in \mathbb{Z}^+$ , let  $G$  be a multigraph with simple degree at least  $2k - 1$ . Then  $G$  has  $k$  disjoint cycles if and only if

$$|V(G)| \geq 3k - 2\ell - \alpha' \quad (1)$$




Remaining graph:  $\min \text{ degree} \geq 2k - \ell - 2\alpha' = 2(k - \ell - \alpha') + \ell$

# Multigraphs

## Theorem (Extension of Corrádi-Hajnal to Multigraphs)

For  $k \in \mathbb{Z}^+$ , let  $G$  be a multigraph with simple degree at least  $2k - 1$ . Then  $G$  has  $k$  disjoint cycles if and only if

$$|V(G)| \geq 3k - 2\ell - \alpha' \quad (1)$$



Remaining graph:  $\min \text{ degree} \geq 2k - 1 - \ell - 2\alpha' = 2(k - \ell - \alpha') + \ell - 1$

# Multigraphs

## Theorem (Extension of Corrádi-Hajnal to Multigraphs)

For  $k \in \mathbb{Z}^+$ , let  $G$  be a multigraph with simple degree at least  $2k - 1$ . Then  $G$  has  $k$  disjoint cycles if and only if

$$|V(G)| \geq 3k - 2\ell - \alpha' \quad (1)$$

Remaining graph:  $\min \text{ degree} \geq 2k - 1 - \ell - 2\alpha' = 2(k - \ell - \alpha') + \ell - 1$

## Corollary

Let  $G$  be a multigraph with simple degree at least  $2k - 1$  for some integer  $k \geq 2$ . Suppose  $G$  contains at least one loop. Then  $G$  has  $k$  disjoint cycles if and only if

$$|V(G)| \geq 3k - 2\ell - \alpha'.$$

# $(2k - 1)$ -connected multigraphs with no $k$ disjoint cycles

## Answer to Dirac's Question for multigraphs:

Let  $k \geq 2$  and  $n \geq k$ . Let  $G$  be an  $n$ -vertex graph with simple degree at least  $2k - 1$  and **no loops**. Let  $F$  be the simple graph induced by the strong edges of  $G$ ,  $\alpha' = \alpha'(F)$ , and  $k' = k - \alpha'$ . Then  $G$  does not contain  $k$  disjoint cycles if and only if one of the following holds:

- $n + \alpha' < 3k$ ;
- $|F| = 2\alpha'$  (i.e.,  $F$  has a perfect matching) and either (i)  $k'$  is odd and  $G - F = Y_{k',k'}$ , or (ii)  $k' = 2 < k$  and  $G - F$  is a wheel with 5 spokes;
- $G$  is extremal and either (i) some big set is not incident to any strong edge, or (ii) for some two distinct big sets  $I_j$  and  $I_{j'}$ , all strong edges intersecting  $I_j \cup I_{j'}$  have a common vertex outside of  $I_j \cup I_{j'}$ ;
- $n = 2\alpha' + 3k'$ ,  $k'$  is odd, and  $F$  has a superstar  $S = \{v_0, \dots, v_s\}$  with center  $v_0$  such that either (i)  $G - (F - S + v_0) = Y_{k'+1,k'}$ , or (ii)  $s = 2$ ,  $v_1 v_2 \in E(G)$ ,  $G - F = Y_{k'-1,k'}$  and  $G$  has no edges between  $\{v_1, v_2\}$  and the set  $X_0$  in  $G - F$ ;
- $k = 2$  and  $G$  is a wheel, where some spokes could be strong edges;
- $k' = 2$ ,  $|F| = 2\alpha' + 1 = n - 5$ , and  $G - F = C_5$ .

# $(2k - 1)$ -connected multigraphs with no $k$ disjoint cycles


## Answer to Dirac's Question for multigraphs:

Let  $k \geq 2$  and  $n \geq k$ . Let  $G$  be an  $n$ -vertex graph with simple degree at least  $2k - 1$  and no loops. Let  $F$  be the simple graph induced by the strong edges of  $G$ ,  $\alpha' = \alpha'(F)$ , and  $k' = k - \alpha'$ . Then  $G$  does not contain  $k$  disjoint cycles if and only if one of the following holds:

- $n + \alpha' < 3k$ ;
- $|F| = 2\alpha'$  (i.e.,  $F$  has a perfect matching) and either (i)  $k'$  is odd and  $G - F = Y_{k', k'}$ , or (ii)  $k' = 2 < k$  and  $G - F$  is a wheel with 5 spokes;
- $G$  is extremal and either (i) some big set is not incident to any strong edge, or (ii) for some two distinct big sets  $I_j$  and  $I_{j'}$ , all strong edges intersecting  $I_j \cup I_{j'}$  have a common vertex outside of  $I_j \cup I_{j'}$ ;
- $n = 2\alpha' + 3k'$ ,  $k'$  is odd, and  $F$  has a superstar  $S = \{v_0, \dots, v_s\}$  with center  $v_0$  such that either (i)  $G - (F - S + v_0) = Y_{k'+1, k'}$ , or (ii)  $s = 2$ ,  $v_1 v_2 \in E(G)$ ,  $G - F = Y_{k'-1, k'}$  and  $G$  has no edges between  $\{v_1, v_2\}$  and the set  $X_0$  in  $G - F$ ;
- $k = 2$  and  $G$  is a wheel, where some spokes could be strong edges;
- $k' = 2$ ,  $|F| = 2\alpha' + 1 = n - 5$ , and  $G - F = C_5$ .

$k'$  odd,  $F$  has a perfect matching

Example:  $k = 8$ ,  $\alpha' = 3$ ,  $k' = 5$ .




# $(2k - 1)$ -connected multigraphs with no $k$ disjoint cycles

## Answer to Dirac's Question for multigraphs:

Let  $k \geq 2$  and  $n \geq k$ . Let  $G$  be an  $n$ -vertex graph with simple degree at least  $2k - 1$  and no loops. Let  $F$  be the simple graph induced by the strong edges of  $G$ ,  $\alpha' = \alpha'(F)$ , and  $k' = k - \alpha'$ . Then  $G$  does not contain  $k$  disjoint cycles if and only if one of the following holds:

- $n + \alpha' < 3k$ ;
- $|F| = 2\alpha'$  (i.e.,  $F$  has a perfect matching) and either (i)  $k'$  is odd and  $G - F = Y_{k',k'}$ , or (ii)  $k' = 2 < k$  and  $G - F$  is a wheel with 5 spokes;
- $G$  is extremal and either (i) **some big set is not incident to any strong edge**, or (ii) for some two distinct big sets  $I_j$  and  $I_{j'}$ , all strong edges intersecting  $I_j \cup I_{j'}$  have a common vertex outside of  $I_j \cup I_{j'}$ ;
- $n = 2\alpha' + 3k'$ ,  $k'$  is odd, and  $F$  has a superstar  $S = \{v_0, \dots, v_s\}$  with center  $v_0$  such that either (i)  $G - (F - S + v_0) = Y_{k'+1,k'}$ , or (ii)  $s = 2$ ,  $v_1 v_2 \in E(G)$ ,  $G - F = Y_{k'-1,k'}$  and  $G$  has no edges between  $\{v_1, v_2\}$  and the set  $X_0$  in  $G - F$ ;
- $k = 2$  and  $G$  is a wheel, where some spokes could be strong edges;
- $k' = 2$ ,  $|F| = 2\alpha' + 1 = n - 5$ , and  $G - F = C_5$ .

# Big independent set, incident to no multiple edges



# Equitable Coloring

# Equitable Coloring


## Definition

An *equitable  $k$ -coloring* of a graph  $G$  is a proper coloring of  $V(G)$  such that any two color classes differ in size by at most one.

# Equitable Coloring

## Definition


An *equitable  $k$ -coloring* of a graph  $G$  is a proper coloring of  $V(G)$  such that any two color classes differ in size by at most one.



# Equitable Coloring

## Definition

An *equitable  $k$ -coloring* of a graph  $G$  is a proper coloring of  $V(G)$  such that any two color classes differ in size by at most one.



# Equitable Coloring and Cycles


$$n = 3k$$

If  $G$  has  $n = 3k$  vertices, then  $G$  has an equitable  $k$ -coloring if and only if  $\overline{G}$  has  $k$  disjoint cycles (all triangles).

# Equitable Coloring and Cycles

$$n = 3k$$




If  $G$  has  $n = 3k$  vertices, then  $G$  has an equitable  $k$ -coloring if and only if  $\overline{G}$  has  $k$  disjoint cycles (all triangles).



# Equitable Coloring and Cycles

$$n = 3k$$

If  $G$  has  $n = 3k$  vertices, then  $G$  has an equitable  $k$ -coloring if and only if  $\overline{G}$  has  $k$  disjoint cycles (all triangles).



Hajnal-Szemerédi, 1970

If  $k \geq \Delta(G) + 1$ , then  $G$  is equitably  $k$ -colorable.

Hajnal-Szemerédi, 1970

If  $k \geq \Delta(G) + 1$ , then  $G$  is equitably  $k$ -colorable.

Chen-Lih-Wu **Conjecture**

If  $\chi(G), \Delta(G) \leq k$ , and if  $k$  is odd  $K_{k,k} \not\subseteq G$ , then  $G$  is equitably  $k$ -colorable.

## Chen-Lih-Wu Conjecture

If  $\chi(G), \Delta(G) \leq k$  and  $K_{k,k} \not\subseteq G$ , then  $G$  is equitably  $k$ -colorable.

# Ore Conditions

## Chen-Lih-Wu Conjecture

If  $\chi(G), \Delta(G) \leq k$  and  $K_{k,k} \not\subseteq G$ , then  $G$  is equitably  $k$ -colorable.

## Kierstead-Kostochka-Molla-Y, 2014+

If  $G$  is a  $k$ -colorable  $3k$ -vertex graph such that for each edge  $xy$ ,  $d(x) + d(y) \leq 2k + 1$ , then  $G$  is equitably  $k$ -colorable, or is one of several exceptions.

# Ore Conditions

## Chen-Lih-Wu Conjecture

If  $\chi(G), \Delta(G) \leq k$  and  $K_{k,k} \not\subseteq G$ , then  $G$  is equitably  $k$ -colorable.

## Kierstead-Kostochka-Molla-Y, 2014+

If  $G$  is a  $k$ -colorable  $3k$ -vertex graph such that for each edge  $xy$ ,  $d(x) + d(y) \leq 2k + 1$ , then  $G$  is equitably  $k$ -colorable, or is one of several exceptions.

## Equivalent

If  $G$  is a graph on  $3k$  vertices with  $\sigma_2(G) \geq 4k - 3$ , then  $G$  contains  $k$  disjoint cycles, or is one of several exceptions, or  $\overline{G}$  is not  $k$ -colorable.

# Ore Conditions

Kierstead-Kostochka-Molla-Y, 2014+

If  $G$  is a  $k$ -colorable  $3k$ -vertex graph such that for each edge  $xy$ ,  $d(x) + d(y) \leq 2k + 1$ , then  $G$  is equitably  $k$ -colorable, or is one of several exceptions.

Equivalent

If  $G$  is a graph on  $3k$  vertices with  $\sigma_2(G) \geq 4k - 3$ , then  $G$  contains  $k$  disjoint cycles, or is one of several exceptions, or  $\overline{G}$  is not  $k$ -colorable.


KKY, 2014+

For  $k \geq 4$ , if  $G$  is a graph on  $n$  vertices with  $n \geq 3k + 1$  and  $\sigma_2(G) \geq 4k - 3$ , then  $G$  contains  $k$  disjoint cycles if and only if  $\alpha(G) \leq n - 2k$ .


# Exceptions

- $k = 3$

*Equitable coloring:*




*Cycles:*




# Exceptions

- *Equitable coloring:*





*Cycles:*




# Exceptions

- *Equitable coloring:*



*Cycles:*



Thanks for Listening!