

Roots of Polynomials

Here are some tricks for finding roots of polynomials. These tricks work well on exams and homework assignments, where polynomials tend to have integer coefficients and roots that are integers, or at least fractions.

Trick # 1

If r or $-r$ is an integer root of a polynomial $a_n x^n + \cdots + a_1 x + a_0$ with integer coefficients, then r is a factor of the constant term a_0 .

To see that this is true, just observe that for any root $\pm r$

$$a_n(\pm r)^n + \cdots + a_1(\pm r) + a_0 = 0 \implies a_0 = -[a_n(\pm r)^n + \cdots + a_1(\pm r)]$$

Every term on the right hand side is an integer times a strictly positive power of r . So the right hand side, and hence the left hand side, is some integer times r .

Example. $P(x) = x^3 - x^2 + 2$.

The constant term in this polynomial is $2 = 1 \times 2$. So the only candidates for integer roots are ± 1 , ± 2 . Trying each in turn

$$P(1) = 2 \qquad P(-1) = 0 \qquad P(2) = 6 \qquad P(-2) = -10$$

so the only integer root is -1 .

Trick # 2

If b/d or $-b/d$ is a rational root in lowest terms (i.e. b and d are integers with no common factors) of a polynomial $a_n x^n + \cdots + a_1 x + a_0$ with integer coefficients, then the numerator b is a factor of the constant term a_0 and the denominator d is a factor of a_n .

For any root $\pm b/d$

$$a_n(\pm b/d)^n + \cdots + a_1(\pm b/d) + a_0 = 0$$

Multiply through by d^n

$$a_0 d^n = -[a_n(\pm b)^n + a_{n-1}d(\pm b)^{n-1} + \cdots + a_1 d^{n-1}(\pm b)]$$

Every term on the right hand side is an integer times a strictly positive power of b . So the right hand side is some integer times b . The left hand side is $d^n a_0$ and d does not contain

any factor that is a factor of b . So a_0 must be some integer times b . Similarly, every term on the right hand side of

$$a_n(\pm b)^n = -[a_{n-1}d(\pm b)^{n-1} \cdots + a_1d^{n-1}(\pm b) + a_0d^n]$$

is an integer times a strictly positive power of d . So the right hand side is some integer times d . The left hand side is $a_n(\pm b)^n$ and b does not contain any factor that is a factor of d . So a_n must be some integer times d .

Example. $P(x) = 2x^2 - x - 3$.

The constant term in this polynomial is $3 = 1 \times 3$ and the coefficient of the highest power of x is $2 = 1 \times 2$. So the only candidates for integer roots are ± 1 , ± 3 and the only candidates for fractional roots are $\pm \frac{1}{2}$, $\pm \frac{3}{2}$.

$$\begin{array}{lll} P(1) = -2 & P(-1) = 0 & P(\pm 3) = 18 \mp 3 - 3 \neq 0 \\ P(\pm \frac{1}{2}) = 2\frac{1}{4} \mp \frac{1}{2} - 3 \neq 0 & P(\frac{3}{2}) = 2\frac{9}{4} - \frac{3}{2} - 3 = 0 & P(-\frac{3}{2}) = 2\frac{9}{4} + \frac{3}{2} - 3 \neq 0 \end{array}$$

So the roots are -1 and $\frac{3}{2}$.

Trick # 3 – Long Division

Suppose that $P(x)$ is a polynomial of degree p and suppose that you know that r is a root of that polynomial. In other words, suppose you know that $P(r) = 0$. Then it is always possible to factor $(x - r)$ out of $P(x)$. More precisely, it is always possible to find a polynomial $Q(x)$ of degree $p - 1$ such that

$$P(x) = (x - r)Q(x)$$

In sufficiently simple cases, you can probably do this factoring by inspection. For example, $P(x) = x^2 - 4$ has $r = 2$ as a root because $P(2) = 2^2 - 4 = 0$. In this case, $P(x) = (x - 2)(x + 2)$ so that $Q(x) = (x + 2)$. As another example, $P(x) = x^2 - 2x - 3$ has $r = -1$ as a root because $P(-1) = (-1)^2 - 2(-1) - 3 = 1 + 2 - 3 = 0$. In this case, $P(x) = (x + 1)(x - 3)$ so that $Q(x) = (x - 3)$.

Once you have found a root r of a polynomial, even if you cannot factor $(x - r)$ out of the polynomial by inspection, you can find $Q(x)$ by dividing $P(x)$ by $x - r$, using the long division algorithm you learned in public school, but with 10 replaced by x .

Example. $P(x) = x^3 - x^2 + 2$.

Because $P(-1) = (-1)^3 - (-1)^2 + 2 = -1 - 1 + 2 = 0$, $r = -1$ is a root of this polynomial and $x + 1$ must be a factor of $x^3 - x^2 + 2$. So we divide $\frac{x^3 - x^2 + 2}{x + 1}$. The first

term, x^2 , in the quotient is chosen so that when you multiply it by the denominator, $x^2(x + 1) = x^3 + x^2$, the leading term, x^3 , matches the leading term in the numerator, $x^3 - x^2 + 2$, exactly.

$$x + 1 \overline{\begin{array}{r} x^2 \\ x^3 - x^2 + 2 \\ x^3 + x^2 \end{array}}$$

When you subtract $x^2(x + 1) = x^3 + x^2$ from the numerator $x^3 - x^2 + 2$ you get the remainder $-2x^2 + 2$. Just like in public school, the 2 is not normally “brought down” until it is actually needed.

$$x + 1 \overline{\begin{array}{r} x^2 \\ x^3 - x^2 + 2 \\ x^3 + x^2 \\ -2x^2 \end{array}}$$

The next term, $-2x$, in the quotient is chosen so that when you multiply it by the denominator, $-2x(x + 1) = -2x^2 - 2x$, the leading term $-2x^2$ matches the leading term in the remainder exactly.

$$x + 1 \overline{\begin{array}{r} x^2 - 2x \\ x^3 - x^2 + 2 \\ x^3 + x^2 \\ -2x^2 \\ -2x^2 - 2x \end{array}}$$

And so on.

$$x + 1 \overline{\begin{array}{r} x^2 - 2x + 2 \\ x^3 - x^2 + 2 \\ x^3 + x^2 \\ -2x^2 \\ -2x^2 - 2x \\ 2x + 2 \\ 2x + 2 \\ 0 \end{array}}$$

Note that we finally end up with a remainder 0. A nonzero remainder would have signalled a computational error, since we know that the denominator $x - (-1)$ must divide the numerator $x^3 - x^2 + 2$ exactly. We conclude that

$$(x + 1)(x^2 - 2x + 2) = x^3 - x^2 + 2$$

To check this, just multiply out the left hand side explicitly.

There is an alternative to long division that involves more writing. In the previous example, we know that $\frac{x^3-x^2+2}{x+1}$ must be a polynomial (since -1 is a root of the numerator) of degree 2. So

$$\frac{x^3 - x^2 + 2}{x + 1} = ax^2 + bx + c$$

for some, as yet unknown, coefficients a , b and c . Cross multiplying and simplifying

$$\begin{aligned}x^3 - x^2 + 2 &= (ax^2 + bx + c)(x + 1) \\ &= ax^3 + (a + b)x^2 + (b + c)x + c\end{aligned}$$

Matching coefficients of the various powers of x on the left and right hand sides

$$\begin{aligned}\text{coefficient of } x^3: & \quad a = 1 \\ \text{coefficient of } x^2: & \quad a + b = -1 \\ \text{coefficient of } x^1: & \quad b + c = 0 \\ \text{coefficient of } x^0: & \quad c = 2\end{aligned}$$

tells us directly that $a = 1$ and $c = 2$. Subbing $a = 1$ into $a + b = -1$ tells us that $1 + b = -1$ and hence $b = -2$.