

Equality of Mixed Partial

Theorem. *If the partial derivatives $\frac{\partial^2 f}{\partial x \partial y}$ and $\frac{\partial^2 f}{\partial y \partial x}$ exist and are continuous at (x_0, y_0) , then*

$$\frac{\partial^2 f}{\partial x \partial y}(x_0, y_0) = \frac{\partial^2 f}{\partial y \partial x}(x_0, y_0)$$

Proof: Here is an outline of the proof. The details are given as footnotes at the end of the outline. Fix x_0 and y_0 and define⁽¹⁾

$$F(h, k) = \frac{1}{hk} [f(x_0 + h, y_0 + k) - f(x_0, y_0 + k) - f(x_0 + h, y_0) + f(x_0, y_0)]$$

Then, by the mean value theorem,

$$\begin{aligned} F(h, k) &\stackrel{2}{=} \frac{1}{h} \left[\frac{\partial f}{\partial y}(x_0 + h, y_0 + \theta_1 k) - \frac{\partial f}{\partial y}(x_0, y_0 + \theta_1 k) \right] \\ &\stackrel{3}{=} \frac{\partial}{\partial x} \frac{\partial f}{\partial y}(x_0 + \theta_2 h, y_0 + \theta_1 k) \\ F(h, k) &\stackrel{4}{=} \frac{1}{k} \left[\frac{\partial f}{\partial x}(x_0 + \theta_3 h, y_0 + k) - \frac{\partial f}{\partial x}(x_0 + \theta_3 h, y_0) \right] \\ &\stackrel{5}{=} \frac{\partial}{\partial y} \frac{\partial f}{\partial x}(x_0 + \theta_3 h, y_0 + \theta_4 k) \end{aligned}$$

for some $0 < \theta_1, \theta_2, \theta_3, \theta_4 < 1$. All of $\theta_1, \theta_2, \theta_3, \theta_4$ depend on x_0, y_0, h, k . Hence

$$\frac{\partial}{\partial x} \frac{\partial f}{\partial y}(x_0 + \theta_2 h, y_0 + \theta_1 k) = \frac{\partial}{\partial y} \frac{\partial f}{\partial x}(x_0 + \theta_3 h, y_0 + \theta_4 k)$$

for all h and k . Taking the limit $(h, k) \rightarrow (0, 0)$ and using the assumed continuity of both partial derivatives at (x_0, y_0) gives

$$\frac{\partial}{\partial x} \frac{\partial f}{\partial y}(x_0, y_0) = \frac{\partial}{\partial y} \frac{\partial f}{\partial x}(x_0, y_0)$$

The Details

- (1) We define $F(h, k)$ in this way because both partial derivatives $\frac{\partial^2 f}{\partial x \partial y}(x_0, y_0)$ and $\frac{\partial^2 f}{\partial y \partial x}(x_0, y_0)$ are defined as limits of $F(h, k)$ as $h, k \rightarrow 0$. For example,


$$\begin{aligned} \frac{\partial}{\partial x} \frac{\partial f}{\partial y}(x_0, y_0) &= \lim_{k \rightarrow 0} \frac{1}{k} \left[\frac{\partial f}{\partial x}(x_0, y_0 + k) - \frac{\partial f}{\partial x}(x_0, y_0) \right] \\ &= \lim_{k \rightarrow 0} \frac{1}{k} \left[\lim_{h \rightarrow 0} \frac{f(x_0 + h, y_0 + k) - f(x_0, y_0 + k)}{h} - \lim_{h \rightarrow 0} \frac{f(x_0 + h, y_0) - f(x_0, y_0)}{h} \right] \\ &= \lim_{k \rightarrow 0} \lim_{h \rightarrow 0} \frac{f(x_0 + h, y_0 + k) - f(x_0, y_0 + k) - f(x_0 + h, y_0) + f(x_0, y_0)}{hk} \\ &= \lim_{k \rightarrow 0} \lim_{h \rightarrow 0} F(h, k) \end{aligned}$$

Similarly,

$$\begin{aligned} \frac{\partial}{\partial y} \frac{\partial f}{\partial x}(x_0, y_0) &= \lim_{h \rightarrow 0} \frac{1}{h} \left[\frac{\partial f}{\partial y}(x_0 + h, y_0) - \frac{\partial f}{\partial y}(x_0, y_0) \right] \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left[\lim_{k \rightarrow 0} \frac{f(x_0 + h, y_0 + k) - f(x_0 + h, y_0)}{k} - \lim_{k \rightarrow 0} \frac{f(x_0, y_0 + k) - f(x_0, y_0)}{k} \right] \\ &= \lim_{h \rightarrow 0} \lim_{k \rightarrow 0} \frac{f(x_0 + h, y_0 + k) - f(x_0 + h, y_0) - f(x_0, y_0 + k) + f(x_0, y_0)}{hk} \\ &= \lim_{h \rightarrow 0} \lim_{k \rightarrow 0} F(h, k) \end{aligned}$$

- (2) The mean value theorem says that, for any differentiable function $\varphi(x)$, the slope of the line joining the points $(x_0, \varphi(x_0))$ and $(x_0 + k, \varphi(x_0 + k))$ on the graph of φ is the same as the slope of the tangent to the graph at some point between x_0 and $x_0 + k$. This is, there is some $0 < \theta_1 < 1$ such that

$$\frac{\varphi(x_0+k) - \varphi(x_0)}{k} = \frac{d\varphi}{dx}(x_0 + \theta_1 k)$$



Applying this with x replaced by y and φ replaced by $G(y) = f(x_0 + h, y) - f(x_0, y)$ gives

$$\begin{aligned} \frac{G(y_0+k) - G(y_0)}{k} &= \frac{dG}{dy}(y_0 + \theta_1 k) \quad \text{for some } 0 < \theta_1 < 1 \\ &= \frac{\partial f}{\partial y}(x_0 + h, y_0 + \theta_1 k) - \frac{\partial f}{\partial y}(x_0, y_0 + \theta_1 k) \end{aligned}$$

Hence, for some $0 < \theta_1 < 1$,

$$F(h, k) = \frac{1}{h} \left[\frac{G(y_0+k) - G(y_0)}{k} \right] = \frac{1}{h} \left[\frac{\partial f}{\partial y}(x_0 + h, y_0 + \theta_1 k) - \frac{\partial f}{\partial y}(x_0, y_0 + \theta_1 k) \right]$$

- (3) Define $H(x) = \frac{\partial f}{\partial y}(x, y_0 + \theta_1 k)$. By the mean value theorem

$$\begin{aligned} F(h, k) &= \frac{1}{h} \left[H(x_0 + h) - H(x_0) \right] \\ &= \frac{dH}{dx}(x_0 + \theta_2 h) \quad \text{for some } 0 < \theta_2 < 1 \\ &= \frac{\partial}{\partial x} \frac{\partial f}{\partial y}(x_0 + \theta_2 h, y_0 + \theta_1 k) \end{aligned}$$

- (4) Define $A(x) = f(x, y_0 + k) - f(x, y_0)$. By the mean value theorem

$$\begin{aligned} F(h, k) &= \frac{1}{k} \left[\frac{A(x_0+h) - A(x_0)}{h} \right] \\ &= \frac{1}{k} \frac{dA}{dx}(x_0 + \theta_3 h) \quad \text{for some } 0 < \theta_3 < 1 \\ &= \frac{1}{k} \left[\frac{\partial f}{\partial x}(x_0 + \theta_3 h, y_0 + k) - \frac{\partial f}{\partial x}(x_0 + \theta_3 h, y_0) \right] \end{aligned}$$

- (5) Define $B(y) = \frac{\partial f}{\partial x}(x_0 + \theta_3 h, y)$. By the mean value theorem

$$\begin{aligned} F(h, k) &= \frac{1}{k} \left[B(y_0 + k) - B(y_0) \right] \\ &= \frac{dB}{dy}(y_0 + \theta_4 k) \quad \text{for some } 0 < \theta_4 < 1 \\ &= \frac{\partial}{\partial y} \frac{\partial f}{\partial x}(x_0 + \theta_3 h, y_0 + \theta_4 k) \end{aligned}$$

■