


# Linear Regression

Imagine an experiment in which you measure one quantity, call it  $y$ , as a function of a second quantity, say  $x$ . For example,  $y$  could be the current that flows through a resistor when a voltage  $x$  is applied to it. Suppose that you measure  $n$  data points  $(x_1, y_1), \dots, (x_n, y_n)$  and that you wish to find the straight line  $y = mx + b$  that fits the data best. If the data point



$(x_i, y_i)$  were to land exactly on the line  $y = mx + b$  we would have  $y_i = mx_i + b$ . If it doesn't land exactly on the line, the vertical distance between  $(x_i, y_i)$  and the line  $y = mx + b$  is  $|y_i - mx_i - b|$ . That is, the discrepancy between the measured value of  $y_i$  and the corresponding idealized value on the line is  $|y_i - mx_i - b|$ . One measure of the total discrepancy for all data points is  $\sum_{i=1}^n |y_i - mx_i - b|$ . A more convenient measure, which avoids the absolute value signs, is

$$D(m, b) = \sum_{i=1}^n (y_i - mx_i - b)^2$$

We will now find the values of  $m$  and  $b$  that give the minimum value of  $D(m, b)$ . The corresponding line  $y = mx + b$  is generally viewed as the line that fits the data best.

You learned in your first Calculus course that the value of  $m$  that gives the minimum value of a function of one variable  $f(m)$  obeys  $f'(m) = 0$ . The analogous statement for functions of two variables is the following. First pretend that  $b$  is just a constant and compute the derivative of  $D(m, b)$  with respect to  $m$ . This is called the partial derivative of  $D(m, b)$  with respect to  $m$  and denoted  $\frac{\partial D}{\partial m}(m, b)$ . Next pretend that  $m$  is just a constant and compute the derivative of  $D(m, b)$  with respect to  $b$ . This is called the partial derivative of  $D(m, b)$  with respect to  $b$  and denoted  $\frac{\partial D}{\partial b}(m, b)$ . If  $(m, b)$  gives the minimum value of  $D(m, b)$ , then

$$\frac{\partial D}{\partial m}(m, b) = \frac{\partial D}{\partial b}(m, b) = 0$$

For our specific  $D(m, b)$

$$\begin{aligned} \frac{\partial D}{\partial m}(m, b) &= \sum_{i=1}^n 2(y_i - mx_i - b)(-x_i) \\ \frac{\partial D}{\partial b}(m, b) &= \sum_{i=1}^n 2(y_i - mx_i - b)(-1) \end{aligned}$$

It is important to remember here that all of the  $x_i$ 's and  $y_i$ 's here are given numbers. The only unknowns are  $m$  and  $b$ . The two partials are of the forms

$$\begin{aligned}\frac{\partial D}{\partial m}(m, b) &= 2c_{xx}m + 2c_x b - 2c_{xy} \\ \frac{\partial D}{\partial b}(m, b) &= 2c_x m + 2nb - 2c_y\end{aligned}$$

where the various  $c$ 's are just given numbers whose values are

$$c_{xx} = \sum_{i=1}^n x_i^2 \quad c_x = \sum_{i=1}^n x_i \quad c_{xy} = \sum_{i=1}^n x_i y_i \quad c_y = \sum_{i=1}^n y_i$$

So the value of  $(m, b)$  that gives the minimum value of  $D(m, b)$  is determined by

$$c_{xx}m + c_x b = c_{xy} \quad (1)$$

$$c_x m + nb = c_y \quad (2)$$

This is a system of two linear equations in the two unknowns  $m$  and  $b$ , which is easy to solve:

$$n(1) - c_x(2) : \quad [nc_{xx} - c_x^2]m = nc_{xy} - c_x c_y \quad \implies \quad m = \frac{nc_{xy} - c_x c_y}{nc_{xx} - c_x^2}$$

$$c_x(1) - c_{xx}(2) : \quad [c_x^2 - nc_{xx}]b = c_x c_{xy} - c_{xx} c_y \quad \implies \quad b = \frac{c_{xx} c_y - c_x c_{xy}}{nc_{xx} - c_x^2}$$