

Lecture Notes on Measure Theory and Integration

7 — The Radon-Nikodym Theorem

Joel Feldman

University of British Columbia

November 28, 2019

7 The Radon-Nikodym Theorem

Let $(X, \mathcal{M}, \mu)$ be a measure space. One easy way to build another measure from μ is to take a measurable nonnegative function f and set

$$\nu(E) = \int_E f(x) d\mu(x) \quad \text{for all } E \in \mathcal{M}$$

In this case, one usually writes $d\nu = f d\mu$.

Now suppose that someone gives you two measures, μ and ν , on $\mathcal{M}$. How can you tell if $d\nu = f d\mu$ for some f ? There is one obvious necessary condition. If $d\nu = f d\mu$, for some function f (even if you don't know what it is), and if $E \in \mathcal{M}$ is a null set for μ (i.e. $\mu(E) = 0$), then

$$\nu(E) = \int_E f(x) d\mu(x) = \int \overbrace{f \chi_E}^{=0 \text{ a.e.}} d\mu = \int 0 d\mu = 0$$

This property is given a name.

Definition 7.1. Let ν be a signed measure on $(X, \mathcal{M})$ and μ be a (positive) measure on $(X, \mathcal{M})$. Then ν is said to be **absolutely continuous** with respect to μ , denoted $\nu \ll \mu$, if

$$E \in \mathcal{M}, \mu(E) = 0 \implies \nu(E) = 0$$

In this part of the course we prove the Radon-Nikodym theorem, which says that, for σ -finite measures, absolute continuity is also a sufficient condition for ν to be of the form $d\nu = f d\mu$. Here is some prep.

Proposition 7.2. Let ν be a finite signed measure (i.e. ν_+ and ν_- are both finite) and μ a (positive) measure on $(X, \mathcal{M})$. Then

$$\nu \ll \mu \iff \forall \varepsilon > 0 \exists \delta > 0 \text{ such that } E \in \mathcal{M}, \mu(E) < \delta \implies |\nu(E)| < \varepsilon$$

Proof.

Part 1: Reduction to the case that ν is positive:

Use $|\nu|$ to denote the measure $\nu_+ + \nu_-$. Let $X = P \cup N$ be a Hahn decomposition (of Theorem 6.5) for ν . So

$$\nu_+(E) = \nu(E \cap P) \quad \nu_-(E) = -\nu(E \cap N)$$

For the left hand side of the statement of the theorem

$$\begin{aligned}
\nu \ll \mu &\iff \left(\overbrace{\mu(E) = 0}^{\iff \mu(E \cap P) + \mu(E \cap N) = 0} \implies \nu(E) = 0 \right) \\
&\iff \left(\mu(E) = 0 \implies \nu_+(E) = \nu_-(E) = 0 \right) \\
&\iff \left(\mu(E) = 0 \implies |\nu|(E) = 0 \right) \\
&\iff |\nu| \ll \mu
\end{aligned}$$

Call the statement on the right hand side of the theorem S_ν . That is, S_ν is “ $\forall \varepsilon > 0 \exists \delta > 0$ such that $E \in \mathcal{M}, \mu(E) < \delta \implies |\nu(E)| < \varepsilon$ ”. We now show that S_ν is true if and only if $S_{|\nu|}$ is true.

$$\begin{aligned}
S_\nu &\implies \left(\forall \varepsilon > 0 \exists \delta > 0 \text{ such that } E \in \mathcal{M}, \mu(E) < \delta \implies |\nu(E)| < \frac{\varepsilon}{2} \right) \\
&\hspace{15em} \text{(just replaced } \varepsilon \text{ by } \frac{\varepsilon}{2} \text{)} \\
&\implies \left(\forall \varepsilon > 0 \exists \delta > 0 \text{ s.t. } E \in \mathcal{M}, \mu(E) < \delta \implies |\nu(E \cap N)|, |\nu(E \cap P)| < \frac{\varepsilon}{2} \right) \\
&\hspace{15em} \text{(since } \mu(E) < \delta \text{ implies } \mu(E \cap N), \mu(E \cap P) < \delta \text{)} \\
&\implies \left(\forall \varepsilon > 0 \exists \delta > 0 \text{ s.t. } E \in \mathcal{M}, \mu(E) < \delta \implies |\nu|(E) < \varepsilon \right) \\
&\implies S_{|\nu|}
\end{aligned}$$

and

$$\begin{aligned}
S_{|\nu|} &\implies \left(\forall \varepsilon > 0 \exists \delta > 0 \text{ such that } E \in \mathcal{M}, \mu(E) < \delta \implies |\nu|(E) < \varepsilon \right) \\
&\implies \left(\forall \varepsilon > 0 \exists \delta > 0 \text{ such that } E \in \mathcal{M}, \mu(E) < \delta \implies |\nu(E)| < \varepsilon \right) \\
&\hspace{15em} \text{(since } |\nu(E)| \leq |\nu|(E) \text{)} \\
&\implies S_\nu
\end{aligned}$$

Part 2: Proof of $\Leftarrow$ in the case that ν is positive:

We are assuming that S_ν is true and have to prove that $\nu \ll \mu$. Let $E \in \mathcal{M}$. Then

$$\begin{aligned}
\mu(E) = 0 &\xrightarrow{S_\nu} |\nu(E)| < \varepsilon \text{ for all } \varepsilon > 0 \\
&\implies \nu(E) = 0
\end{aligned}$$

so that $\nu \ll \mu$.

Part 3: Proof of $\implies$ in the case that ν is positive:

We are assuming that $\nu \ll \mu$ and have to prove that S_ν is true. Assume that S_ν is false. Then there is an $\varepsilon > 0$ and, for every $n \in \mathbb{N}$, there is an $E_n \in \mathcal{M}$ such that $\mu(E_n) < \frac{1}{2^n}$ and $\nu(E_n) \geq \varepsilon$. Set

$$F_k = \bigcup_{n=k}^{\infty} E_n \quad F = \bigcap_{k=1}^{\infty} F_k$$

Then

$$\mu(F_k) \leq \sum_{n=k}^{\infty} \frac{1}{2^n} = \frac{2}{2^k} \implies \mu(F) = \lim_{k \rightarrow \infty} \mu(F_k) = 0$$

but

$$\nu(F_k) \geq \nu(E_k) \geq \varepsilon \xrightarrow{\nu \text{ finite!}} \nu(F) = \lim_{k \rightarrow \infty} \nu(F_k) \geq \varepsilon$$

contradicting the assumption that $\nu \ll \mu$.

□

Corollary 7.3. Let $f \in L^1(\mu)$. Then

$$\forall \varepsilon > 0 \exists \delta > 0 \text{ such that } E \in \mathcal{M}, \mu(E) < \delta \implies \left| \int_E f(x) d\mu(x) \right| < \varepsilon$$

Proof. Apply Proposition 7.2 with $\nu(E) = \int_E f(x) d\mu(x)$.

□

THEOREM 7.4 (The Radon-Nikodym Theorem). Let ν be a σ -finite signed measure and μ be a σ -finite (positive) measure on $(X, \mathcal{M})$. There exist unique σ -finite signed measures λ, ρ on $(X, \mathcal{M})$ such that

$$\lambda \perp \mu \quad \rho \ll \mu \quad \nu = \lambda + \rho$$

Moreover, there is an extended μ -integrable function $f : X \rightarrow \mathbb{R}$ such that $d\rho = f d\mu$. Any two such functions are equal almost everywhere.

First, we have a preliminary

Lemma 7.5. Let μ, ν be finite (positive) measures on $(X, \mathcal{M})$. Then either $\nu \perp \mu$ or

$$\exists \varepsilon > 0, E \in \mathcal{M} \text{ such that } \mu(E) > 0 \text{ and } \nu \geq \varepsilon \mu \text{ on } E$$

That is, $\mu(E) > 0$ and E is a positive set for $\nu - \varepsilon \mu$.

Proof. Let, for each $n \in \mathbb{N}$, $X = P_n \cup N_n$ be a Hahn decomposition for $\nu - \frac{1}{n}\mu$. Let

$$P = \bigcup_{n=1}^{\infty} P_n \quad N = \bigcap_{n=1}^{\infty} N_n = P^c$$

Then

$$\begin{aligned} & \text{For every } n \in \mathbb{N}, N_n \text{ is a negative set for } \nu - \frac{1}{n}\mu \\ & \implies N \text{ is a negative set for } \nu - \frac{1}{n}\mu \text{ for all } n \in \mathbb{N} \\ & \implies \nu(N) - \frac{1}{n}\mu(N) \leq 0 \text{ for all } n \in \mathbb{N} \\ & \implies 0 \leq \nu(N) \leq \frac{1}{n}\mu(N) \text{ for all } n \in \mathbb{N} \\ & \implies \nu(N) = 0 \end{aligned}$$

If $\mu(P) = 0$, then $\nu \perp \mu$.

If $\mu(P) > 0$, then, by continuity from below, $\mu(P_n) > 0$ for some $n \in \mathbb{N}$, and then P_n is a positive set for $\nu - \frac{1}{n}\mu$. □

Proof of Theorem 7.4.

Part 1: A little motivation in the case that μ, ν are finite and positive:

We want to guess a function f so that, if we set $d\rho = f d\mu$, then $\lambda = \nu - \rho$ is mutually singular with respect to μ . So let's suppose that we already know that

$$\begin{array}{llll} \nu = \lambda + \rho & d\rho = f d\mu & \lambda \perp \mu & \\ X = L \cup M & L \cap M = \emptyset & \lambda(M) = 0 & \mu(L) = 0 \end{array}$$

for some (unknown) λ, ρ, L, M , and see what conditions (that do not mention λ, ρ, L, M — which we do not yet know) f has to obey.

◦ Since ν is positive and $\lambda(E) = \lambda(E \cap L) = \nu(E \cap L) \geq 0$,

$$\nu(E) = \lambda(E) + \int_E f d\mu \geq \int_E f d\mu \quad \text{for all } E \in \mathcal{M}$$

and we'll use

$$\mathcal{F} = \left\{ \varphi : X \rightarrow [0, \infty], \text{ measurable} \mid \int_E \varphi d\mu \leq \nu(E) \forall E \in \mathcal{M} \right\}$$

as the set of possible candidates for the function f .

- Think of $\nu(X)$, $\lambda(X) = \nu(L)$ and $\rho(X) = \nu(M)$ as being the “mass” of X , L and M , respectively. We will try to pick f to be the candidate $\varphi \in \mathcal{F}$ which puts as much mass, $\int_X \varphi d\mu$, as possible into $\rho(X)$. The remaining mass, which cannot be put into $\rho(X)$ because it is not in M , will go into $\lambda(X)$, i.e. into L .

Part 2: Proof of existence in the case that μ, ν are finite and positive:

The set $\mathcal{F}$ of candidates for the function f has the properties

- $\mathcal{F} \neq \emptyset$ since $\varphi = 0 \in \mathcal{F}$
- if $\varphi, \psi \in \mathcal{F}$, then $\zeta = \max\{\varphi, \psi\} \in \mathcal{F}$ since, writing $A = \{x \in X \mid \varphi(x) > \psi(x)\}$,

$$\begin{aligned} \int_E \zeta d\mu &= \int_{A \cap E} \varphi d\mu + \int_{E \setminus A} \psi d\mu \\ &\leq \nu(A \cap E) + \nu(E \setminus A) = \nu(E) \end{aligned}$$

Set

$$a = \sup \left\{ \int_X \varphi d\mu \mid \varphi \in \mathcal{F} \right\} \leq \nu(X) < \infty$$

and choose $\{\varphi_n\}_{n \in \mathbb{N}} \subset \mathcal{F}$ such that $\lim_{n \rightarrow \infty} \int_X \varphi_n d\mu = a$. Then set

$$g_n(x) = \max\{\varphi_1(x), \dots, \varphi_n(x)\} \quad f(x) = \sup_n \varphi_n(x)$$

Then

- $g_n \in \mathcal{F}$, for all $n \in \mathbb{N}$
- $g_n(x)$ increases to $f(x)$, as $n \rightarrow \infty$, for all $x \in X$ and
- $a \geq \int_X g_n d\mu \geq \int_X \varphi_n d\mu \xrightarrow{n \rightarrow \infty} a$

which forces

$$\lim_{n \rightarrow \infty} \int_X g_n d\mu = a$$

By the monotone convergence theorem

$$\int_E f d\mu = \lim_{n \rightarrow \infty} \int_E g_n d\mu \leq \nu(E)$$

So

$$f \in \mathcal{F} \quad \int_X f d\mu = a < \infty \quad 0 \leq f < \infty \text{ a.e.}$$

Now we set

$$d\rho = f d\mu \quad \lambda = \nu - \rho$$

Then

- $\nu = \lambda + \rho$
- $\rho \ll \mu$, ρ is positive
- λ is positive, since $f \in \mathcal{F}$ so that $\rho(E) = \int_E f d\mu \leq \nu(E)$ for all $E \in \mathcal{M}$
- $\lambda \perp \mu$

Proof: If not, then by the preliminary Lemma 7.5, there is an $\varepsilon > 0$ and an $E_0 \in \mathcal{M}$ with $\mu(E_0) > 0$ and $\lambda \geq \varepsilon\mu$ on E_0 . Set $d\rho' = \varepsilon\chi_{E_0} d\mu$. Then $\rho' \leq \lambda = \nu - \rho$, or $\rho + \rho' \leq \nu$. That is

$$(f + \varepsilon\chi_{E_0}) d\mu \leq d\nu \implies f + \varepsilon\chi_{E_0} \in \mathcal{F} \text{ and}$$

$$\int_X (f + \varepsilon\chi_{E_0}) d\mu = a + \varepsilon\mu(E_0) > a$$

which contradicts the definition of a .

Part 3: Proof of uniqueness in the case that μ, ν are finite and positive:

Suppose that

$$\begin{aligned} \nu &= \lambda + \rho & d\rho &= f d\mu & X &= E \cup F & E \cap F &= \emptyset & \mu(E) &= 0 & \lambda(F) &= 0 \\ \nu &= \lambda' + \rho' & d\rho' &= f' d\mu & X &= E' \cup F' & E' \cap F' &= \emptyset & \mu(E') &= 0 & \lambda'(F') &= 0 \end{aligned}$$

We want to show that $\lambda = \lambda'$, $\rho = \rho'$ and $f = f'$ a.e.. We have $\lambda - \lambda' = \rho' - \rho$ and

- $\lambda - \lambda' \perp \mu$ since $\mu(E \cup E') = 0$ and $(\lambda - \lambda')(G) = 0$ for all $G \subset \overbrace{(E \cup E')^c}^{F \cap F'}$.
- $\rho' - \rho = (f' - f) d\mu \ll \mu$

which implies that, for all $S \in \mathcal{M}$,

$$\begin{aligned} (\lambda - \lambda')(S) &= (\lambda - \lambda')(S \cap (E \cup E')) \\ &= (\rho' - \rho)(S \cap (E \cup E')) \\ &= 0 \quad \text{since } \mu(S \cap (E \cup E')) = 0 \\ \implies (\rho' - \rho)(S) &= 0 \end{aligned}$$

which implies that $\lambda = \lambda'$ and that $\rho = \rho'$. Finally, to show that $f' - f = 0$ a.e., set, for each $n \in \mathbb{N}$, $P_n = \{x \mid f'(x) \geq f(x) + 1/n\}$ and $N_n = \{x \mid f(x) \geq f'(x) + 1/n\}$.

Then

$$\begin{aligned}
(\rho' - \rho)(P_n) = 0 &\implies \int_{P_n} \overbrace{(f' - f)}^{\geq 1/n} d\mu = 0 \implies \mu(P_n) = 0 \\
(\rho' - \rho)(N_n) = 0 &\implies \int_{N_n} \overbrace{(f' - f)}^{\leq -1/n} d\mu = 0 \implies \mu(N_n) = 0
\end{aligned}$$

Part 4: Proof in the case that μ, ν are σ -finite and positive:

Write

$$\begin{aligned}
X &= \bigcup_{m=1}^{\infty} E_m \quad \text{with the } E_m \text{'s disjoint and each } \mu(E_m) < \infty \\
X &= \bigcup_{n=1}^{\infty} F_n \quad \text{with the } F_n \text{'s disjoint and each } \nu(F_n) < \infty
\end{aligned}$$

Then

$$X = \bigcup_{m,n=1}^{\infty} E_m \cap F_n$$

with the $E_m \cap F_n$'s all disjoint and each $\mu(E_m \cap F_n), \nu(E_m \cap F_n) < \infty$. Set, for each $m, n \in \mathbb{N}$,

$$\begin{aligned}
\mu_{m,n}(A) &= \mu(A \cap E_m \cap F_n) \\
\nu_{m,n}(A) &= \nu(A \cap E_m \cap F_n)
\end{aligned}$$

By the previous step, we have

$$d\nu_{m,n} = d\lambda_{m,n} + f_{m,n} d\mu_{m,n} \quad \text{with } \lambda_{m,n} \perp \mu_{m,n}$$

It now suffices to set

$$\lambda = \sum_{m,n=1}^{\infty} \lambda_{m,n} \quad f = \sum_{m,n=1}^{\infty} f_{m,n} \chi_{E_m \cap F_n}$$

Part 4: Proof in the general case:

Just apply the previous case separately to ν_+ and ν_- .

□

Remark 7.6.

Here is some notation and terminology.

- (a) The decomposition $\nu = \lambda + \rho$ with $\lambda \perp \mu$ and $\rho \ll \mu$ is called the Lebesgue decomposition of ν with respect to μ .
- (b) If $\nu \ll \mu$, then $d\nu = f d\mu$ for some f . This is called the Radon-Nikodym theorem and we write $f = \frac{d\nu}{d\mu}$.