

**Math 308, Section 101**  
**First Midterm Exam**  
October 8, 2004  
Duration: 50 minutes

Name: \_\_\_\_\_ Student Number: \_\_\_\_\_

**Do not open this test until instructed to do so!** This exam should have 8 pages, including this cover sheet. No textbooks, notes, calculators, or other aids are allowed. Turn off any cell phones, pagers, etc. that could make noise during the exam. You must remain in this room until you have finished the exam.

**All your solutions must be written clearly and understandably.** Use complete sentences and explain why your mathematical statements are relevant to the problem. You should always write enough to demonstrate that you're not just guessing the answer. Use the backs of the pages if necessary. You will find some of the questions quite easy; try to solve these first. Good luck!

**Be aware of these UBC rules governing examinations:**

- (i) Each candidate must be prepared to produce, upon request, a Library / AMS card for identification.
- (ii) Candidates are not permitted to ask questions of the invigilators, except in cases of supposed errors or ambiguities in examination questions.
- (iii) No candidate shall be permitted to enter the examination room after the expiration of one-half hour from the scheduled starting time, or to leave during the first half hour of the examination.
- (iv) Candidates suspected of any of the following, or similar, dishonest practices shall be immediately dismissed from the examination and shall be liable to disciplinary action.
  - Having at the place of writing any books, papers or memoranda, calculators, computers, audio or video cassette players or other memory aid devices, other than those authorized by the examiners.
  - Speaking or communicating with other candidates.
  - Purposely exposing written papers to the view of other candidates. The plea of accident or forgetfulness shall not be received.
- (v) Candidates must not destroy or mutilate any examination material; must hand in all examination papers; and must not take any examination material from the examination room without permission of the invigilator.

| Problem | Out of | Score |
|---------|--------|-------|
| 1       | 4      |       |
| 2       | 6      |       |
| 3       | 6      |       |
| 4       | 8      |       |

| Problem      | Out of | Score |
|--------------|--------|-------|
| 5            | 4      |       |
| 6            | 6      |       |
| 7            | 8      |       |
| 8            | 8      |       |
| <b>Total</b> | 50     |       |

*These facts are here for your reference. This is not a complete list; you may quote any facts we have covered, as long as you make it clear what fact you are using. You are not required to quote them by theorem number, lemma number, and so on (but you may if you wish to). Please do not remove this page from the test booklet.*

**Fact.** Given a ray  $AB$  and a positive number  $x$ , there is exactly one point  $C$  on the ray  $AB$  such that  $|AC| = x$ .

**Theorem 1.4.1.** Let  $P$  be a point not on the line  $\ell$ , and let  $Q$  be a point on  $\ell$  such that  $PQ$  is perpendicular to  $\ell$ . Let  $\ell_2$  be the line through  $P$  that is parallel to  $\ell$ . Then  $\ell_2$  is perpendicular to  $PQ$ .

**Corollary 1.4.4.** Suppose a line  $\ell$  intersects two distinct lines  $\ell_1$  and  $\ell_2$  so that the opposite interior angles are equal. Then  $\ell_1$  and  $\ell_2$  are parallel. [Note 1: In particular, if  $\ell$  is perpendicular to both  $\ell_1$  and  $\ell_2$ , then  $\ell_1$  and  $\ell_2$  are parallel.] [Note 2: The converse is also true: if  $\ell_1$  and  $\ell_2$  are parallel, then the opposite interior angles formed by  $\ell$  are equal.]

**Fact.** Suppose that  $\ell_1$  and  $\ell_2$  are both perpendicular to some other line  $\ell$ . If  $\ell_1$  and  $\ell_2$  have a point in common, then  $\ell_1 = \ell_2$ .

**Pons Asinorum.** The base angles of an isosceles triangle are equal: if  $|AB| = |AC|$ , then  $\angle ABC = \angle ACB$ . [Note: the converse is also true: if  $\angle ABC = \angle ACB$ , then  $|AB| = |AC|$ .]

**Star Trek Lemma.** The measure of an inscribed angle equals half of the angular measure of the arc it subtends. [Note: the angular measure of an arc  $PQ$  on a circle centered at  $O$  is equal to the measure of the central angle  $\angle POQ$ .]

**Bow Tie Lemma.** Let  $A, A', B$ , and  $C$  lie on a circle, and suppose that  $\angle BAC$  and  $\angle BA'C$  subtend the same arc. Then  $\angle BAC = \angle BA'C$ .

**Theorem 1.7.1.** Let  $B'$  and  $C'$  be on the sides  $AB$  and  $AC$ , respectively, of  $\triangle ABC$ . Then  $B'C'$  is parallel to  $BC$  if and only if  $|AB'|/|AB| = |AC'|/|AC|$ .

**Corollary 1.7.4.** If  $\triangle ABC \sim \triangle A'B'C'$ , then  $|A'B'|/|AB| = |A'C'|/|AC| = |B'C'|/|BC|$ . [Note: the converse is also true: if  $|A'B'|/|AB| = |A'C'|/|AC| = |B'C'|/|BC|$ , then  $\triangle ABC \sim \triangle A'B'C'$ .]

**SAS for Similarity.** Let  $\triangle ABC$  and  $\triangle A'B'C'$  be two triangles such that  $\angle BAC = \angle B'A'C'$  and  $|A'B'|/|AB| = |A'C'|/|AC|$ . Then  $\triangle ABC \sim \triangle A'B'C'$ .

**Power of the Point Theorem.** Let  $P$  be any point and  $\mathcal{C}$  any circle, and let  $\ell_1$  and  $\ell_2$  be two lines through  $P$ . Suppose that  $\ell_1$  intersects  $\mathcal{C}$  in the two points  $Q$  and  $Q'$ , and  $\ell_2$  intersects  $\mathcal{C}$  in the two points  $R$  and  $R'$ . Then  $|PQ||PQ'| = |PR||PR'|$ .

**Theorem 1.9.1.** Let  $AA'$ ,  $BB'$ , and  $CC'$  be the three medians of the triangle  $\triangle ABC$ . Let  $G$  be the centroid of  $\triangle ABC$ . Then  $|AG| = 2|A'G|$ ,  $|BG| = 2|B'G|$ , and  $|CG| = 2|C'G|$ .

**Heron's Formula.** Let  $s$  be the semiperimeter of  $\triangle ABC$ . Then the area of  $\triangle ABC$  is given by the formula  $|\triangle ABC| = \sqrt{s(s-a)(s-b)(s-c)}$ . [Note:  $a$ ,  $b$ , and  $c$  are the lengths of the sides of  $\triangle ABC$ , and  $s = \frac{1}{2}(a+b+c)$ .]

**Law of Cosines.** For any triangle  $\triangle ABC$ , we have  $c^2 = a^2 + b^2 - 2ab \cos C$ .

**Extended Law of Sines.** Let  $R$  be the circumradius of  $\triangle ABC$ . Then  $a/\sin A = b/\sin B = c/\sin C = 2R$ . [Note: the circumradius of a triangle is the radius of the circle that can be circumscribed around it.]

**Theorem 1.10.2.** Let  $r$  be the inradius of  $\triangle ABC$ , and let  $s$  be the semiperimeter of  $\triangle ABC$ . Then the area of  $\triangle ABC$  is given by the formula  $|\triangle ABC| = rs$ . [Note: the inradius of a triangle is the radius of the circle that can be inscribed inside the triangle.]