

Math 534. Written problems, set 1.

- (1) Let \mathfrak{g} be a Lie algebra.
 - (a) Show that if \mathfrak{h} is an ideal in \mathfrak{g} , then $[\mathfrak{h}, \mathfrak{h}]$ is also an ideal in \mathfrak{g} .
 - (b) Let $\mathcal{D}^k \mathfrak{g}$ be the k -th term of the derived series of \mathfrak{g} . Show that it is, in fact, an ideal in \mathfrak{g} .
 - (c) Show that \mathfrak{g} is semisimple if and only if it has no non-trivial *abelian* ideals.
- (2) Humphreys, Exercise 3 on p.5.
- (3) Humphreys, Exercise 6 on p.5.
- (4) **Heisenberg Lie algebra.** Let V be a 3-dimensional vector space over a field F with basis $\{x, y, z\}$. Define the Lie bracket by $[x, y] = z$ and declaring that z is central.
 - (a) Show that this defines a Lie algebra, isomorphic to the space of matrices over F of the form $\begin{pmatrix} 0 & a & c \\ 0 & 0 & b \\ 0 & 0 & 0 \end{pmatrix}$ with the Lie bracket defined by the usual matrix commutator.
 - (b) Show that this is a 2-step nilpotent Lie algebra (i.e. the second term of the lower central series vanishes).