

**Math 100A – WORKSHEET 5**  
**THE CHAIN RULE**

1. THE CHAIN RULE

- (1) We know  $\frac{d}{dy} \sin y = \cos y$ .
- (a) ★ Expand  $\sin(y + h)$  to linear order in  $h$ . Write down the linear approximation to  $\sin y$  about  $y = a$ .
- (b) ★★ Now let  $F(x) = \sin(3x)$ . Expand  $F(x + h)$  to linear order in  $h$ . What is the derivative of  $\sin 3x$ ?

|  |
|--|
| <b>Fact.</b> $(f(g(x)))' = f'(g(x))g'(x)$ or $\frac{d}{dx}(f(g(x))) = \frac{df}{dg} \cdot \frac{dg}{dx}$ . |
|--|

- (2) Write each function as a composition and differentiate
- (a) ★  $e^{3x}$

(b) ★  $\sqrt{2x + 1}$

(c) (Final, 2015) ★  $\sin(x^2)$

(d) ★  $(7x + \cos x)^n$ .

- (3) (Final, 2012) ★★ Let  $f(x) = g(2 \sin x)$  where  $g'(\sqrt{2}) = \sqrt{2}$ . Find  $f'(\frac{\pi}{4})$ .

- (4) Differentiate  
 (a)  $\star 7x + \cos(x^n)$

(b)  $\star e^{\sqrt{\cos x}}$

(c)  $\star$  (Final 2012)  $e^{(\sin x)^2}$

- (5)  $\star\star$  Suppose  $f, g$  are differentiable functions with  $f(g(x)) = x^3$ . Suppose that  $f'(g(4)) = 5$ . Find  $g'(4)$ .

## 2. LOGARITHMIC DIFFERENTIATION

$$\log_b(b^x) = b^{\log_b x} = x$$

$$\log_b(xy) = \log_b x + \log_b y$$

$$\log_b(x^y) = y \log_b x$$

$$\log_b \frac{1}{x} = -\log_b x$$

$$\mathbf{Fact.} \quad \frac{d}{dx} \log x = \frac{1}{x}$$

(6)  $\star \log(e^{10}) =$

$\log(2^{100}) =$

(in terms of  $\log 2$ )

- (7)  $\star$  Differentiate

(a)  $\frac{d(\log(ax))}{dx} =$

$\frac{d}{dt} \log(t^2 + 3t) =$

(b)  $\star \frac{d}{dx} x^2 \log(1 + x^2) =$

$\frac{d}{dr} \frac{1}{\log(2 + \sin r)} =$

- (8) \*\* (Logarithmic differentiation) Use  $\log(fg) = \log f + \log g$  to differentiate  $y = (x^2 + 1) \cdot \sin x \cdot \frac{1}{\sqrt{x^3+3}} \cdot e^{\cos x}$ .

- (9) Differentiate using  $f' = f \times (\log f)'$

(a)  $x^n$

(b)  $x^x$

(c) \*\*  $(\log x)^{\cos x}$

- (d) (Final, 2014)  $x$  Let  $y = x^{\log x}$ . Find  $\frac{dy}{dx}$  in terms of  $x$  only.

### 3. MORE PROBLEMS

- (10) **\*\***Let  $f(x) = g(x)^{h(x)}$ . Find a formula for  $f'$  in terms of  $g'$  and  $h'$ .
- (11) Let  $f(\theta) = \sin^2 \theta + \cos^2 \theta$ . Find  $\frac{df}{d\theta}$  without using trigonometric identities. Evaluate  $f(0)$  and conclude that  $\sin^2 \theta + \cos^2 \theta = 1$  for all  $\theta$ .
- (12) (“Inverse function rule”) suppose  $f(g(x)) = x$  for all  $x$ .
- (a) Show that  $f'(g(x)) = \frac{1}{g'(x)}$ .
- (b) Suppose  $g(x) = e^x$ ,  $f(y) = \log y$ . Show that  $f(g(x)) = x$  and conclude that  $(\log y)' = \frac{1}{y}$ .
- (c) Suppose  $g(\theta) = \sin \theta$ ,  $f(x) = \arcsin x$  so that  $f(g(\theta)) = \theta$ . Show that  $f'(x) = \frac{1}{\sqrt{1-x^2}}$ .
- (13) (Final, 2015) **\*\*** Let  $xy^2 + x^2y = 2$ . Find  $\frac{dy}{dx}$  at the point  $(1, 1)$ .