

Math 257/316, Midterm 1, Section 202

15 February 2012

**Instructions.** The duration of the exam is 55 minutes. Answer all questions. Calculators are not allowed.

Maximum score 80.

1. Consider the second order differential equation:

$$Ly = 9x^2(1-x)y'' + 9xy' - y = 0 \quad (1)$$

- (a) Classify the points  $0 \leq x < \infty$  as ordinary points, regular singular points, or irregular singular points.

[10 marks]

- (b) If you were given  $y(-1) = 2$  and  $y'(-1) = 4$ , what form of series expansion would you assume (**Do not** determine the expansion coefficients of this series)? What would be the minimal radius of convergence of this series?

[5 marks]

- (c) If you required a solution about  $x = 1$ , what form of series expansion would you assume (**Do not** determine the expansion coefficients of this series)? What is the behavior of the solution as  $x \rightarrow 1$ ? What would be the minimal radius of convergence of this series?

[5 marks]

- (d) Use the appropriate series expansion about the point  $x = 0$  to determine two independent solutions to (1). You only need to determine the first three non-zero terms in each case.

[30 marks]

2. Apply the method of separation of variables to determine the solution to the one dimensional heat equation with the following Mixed homogeneous boundary conditions:

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}, \quad 0 < x < \pi, \quad t > 0$$

$$\text{BC} : \frac{\partial u(0, t)}{\partial x} = 0 = u(\pi, t)$$

$$\text{IC} : u(x, 0) = x(\pi - x)$$

Hint: It may be useful to know that:

$$\frac{2}{\pi} \int_0^\pi x(\pi - x) \cos\left(\frac{2n+1}{2}x\right) dx = \frac{8}{\pi} \frac{4(-1)^n - (2n+1)\pi}{(2n+1)^3}$$

[30 marks]