

On Tensor Products of Polynomial Representations

Kevin Purbhoo and Stephanie van Willigenburg


Abstract. We determine the necessary and sufficient combinatorial conditions for which the tensor product of two irreducible polynomial representations of $GL(n, \mathbb{C})$ is isomorphic to another. As a consequence we discover families of Littlewood–Richardson coefficients that are non-zero, and a condition on Schur non-negativity.

1 Introduction

It is well known that the representation theory of $GL(n, \mathbb{C})$ is intimately connected to the combinatorics of partitions [8, Ch. 7, Appendix 2]. Before we address the main problem in this paper that concerns the representations of $GL(n, \mathbb{C})$, we will briefly review this connection.

Recall that a *partition* λ of a positive integer m , denoted $\lambda \vdash m$, is a list of positive integers $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_{\ell(\lambda)} > 0$ whose sum is m . We call m the *size* of λ , the λ_i the *parts* of λ and $\ell(\lambda)$ the *length* of λ . We also let $\lambda = 0$ be the unique partition of 0, called the *empty partition* of length 0. Every partition corresponds naturally to a (*Ferrers*) *diagram* of *shape* λ , which consists of an array of m boxes such that there are λ_i left justified boxes in row i , where the rows are read from top to bottom. By abuse of notation we also denote this diagram by λ . In the following example the boxes are denoted by \times .

Example 1.1



Moreover, given partitions λ, μ such that $\lambda_i \geq \mu_i$ for all $1 \leq i \leq \ell(\mu)$, if we consider the boxes of μ to be situated in the top left corner of λ , then we say that μ is a *subdiagram* of λ , and the *skew diagram* of shape λ/μ is the array of boxes contained in λ , but not in μ . Again we abuse notation and denote this skew diagram by λ/μ .

Received by the editors April 11, 2006; revised December 1, 2007.
 The authors were supported in part by the National Sciences and Engineering Research Council of Canada.
 AMS subject classification: Primary: 05E05; secondary: 05E10, 20C30.
 Keywords: polynomial representation, symmetric function, Littlewood–Richardson coefficient, Schur non-negative.
 ©Canadian Mathematical Society 2008.

Example 1.2

$$43211/21 = \begin{array}{cc} & \times \times \\ & \times \times \\ \times & \times \\ & \times \\ & \times \end{array}$$

Furthermore, given a (skew) diagram, we can fill the boxes with positive integers to form a *tableau* T , and if T contains $c_1(T)$ 1s, $c_2(T)$ 2s, \dots , then we say it has *content* $c(T) = c_1(T)c_2(T)\dots$. With this in mind we are able to state the connection between $GL(n, \mathbb{C})$ and partitions of n as follows.

The irreducible polynomial representations ϕ^λ of $GL(n, \mathbb{C})$ are indexed by partitions λ such that $\ell(\lambda) \leq n$ and given two irreducible polynomial representations of $GL(n, \mathbb{C})$, ϕ^μ and ϕ^ν , one has

$$\text{char}(\phi^\mu \otimes \phi^\nu) = \sum_{\ell(\lambda) \leq n} c_{\mu\nu}^\lambda \text{char } \phi^\lambda,$$

where $c_{\mu\nu}^\lambda$ is the number of tableaux T of shape λ/μ such that

- (i) the entries in the rows weakly increase from left to right;
- (ii) the entries in the columns strictly increase from top to bottom;
- (iii) $c(T) = \nu_1\nu_2\dots$;
- (iv) when we read the entries from right to left and top to bottom the number of i s we have read is always greater than or equal to the number of $(i + 1)$ s we have read.

This method for computing the $c_{\mu\nu}^\lambda$ is called the *Littlewood–Richardson rule*. As one might expect the $c_{\mu\nu}^\lambda$ are called *Littlewood–Richardson coefficients*. Observe that we could have equally well chosen conditions (i)–(iv) to read

- (i) the entries in the rows weakly increase from *right to left*;
- (ii) the entries in the columns strictly increase from *bottom to top*;
- (iii) $c(T) = \nu_1\nu_2\dots$;
- (iv) when we read the entries from *left to right and bottom to top* the number of i s we have read is always greater than or equal to the number of $(i + 1)$ s we have read.

For convenience we will call this the *reverse Littlewood–Richardson rule*.

Example 1.3 To illustrate both rules we now compute $c_{21,21}^{321}$. We will replace each box with the number it contains.

Using the Littlewood–Richardson rule we obtain $c_{21,21}^{321} = 2$ from the tableaux

$$\begin{array}{c} 1 \\ 2 \\ 1 \end{array} \quad \text{and} \quad \begin{array}{c} 1 \\ 1 \\ 2 \end{array}.$$

Meanwhile, using the reverse Littlewood–Richardson rule we also obtain $c_{21,21}^{321} = 2$

from the tableaux

$$\begin{array}{c} 1 \\ 2 \\ 1 \end{array} \quad \text{and} \quad \begin{array}{c} 2 \\ 1 \\ 1 \end{array}.$$

Another place where Littlewood–Richardson coefficients arise is in the algebra of symmetric functions, $\Lambda = \bigoplus_{m \geq 0} \Lambda^m$, which is a subalgebra of $\mathbb{Z}[[x_1, x_2, \dots]]$ that is invariant under the natural action of the symmetric group. Each Λ^m is spanned by $\{s_\lambda\}_{\lambda \vdash m}$, where $s_0 := 1$ and

$$(1.1) \quad s_\lambda := \sum_T x^T.$$

The sum is over all tableaux T that satisfy conditions (i) and (ii) of the Littlewood–Richardson rule and $x^T := \prod_i x_i^{c_i(T)}$. For partitions λ, μ, ν the structure coefficients of these *Schur functions* satisfy

$$s_\mu s_\nu = \sum_\lambda c_{\mu\nu}^\lambda s_\lambda,$$

where the $c_{\mu\nu}^\lambda$ are again Littlewood–Richardson coefficients.

Similarly we can define the algebra of symmetric polynomials on n variables by setting $x_{n+1} = x_{n+2} = \dots = 0$ above and working with *Schur polynomials* $s_\lambda(x_1, \dots, x_n)$. Observe that by Definition (1.1) if $\ell(\lambda) > n$, then $s_\lambda(x_1, \dots, x_n) = 0$. The motivation for restricting to n variables is that the irreducible representations of $\text{GL}(n, \mathbb{C})$ can be indexed such that

$$(1.2) \quad \text{char } \phi^\lambda = s_\lambda(x_1, \dots, x_n).$$

See [2, 8] for further details.

2 Identical Tensor Products

We now begin to address the main problem of the paper, that is, to determine for which partitions λ, μ, ν, ρ we have

$$(2.1) \quad \phi^\lambda \otimes \phi^\mu \cong \phi^\nu \otimes \phi^\rho$$

for irreducible polynomial representations of $\text{GL}(n, \mathbb{C})$.

For ease of notation, we assume n is fixed throughout the remainder of the paper. Additionally, since $s_\lambda(x_1, \dots, x_n) = 0$ for $\ell(\lambda) > n$, we assume that all partitions have at most n parts. We extend our partitions to exactly n parts by appending a string of $n - \ell(\lambda)$ 0s. For example, if $n = 4$, then $\lambda = 32$ becomes $\lambda = 3200$.

We now define an operation on diagrams that will be useful later.

Definition 2.1 Given partitions λ and μ and an integer s such that $0 \leq s \leq n - 1$, the s -cut of λ and μ is the partition whose parts are


$$\lambda_1 + \mu_1, \quad \lambda_2 + \mu_2, \quad \dots, \quad \lambda_s + \mu_s, \\ \lambda_{s+1} + \mu_n, \quad \lambda_{s+2} + \mu_{n-1}, \quad \dots, \quad \lambda_{n-1} + \mu_{s+2}, \quad \lambda_n + \mu_{s+1},$$

listed in weakly decreasing order.

Remark 2.1. Diagrammatically we can think of the s -cut of λ and μ as

- (i) aligning the top rows of λ and μ , then
- (ii) cutting the diagrams λ and μ between the s and $s + 1$ rows,
- (iii) taking the rows of μ (or λ) below the cut and rotating them by 180° ,
- (iv) appending the newly aligned rows and sorting into weakly decreasing row length to make a diagram.

Example 2.2 If $n = 6$, then the 2-cut of 432110 and 543200 is 973321. This example can be viewed diagrammatically as the following.



It transpires that the s -cut of λ and μ yields a condition on Littlewood–Richardson coefficients.

Lemma 2.3 If λ , μ , and s are as in Definition 2.1 and κ is the s -cut of λ and μ then $c_{\lambda\mu}^\kappa > 0$.

Proof Observe that since the Littlewood–Richardson and the reverse Littlewood–Richardson rule yield the same coefficients, there must be a bijection ψ between the tableaux generated by each. This bijection will play a key role in the proof.

Consider creating a tableau T of shape κ/λ , where $\kappa_i = \lambda_i + \mu_i$ for $1 \leq i \leq s$, that will contribute towards the coefficient $c_{\lambda\mu}^\kappa$. If we use the Littlewood–Richardson rule, then it is clear that for $1 \leq i \leq s$ we must fill the boxes of the i -th row with the μ_i i s. Now all that remains for us to do is to fill the remaining boxes of T with μ_{s+1} $(s + 1)$ s, \dots , μ_n n s. To do this we create a tableau T' of shape $\kappa_{s+1} \cdots \kappa_n / \lambda_{s+1} \cdots \lambda_n = \kappa / \kappa_1 \cdots \kappa_s \lambda_{s+1} \cdots \lambda_n$ that will contribute towards the coefficient $c_{\alpha\beta}^\gamma$ where $\alpha = \lambda_{s+1} \cdots \lambda_n$, $\beta = \mu_{s+1} \cdots \mu_n$ and $\gamma = \kappa_{s+1} \cdots \kappa_n$. We do this as follows.

Fill the box at the bottom of each column from left to right with μ_{s+1} 1 s. Then repeat on the remaining boxes with the μ_{s+2} 2 s. Iterate this procedure until the boxes are full. Observe by the reverse Littlewood–Richardson rule that this filling contributes 1 to the coefficient $c_{\alpha\beta}^\gamma$. Now using ψ , create a tableau T'' of the same shape that satisfies the Littlewood–Richardson rule and increase each entry by s , forming a tableau T''' . Placing the entries of T''' in the naturally corresponding boxes of T we see we have a tableau that contributes 1 to the coefficient $c_{\lambda\mu}^\kappa$ by the Littlewood–Richardson rule and indeed $c_{\lambda\mu}^\kappa > 0$. ■

Definition 2.4 If λ , μ , and s are as in Definition 2.1, then the s -poset of λ and μ is the set of all partitions κ such that

- (i) $c_{\lambda\mu}^\kappa > 0$,
- (ii) $\kappa_i = \lambda_i + \mu_i$ for all $1 \leq i \leq s$,

which are ordered lexicographically, that is, $\kappa > \kappa'$ if and only if there exists some i , where $1 \leq i \leq n$, such that $\kappa_1 = \kappa'_1, \dots, \kappa_{i-1} = \kappa'_{i-1}$ and $\kappa_i > \kappa'_i$.

Lemma 2.5 *If λ, μ , and s are as in Definition 2.1, then the s -cut of λ and μ is the unique minimal element in the s -poset of λ and μ .*

Proof Let ξ be any element in the s -poset of λ and μ and let U be any tableau that will contribute towards the coefficient $c_{\lambda\mu}^\xi$ via the Littlewood–Richardson rule. As in the proof of Lemma 2.3, it is clear that for $1 \leq j \leq s$ we have that j appears in every box of row j . Now consider the natural subtableau of shape $\xi_{s+1} \cdots \xi_n / \lambda_{s+1} \cdots \lambda_n$, which we denote by \bar{U} . Note that if we subtract s from every entry in \bar{U} , then we obtain a tableau that contributes towards $c_{(\lambda_{s+1} \cdots \lambda_n)(\mu_{s+1} \cdots \mu_n)}^{(\xi_{s+1} \cdots \xi_n)}$ via the Littlewood–Richardson rule. If we then apply the bijection ψ to rearrange these new entries, we obtain a tableau U' that contributes towards $c_{(\lambda_{s+1} \cdots \lambda_n)(\mu_{s+1} \cdots \mu_n)}^{(\xi_{s+1} \cdots \xi_n)}$ via the reverse Littlewood–Richardson rule.

Now let κ be the s -cut of λ and μ . Let T and T' be the tableaux constructed in the proof of Lemma 2.3. Recall that T contributes towards the coefficient $c_{\lambda\mu}^\kappa$ via the Littlewood–Richardson rule, and that T' contributes towards $c_{(\lambda_{s+1} \cdots \lambda_n)(\mu_{s+1} \cdots \mu_n)}^{(\kappa_{s+1} \cdots \kappa_n)}$ via the reverse Littlewood–Richardson rule.

We now consider transforming T' into U' as follows. Since T' and U' both have content μ , we can map the boxes of T' bijectively to the boxes of U' such that the k -th box containing i from the left in T' maps to the k -th box containing i from the left in U' . This bijection factors as follows. First move each box in T' horizontally, so that it is in the same column as the corresponding box in U' . Then move each box vertically to form U' . By the construction of T' the entries are as left justified and low as possible, and so this transformation necessarily moves each box rightwards and upwards. It follows that κ , the shape of T' , is lexicographically less than or equal to ξ , the shape of U' , and we are done. ■

Recall that λ_n is the number of columns of length n in the diagram λ , and thus $(\lambda_n)^n$ is a subdiagram of λ . Define $\lambda^- := \lambda / (\lambda_n)^n$. Notice that λ^- is a Ferrers diagram, with at most $n - 1$ rows, and the number of columns of length $n - 1$ is λ_{n-1}^- . We therefore define $\lambda^{--} := \lambda^- / (\lambda_{n-1}^-)^{n-1}$. Notice that by (1.1) we have the factorization

$$(2.2) \quad s_\lambda(x_1, \dots, x_n) = (x_1 \cdots x_n)^{\lambda_n} s_{\lambda^-}(x_1, \dots, x_n),$$

and moreover $x_1 \cdots x_n$ does not divide $s_{\lambda^-}(x_1, \dots, x_n)$.

Theorem 2.6 *$\phi^\lambda \otimes \phi^\mu \cong \phi^\nu \otimes \phi^\rho$ as representations of $GL(n)$ if and only if $\lambda_n + \mu_n = \nu_n + \rho_n$ and $\{\lambda^-, \mu^-\} = \{\nu^-, \rho^-\}$ as multisets.*

An alternative proof, previously unknown to the authors, appears in [6].

Proof We will show that

$$(2.3) \quad s_\lambda(x_1, \dots, x_n) s_\mu(x_1, \dots, x_n) = s_\nu(x_1, \dots, x_n) s_\rho(x_1, \dots, x_n)$$

if and only if $\lambda_n + \mu_n = \nu_n + \rho_n$ and $\{\lambda^-, \mu^-\} = \{\nu^-, \rho^-\}$. The theorem then follows, using (1.2).

One direction is immediate. Suppose $\lambda_n + \mu_n = \nu_n + \rho_n$ and $\{\lambda^-, \mu^-\} = \{\nu^-, \rho^-\}$, then by (2.2) we have

$$\begin{aligned} s_\lambda(x_1, \dots, x_n) s_\mu(x_1, \dots, x_n) &= (x_1 \cdots x_n)^{\lambda_n + \mu_n} s_{\lambda^-}(x_1, \dots, x_n) s_{\mu^-}(x_1, \dots, x_n) \\ &= (x_1 \cdots x_n)^{\nu_n + \rho_n} s_{\nu^-}(x_1, \dots, x_n) s_{\rho^-}(x_1, \dots, x_n) \\ &= s_\nu(x_1, \dots, x_n) s_\rho(x_1, \dots, x_n). \end{aligned}$$

For the opposite direction, assume that (2.3) holds. We first show that $\lambda_n + \mu_n = \nu_n + \rho_n$. If they were not equal, say $\lambda_n + \mu_n > \nu_n + \rho_n$, then by (2.2), we would have

$$\begin{aligned} (x_1 \cdots x_n)^{\lambda_n + \mu_n - \nu_n - \rho_n} s_{\lambda^-}(x_1, \dots, x_n) s_{\mu^-}(x_1, \dots, x_n) \\ = s_{\nu^-}(x_1, \dots, x_n) s_{\rho^-}(x_1, \dots, x_n), \end{aligned}$$

which is impossible since $x_1 \cdots x_n$ does not divide the right-hand side. Similarly we cannot have $\lambda_n + \mu_n < \nu_n + \rho_n$. Thus, we see furthermore that

$$(2.4) \quad s_{\lambda^-}(x_1, \dots, x_n) s_{\mu^-}(x_1, \dots, x_n) = s_{\nu^-}(x_1, \dots, x_n) s_{\rho^-}(x_1, \dots, x_n).$$

Let $S(n)$ be the assertion that the equation (2.4) holds only if $\{\lambda^-, \mu^-\} = \{\nu^-, \rho^-\}$. To complete the proof of the theorem, it remains to show that $S(n)$ is true for all n . We prove this by induction.

The base case $n = 1$ is trivial, since each of $\lambda^-, \mu^-, \nu^-, \rho^-$ is necessarily the empty partition.

Now assume that $S(1), \dots, S(n-1)$ are true. In particular this assumption implies that the theorem holds for smaller values of n . Furthermore, assume that (2.4) holds. Let

$$a := \lambda_{n-1}^- \quad b := \mu_{n-1}^- \quad c := \nu_{n-1}^- \quad d := \rho_{n-1}^-.$$

Since (2.4) implies

$$s_{\lambda^-}(x_1, \dots, x_{n-1}) s_{\mu^-}(x_1, \dots, x_{n-1}) = s_{\nu^-}(x_1, \dots, x_{n-1}) s_{\rho^-}(x_1, \dots, x_{n-1}),$$

by our inductive hypothesis we must have

$$a + b = c + d \quad \text{and} \quad \{\lambda^{--}, \mu^{--}\} = \{\nu^{--}, \rho^{--}\}.$$

Assume without loss of generality that $\lambda^{--} = \nu^{--} =: \alpha$ and $\mu^{--} = \rho^{--} =: \beta$. To show that $\{\lambda^-, \mu^-\} = \{\nu^-, \rho^-\}$, we need to check that $a = c$ and $b = d$, or that $a = d, b = c$, and $\alpha = \beta$.

To show this we note that if (2.4) holds, then for all $s, 0 \leq s \leq n - 1$, the s -poset of λ^- and μ^- , must be the same as the s -poset of ν^- and ρ^- . Thus by Lemma 2.5, the s -cut of λ^- and μ^- must be the same as the s -cut of ν^- and ρ^- .

The s -cut of λ^- and μ^- has part sizes

$$\begin{aligned} a + b + \alpha_j + \beta_j & \quad 1 \leq j \leq s, \\ a + b + \alpha_{s+j} + \beta_{n-j+1} & \quad 2 \leq j \leq n - s - 1, \\ a + \alpha_{s+1}, & \\ b + \beta_{s+1}, & \end{aligned}$$

whereas the s -cut of ν^- and ρ^- has part sizes

$$\begin{aligned} a + b + \alpha_j + \beta_j & \quad 1 \leq j \leq s, \\ a + b + \alpha_{s+j} + \beta_{n-j+1} & \quad 2 \leq j \leq n - s - 1, \\ c + \alpha_{s+1}, & \\ d + \beta_{s+1}. & \end{aligned}$$

These lists must agree. Consequently we must have

$$a + \alpha_{s+1} = c + \alpha_{s+1} \quad \text{or} \quad a + \alpha_{s+1} = d + \beta_{s+1},$$

for all s . If, for any s , we are in the first situation, then $a = c$ and $b = d$ as desired. If not, then

$$a + \alpha_{s+1} = d + \beta_{s+1} \quad \text{and} \quad c + \alpha_{s+1} = b + \beta_{s+1}$$

for all $0 \leq s \leq n - 1$. In particular, since $\alpha_{n-1} = \beta_{n-1} = 0$, we have $a = d$ and $b = c$, ensuring $\alpha_j = \beta_j$ for $1 \leq j \leq n - 1$. ■

Example 2.7 If $n = 3$,

$$\begin{array}{ccc} \times & \times & \times & \times & \times & & & & \times & \times \\ \lambda = & \times & \times & \times & & & & & \times & \times \\ & \times & \times & & & & & & \times & \times \end{array} \quad \text{and} \quad \begin{array}{ccc} & & & & & & & & \times & \times \\ & & & & & & & & \times & \times \\ & & & & & & & & \times & \times \end{array}$$

then $\phi^\lambda \otimes \phi^\mu \cong \phi^\nu \otimes \phi^\rho$ if and only if $\{\nu, \rho\}$ is equal to one of

$$\{\lambda, \mu\}, \quad \left\{ \begin{array}{ccc} \times & \times & \times & \times & \times & \times & \times \\ \times & \times & & \times & \times & \times & \times \\ \times & & & \times & & & \end{array} \right\} \quad \text{or} \quad \left\{ \begin{array}{ccc} \times & \times & \times & \times & \times & \times & \times \\ \times & & & \times & \times & \times & \times \\ & & & \times & & & \times & \times \end{array} \right\}.$$

We consequently obtain a strict lower bound on n , in terms of the size of the partitions, to guarantee that (2.1) has only trivial solutions.

Corollary 2.8 Suppose m, m' are non-negative integers. If $n > \max\{m, m'\}$, then for any partitions $\lambda \vdash m$ and $\mu \vdash m'$, we have that

$$(2.5) \quad \phi^\lambda \otimes \phi^\mu \cong \phi^\nu \otimes \phi^\rho$$

has only the trivial solution $\{\nu, \rho\} = \{\lambda, \mu\}$. If $\min\{m, m'\} \geq 2$ and $n \leq \max\{m, m'\}$, then there exist $\lambda \vdash m$ and $\mu \vdash m'$ for which (2.5) has non-trivial solutions.

3 Schur Non-Negativity

Recently the question of Schur non-negativity has received much attention; see for example [4, 7]. The notion of Schur non-negativity is of interest as it arises in the study of algebraic geometry [1], quantum groups [3], and branching problems in representation theory [5].

One of the most basic Schur non-negativity questions is the following. Given partitions λ, μ, ν, ρ , when is the difference $s_\lambda s_\mu - s_\nu s_\rho$ a non-negative linear combination of Schur functions? Note that if $s_\lambda s_\mu - s_\nu s_\rho$ is Schur non-negative, then the same is certainly true of the corresponding expression in finitely many variables

$$s_\lambda(x_1, \dots, x_n) s_\mu(x_1, \dots, x_n) - s_\nu(x_1, \dots, x_n) s_\rho(x_1, \dots, x_n).$$

The following yields a test for failure of Schur non-negativity.

Corollary 3.1 For $0 \leq s \leq n - 1$, let $\kappa = \kappa_1 \cdots \kappa_n$ be the s -cut of λ and μ , and let $\xi = \xi_1 \cdots \xi_n$ be the s -cut of ν and ρ . Form the sequences

$$\sigma(s) := \kappa_1 \cdots \kappa_s \xi_{s+1} \cdots \xi_n \quad \text{and} \quad \tau(s) := \xi_1 \cdots \xi_s \kappa_{s+1} \cdots \kappa_n.$$

If there exists an s for which $\tau(s)$ is lexicographically greater than $\sigma(s)$, then

$$s_\lambda(x_1, \dots, x_n) s_\mu(x_1, \dots, x_n) - s_\nu(x_1, \dots, x_n) s_\rho(x_1, \dots, x_n)$$

is not Schur non-negative.

Proof Suppose the s -cut of λ and μ is not equal to the s -cut of ν and ρ , and let k be the first index in which they differ. If $k \leq s$, and $\xi_k > \kappa_k$, then by the Littlewood–Richardson rule, $c_{\lambda\mu}^\xi = 0$. On the other hand if $k > s$ and $\kappa_k > \xi_k$, then the same is true by Lemma 2.5. In either case, by Lemma 2.3, $c_{\nu\rho}^\xi > 0$, and thus

$$s_\lambda(x_1, \dots, x_n) s_\mu(x_1, \dots, x_n) - s_\nu(x_1, \dots, x_n) s_\rho(x_1, \dots, x_n)$$

is not Schur non-negative. ■

Example 3.2 Suppose $n = 3$, and

$$\lambda = 310, \quad \mu = 110, \quad \nu = 220, \quad \rho = 200.$$

Then $\sigma(0) = 222 < 321 = \tau(0)$. Thus we can conclude that $s_\lambda s_\mu - s_\nu s_\rho$ is not Schur non-negative. On the other hand, $\sigma(1) = 420 > 411 = \tau(1)$. Thus $s_\nu s_\rho - s_\lambda s_\mu$ is also not Schur non-negative.

Acknowledgements The authors are grateful to Matthew Morin for helpful discussions, Christopher Ryan for providing some data and Mark Skandera for suggesting the problem. They would also like to thank the referee for illuminating comments.

References

- [1] S. Fomin, W. Fulton, C. Li, and Y. Poon, *Eigenvalues, singular values, and Littlewood-Richardson coefficients*. Amer. J. Math. **127**(2005), no. 1, 101–127.
- [2] I. Macdonald, *Symmetric Functions and Hall Polynomials*. Second edition. Oxford University Press, New York, 1995.
- [3] A. Lascoux, B. Leclerc, and J.-Y. Thibon, *Ribbon tableaux, Hall-Littlewood functions, quantum affine algebras and unipotent varieties*. J. Math. Phys. **38**(1997), no. 2, 1041–1068.
- [4] T. Lam, A. Postnikov, and P. Pylyavskyy, *Schur positivity and Schur log-concavity*. Amer. J. Math. **129**(2007), no. 6, 1611–1622.
- [5] A. Okounkov, *Log-concavity of multiplicities with applications to characters of $U(\infty)$* . Adv. Math. **127**(1997), no. 2, 258–282.
- [6] C. Rajan, *Unique decomposition of tensor products of irreducible representations of simple algebraic groups*. Ann. of Math. **160**(2004), no. 2, 683–704.
- [7] B. Rhoades and M. Skandera, *Kazhdan-Lusztig immanants and products of matrix minors*. J. Algebra **304**(2006), no. 2, 793–811.
- [8] R. Stanley, *Enumerative Combinatorics*. Cambridge Studies in Advanced Mathematics 62, Cambridge University Press, Cambridge, 1999.

Department of Mathematics, University of British Columbia, Vancouver, BC, V6T 1Z2
e-mail: kevinp@math.ubc.ca
steph@math.ubc.ca