

Lecture 22: Interpreting D'Alembert's Solution in Space-Time: characteristics, regions of influence and domains of dependence

(Compiled 25 October 2012)

In this lecture we discuss the physical interpretation of the D'Alembert solution in terms of space-time plots. In particular we identify the left and right-moving *characteristics* as well as the domain of dependence of a given point (x_0, t_0) in space-time and the region of influence of a given initial value specified at the point $x_1, 0$. We discuss the evolution of a few simple pulses and track the regions in space-time that are carved out by the intersecting characteristics.

Key Concepts: The one dimensional Wave Equation; D'Alembert's Solution, Characteristics, Domain of Dependence, Region of Influence.

Reference Section: Boyce and Di Prima Section 10.7

22 Space-Time Interpretation of D'Alembert's Solution

In this lecture we discuss the interpretation of D'Alembert's solution

$$u(x, t) = \frac{1}{2} [u_0(x - ct) + u_0(x + ct)] + \frac{1}{2c} \int_{x-ct}^{x+ct} v_0(s) ds \quad (22.1)$$

to the one dimensional wave equation

$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2} \quad (22.2)$$

22.1 Characteristics

In the $x - t$ plane the lines

$$x - ct = x_0 \text{ and } x + ct = x_0 \quad (22.3)$$

are called the *characteristics* that emanate from the point $(x_0, 0)$ in space-time (see figure 1). Characteristics are the lines (or curves of more general hyperbolic problems) along which information is propagated by the equation. To interpret the characteristic lines in the $x - t$ plane, it is useful to rewrite the characteristic equations in the form

$$\begin{aligned} x - ct = x_0 &\Rightarrow t = \frac{1}{c}x - \frac{1}{c}x_0 \\ x + ct = x_0 &\Rightarrow t = -\frac{1}{c}x + \frac{1}{c}x_0 \end{aligned} \quad (22.4)$$





FIGURE 1. The characteristics that emanate from.

22.2 Region of Influence and Domain of Dependence

Region of influence: The lines $x + ct = x_1$ and $x - ct = x_1$ bound the region of influence of the function values at the initial point $(x_1, 0)$. Thus all the solution values $u(x, t)$ are influenced by the value at the point $(x_1, 0)$.

Domain of Dependence: The lines $x = x_0 - ct_0$ and $x = x_0 + ct_0$ that pass through the point (x_0, t_0) bound the domain of dependence. Thus the solution $u(x_0, t_0)$ depends on all the function values in the shaded region.

FIGURE 2. Space-time Region of Influence of the point $(x_1, 0)$ and Domain of Dependence of the point (x_0, t_0) , both of which can be determined from D'Alembert's Solution (22.1).

Example 22.1 A Rectangular pulse Pulse:

$$u(x, 0) = \begin{cases} 1 & |x| < 1 \\ 0 & |x| > 1 \end{cases} \quad (22.5)$$

$$u(x, t) = \frac{1}{2} [u_0(x - ct) + u_0(x + ct)] \quad (22.6)$$

Let $c = 1$.

$t = \frac{1}{2}$:

$$\begin{aligned} x_r - \frac{1}{2} &= 1 \quad \Rightarrow \quad x_r = \frac{3}{2} \quad x_R + \frac{1}{2} &= 1 \quad x_R = \frac{1}{2} \\ x_\ell - \frac{1}{2} &= -1 \quad \Rightarrow \quad x_\ell = -\frac{1}{2} \quad x_L + \frac{1}{2} = -1 \quad x_L = -\frac{3}{2} \end{aligned} \quad (22.7)$$

$t = 1$:

$$\begin{aligned} x_r - 1 &= 1 \quad \Rightarrow \quad x_r = 2 \quad x_R + 1 = 1 \quad \Rightarrow \quad x_R = 0 \\ x_\ell - 1 &= -1 \quad \Rightarrow \quad x_\ell = 0 \quad x_\ell + 1 = -1 \quad \Rightarrow \quad x_L = -2 \end{aligned} \quad (22.8)$$

$t = 2$:

$$\begin{aligned} x_r - 2 &= 1 \quad \Rightarrow \quad x_r = 3 \quad x_R + 2 = 1 \quad \Rightarrow \quad x_R = -1 \\ x_\ell - 2 &= -1 \quad \Rightarrow \quad x_\ell = 1 \quad x_L + 2 = -1 \quad \Rightarrow \quad x_L = -3 \end{aligned} \quad (22.9)$$




FIGURE 3. Top: Space-time representation of the regions in which the solution takes on different values for the rectangular pulse (22.5). Bottom: Cross sections of the solution $u(x, t)$ at times $t = 0$, $1/2c$, $1/c$, and $t > 1/c$